



Health and Welfare
Canada

Santé et Bien-être social
Canada

Canada

Health Services and Promotion Branch Working Paper

HEALTHY ENVIRONMENTS FOR CANADIANS

HSPB 88-12

Bruce M. Small and Associates Limited



HEALTHY ENVIRONMENTS FOR CANADIANS

HSPB 88-12

Bruce M. Small and Associates Limited

R.R. #1, Goodwood, Ontario LOC 1AO

December, 1987

Principal Consultant: Bruce M. Small, P. Eng.
Small and Associates

Principal Researcher: Wendy Priesnitz
Wendy Priesnitz and
Associates

Research Associate: Barbara J. Small
Small and Associates

Supported by Grant No. R-1301-504-5210
from Health and Welfare Canada.

- * This paper was prepared on contract for the Health Promotion Directorate/NHRDP Working Group on Priorities for Health Promotion/Disease Prevention Research. The views expressed herein are not necessarily those of the Department of National Health and Welfare.

Table of Contents

PART I: OVERVIEW	1
Foreword	1
Executive Summary	3
1. Synthesis of Literature Review	5
2. Gaps in Knowledge	8
3. Methodological Considerations	10
4. Suggested Indicators	12
5. Data Needs	14
6. Priority Questions for Research	15
7. Political and Social Considerations	17
8. Implications for Action	19
 PART II: ANALYSIS	 A1
1. Introduction	A1
2. Healthy Environment Issues	A4
2.1 By Setting	A4
2.1.1 At Home	A4
2.1.2 At School	A8
2.1.3 In Workplaces	A12
2.1.4 In Public Places	A19
2.2 By Human Characteristic & Situation	A23
2.2.1 Age	A23
2.2.2 Gender	A32
2.2.3 Family Type	A36
2.2.4 Race & Culture	A38
2.2.5 Illness & Handicap	A40
2.2.6 Orientation and Lifestyle	A46

.../continued

HEALTHY ENVIRONMENTS FOR CANADIANS
A Review for Health and Welfare Canada

Table of Contents (continued)

2.3 By Environmental Resource	A50
2.3.1 Safety	A50
2.3.2 Clean Air	A51
2.3.3 Food and Water	A56
2.3.4 Physical Factors	A60
2.3.5 Social Conditions	A64
2.3.6 Employment and Money	A69
2.3.7 Personal and Social Support	A72
2.3.8 Health Care and Social Services	A74
3. Bringing About Healthier Environments	A79
3.1 Ending Prejudice and Oppression in Canada	A79
3.2 Creating a New Vision of Environmental Choice	A81
3.3 Fostering Technology for Assessing Our Environment	A82
3.4 Accommodating Diversity Through Participation in Design	A84
3.5 Becoming World Experts in Designing Healthy Environments	A85
 PART III: ANNOTATED BIBLIOGRAPHY	 B1 - B244
 PART IV: NAME AND SUBJECT INDEXES	 C1
Name Index	C1-C10
Subject Index	C11-C84

PART I: OVERVIEW**Foreword**

This review and analysis was prepared for the Health Promotion Directorate and the NHRDP Working Group on Priorities for Health Promotion/Disease Prevention Research, of Health and Welfare Canada, under Articles of Agreement for Consulting and Professional Services dated May 4, 1987 between Her Majesty the Queen in right of Canada represented by the Minister of National Health and Welfare and Bruce M. Small and Associates Limited.

This document constitutes a comprehensive review of scientific and other literature, and of other research sources available to the contractor, in the area of healthy environments, addressing the question 'What is a healthy environment for people in Canada?'

Part I: Overview represents the contractor's interpretations of the facts available and the contractor's recommendations to the Government of Canada for making healthy environments accessible to all Canadians, consistent with the Minister of Health's 'Health for All' initiative.

Part II: Analysis represents the contractor's organization of the knowledge found in the literature and research review. No claim of completeness is made for each area of discussion, as the knowledge summarized is limited by the amount of research uncovered.

Part III: Annotated Bibliography lists all relevant literature and research contacts uncovered during the review. Most of the entries are annotated with a summary or abstract of important information contained in the reference.

Part IV: Name and Subject Indexes contains a full alphabetical listing of all principal authors referenced in Parts I and II, and all authors included in Part III. A detailed subject index is provided to all sections, so that readers can access material relevant to their field of interest directly. Page numbers in the index distinguish between Parts I, II and III.

The contractor takes full responsibility for the logic and accuracy of this review. The reader is cautioned, however, that Parts II and III make considerable reference to conclusions drawn by others in research, which we have carefully and critically selected, but not independently verified.

The views expressed herein are not necessarily those of the Department of National Health and Welfare.

Executive Summary

Physical and social environments have an important effect on the health and wellbeing of people in Canada. Although this connection has been known and documented for many years, the knowledge does not appear to have been widely applied within Canada's health care delivery systems and in the design of physical settings and social programs and institutions in Canada.

There are major health and safety issues to be addressed in virtually every physical setting in Canada - in homes, in schools, in the workplace, and in public places, including the natural environment. Creating building designs, operational procedures, building and maintenance materials, and other products that do not lead to polluted indoor air is a common priority in all settings, that would appear to lead to the greatest immediate health benefit.

Many people in Canada are stuck in physical settings and social conditions which are unhealthy for them. Poverty and unemployment, for example, are major contributors to ill health. In many cases, either overt or subtle forms of discrimination (devaluing of one group of people by another) are involved in confining an individual to unhealthy environments. Prejudices based on age, gender, family type, race and culture, illness or handicap, and orientation and lifestyle are still common in Canada. If healthy environments are to be available for everyone in Canada, we must begin on every level to treat each person, no matter what their characteristics, as a person of value. In particular, achieving equal status for women in Canada is essential.

Many of our common environments ignore the true range of physical and cultural differences among people in Canada, and are less accessible than they could be. Some of our specialized environments recognize our physical differences, but forget the rest of our human characteristics and needs, creating environments which may be safe, or accessible, but which are psychologically or socially unhealthy. A new vision of environmental choice and greater individual control over environment is needed.

Creating better environmental designs, eliminating personal and structural prejudices, and applying new knowledge about environmental health within Canada's health care systems, all require a much more detailed understanding of the true diversity of Canada's people, their social and psychological characteristics, their needs and goals, and their physical differences in vulnerability to different environmental stressors. Direct participation of individuals who will be affected by any physical design, social program, or social institution is required.

Research and development of emerging technologies which give detailed, quantified information on the effects of different environments on health will provide the scientific basis on which to proceed to design healthier environments in Canada. Creating healthy environments in Canada, and producing a new generation of safer, cleaner materials and products will enhance Canada's industrial and scientific reputation, and distinguish Canada's products and services from those of other countries in world markets.

1. Synthesis of the Literature Review

The following is a brief synthesis of the major conclusions arising from the literature and from ideas obtained by research contacts. A detailed analysis of research and other information on healthy environments is presented in Part II: "Analysis".

a) our physical environment can affect our health

Dangers in the physical environment contribute significantly to the risk of injury, ill health, or death, for many people in Canada. The physical environment contains some health or safety risks in virtually every Canadian setting: homes, schools, workplaces, and public environments, including natural environments.

b) indoor air pollution is a significant component of the risk

The contamination of indoor air by numerous chemical emissions represents a significant component of the overall risk to health posed by physical environmental factors. Indoor emission sources include industrial processes, building materials, furnishings, combustion devices, maintenance products, pest control substances, personal care or pleasure products and human metabolism.

c) pollutants in food, water, and outdoor air are also stressors

Trace-level contamination of food, drinking water and outdoor air, with a wide array of pollutants of varying toxicities, likewise presents a general chemical stress to which virtually everyone in Canada must adapt.

d) social and economic conditions can definitely affect health

Unemployment, poverty and financial stress, work pressures, family problems, prejudice and oppression, isolation, and lack of control over one's life are all examples of socio-economic conditions which can have an adverse effect on physical and mental health. Conditions previously thought of as merely unfortunate (at least by the fortunate) can be highly stressful and clearly have more health cost than has been commonly acknowledged.

e) there is a wide range of vulnerability in the population

Each individual is unique in his or her reactivity to physical environmental factors. The population contains a wide variety of people, ranging from those who can go about their lives with little or no effect from their immediate physical environment, to individuals who are severely impaired in their day-to-day functioning by one or more specific environmental factors. Any one individual may have a unique array of sensitivities - he or she may be quite insensitive to one factor while being very sensitive to another.

f) limits to adaption to changing environmental conditions

Some people appear to be suffering, at least temporarily, from an inability to adapt to changing conditions, particularly increased pollution load. A healthy environment and lifestyle appears to be one in which an individual has sufficient control to limit the amount of adaptation required at any one time, and to shape the environment to accommodate his or her inherent nature, rather than the other way around.

g) environmental health issues are different for each of us

Environmental health issues vary widely according to different human characteristics and situations. People of different ages, gender, family or household types, races or cultures, health status or mobility, sexual orientation and lifestyle all encounter variations of general environmental health issues, which are unique to their situation. Each different physical setting and each different social and economic condition brings with it new viewpoints and priorities in environmental health. Understanding this diversity of issues is an important key to achieving health for all.

h) the paradox in environmental design

Many environments (e.g. public spaces, offices, schools) do not accommodate well the full range of human characteristics. People who are handicapped, less mobile, or chemically hypersensitive, for example, have been lumped in the category of 'special needs' and often excluded from consideration, even though making environments accessible to them might also benefit the rest of the population, both physically, by many of the health and safety features required, and socially, by their presence. Where special environments have been created, they accommodate well certain unique needs of such individuals, but often ignore the rest of their human characteristics and isolate them from others.

**i) social conditions and being devalued trap
people in unhealthy environments**

Many people in Canada become locked into situations where they are exposed to unhealthy environments. Certain groups are subject to environmental hazards more than others. Devaluing or undervaluing people from certain groups (e.g. women, blue collar workers, handicapped people) can lead to widespread disparities in both social and physical environmental conditions.

j) prejudice is itself unhealthy

It is now clear that victims of prejudice and oppression suffer the direct physical and emotional health effects of such an unhealthy social atmosphere, in addition to the indirect environmental consequences arising from the discrimination they experience. The 'inferiorized' often internalize some of the attitudes toward them and can be subject to a spiral of decrea-

sing self-esteem and the many other problems that develop from that. Many people are undervalued ostensibly because of one characteristic for which they are singled out, which is bad enough, but the nature of prejudice is such that the rest of their humanity is also rejected in the process.

k) social and personal support is important

The extent of an individual's personal and social support network is an important determinant of health. Isolation increases health risk and also guarantees that a person's needs, and value, remain invisible to others.

l) more personal control and participation can reduce hazards

Many people in Canada voice a hopelessness about reducing their health risks from environmental exposures, because they lack sufficient control over their personal and social surroundings to effect the changes they want. Participation of the people of Canada in the creation or modification of their physical and social surroundings is a feasible, and important, step in creating healthier environments. It is critical to introducing into the design process sufficiently detailed information about the full diversity of the people being served. A great deal of knowledge is already available about user participation and need only be implemented.

e) new ways of measuring effects of environment change our views

Advances in measuring pollutants and the ways we are affected by them, are revolutionizing the field of environmental health. Health and performance effects are becoming apparent at levels of exposure previously considered safe, and questions are being posed about the unknown and possibly synergistic effects of mixtures of traces levels of different pollutants. Emerging new technologies provide the basis for scientific selection of healthier environments.

2. Gaps in Knowledge

It would be much easier to determine which environments are healthy, and which are not, if a number of gaps in our knowledge base were filled. These include:

a) detailed health effects on individuals

With few exceptions, we do not now know the actual effects of many environments on individual human beings. Much of the research designed to help decide what is healthy and what is not, is of an epidemiological or other statistical nature, or involves the effects of environment on animals rather than humans. While helpful in determining general environmental standards and identifying major hazards, this level of detail is not sufficient to make truly competent decisions about healthy environments for specific individual human beings, which after all is the basic question each individual in Canada faces, and that health care professionals must also deal with. Imagine a time when each person in Canada could have a profile of their own individual adaptability to different kinds of environments.

b) population profiles of environmental sensitivities

We do not know at the moment, with any degree of accuracy, how sensitivity to environmental insults, either physical or social, varies across the population. Without this, we cannot make competent decisions about design standards for physical, social, or economic environments in Canada. One profile urgently needed is that of people's degree of reactivity to common mixtures of indoor air pollution.

c) profiles of the extent of diversity in Canada

A true picture of the extent of diversity of human characteristics and situations among people in Canada has yet to be painted. Statistics are currently gathered under certain basic characteristics (e.g. gender, marital status, age, diseases, income, education, employment, etc) and usually analyzed separately. But each individual has many diverse characteristics, and a richer picture of these characteristics, and their various combinations, is more useful for physical, social, and economic design decisions. Categorizing people by single characteristics pulls us toward thinking about stereotypes rather than about accurate descriptions of people. How many young, black single women with an interest in learning more about computers are there in Canada, and how can we help them create a healthy environment for themselves? How would our approach differ if we knew more about them?

d) how physical, chemical and social stress affects our bodies and minds

We lack detailed and verified knowledge of the mechanism by which various stresses affect human health adversely. For example, it is known that low-level chemical exposures can trigger many measurable symptoms in sensitive individuals. There is no consensus on the mechanism by which such individuals are affected. Immunology research is teaching us that many body parameters are affected by stress of all kinds, but more work is needed to unravel the remaining details.

e) how to help people heal, who are victims of environmental stress

There is limited experience in helping individuals who have been adversely affected by their environment, to heal the effects. Clinical work is being done with people who become hypersensitive to minute chemical exposures, but there is no consensus yet on appropriate therapy, other than reducing the stress load on the individual. This is a serious gap in light of the fact that many physical and social factors in present-day Canadian environments pose considerable health risks.

f) knowledge of the effects of cleaner environments

We are no longer in the position, in Canada, of having comparison populations who can give us a clear picture of what health could be in a much less polluted and much less stressful environment. Most people in Canada are exposed to complex mixtures of indoor and outdoor contaminants, albeit at trace levels. While lifestyles vary considerably, most Canadians also experience a complex environment of social and economic stress. Competent decisions about the value of less polluted or less stressful environments will require good information about the effects of alternatives. To gather this knowledge, experimental situations will be required.

3. Methodological Considerations

Much of what is needed in the field of environmental health will involve the extension of technology and research methods well beyond the present state-of-the-art. It will involve tackling physical and social stressors without an overlay of hopelessness which presumes, without testing the limits of our ingenuity, that nothing can be done. Some examples of new methodologies needed are as follows:

a) protocols for measuring chemical hazards

It has become important to know, in far more detail than before, how products behave, physically and chemically. We require efficient and economical laboratory protocols for determining what substances are given off into our air and onto our skins, by the materials we handle and the environments we live in, from day to day.

b) protocols for measuring effects of environment

We need well-controlled, scientifically-verified procedures for obtaining accurate answers about the effects of specific environmental exposures on individuals. New and promising technology, such as computerized brainwave analysis and other developments described in this review, may provide the means of doing this.

c) protocols for responding to physical environmental complaints

We need better protocols for responding to situations in which individuals feel that their immediate environment is making them ill. While industrial situations involving high-level exposures have been well addressed, situations where exposure levels are below accepted standards often are not satisfactorily resolved. These will require better detective work, a greater understanding of the possible effect of low level exposures on sensitive individuals, and more direct involvement of people who are affected.

d) protocols for measuring the effects of alternative environments

New procedures, facilities and technology are needed to open up a new field which might best be described by the term 'experimental environmental health'. Sometimes the best way of determining the effect of an environmental stressor is to remove it from the environment, and to measure how we feel without it. This would open the way to using individuals as their own scientific controls, and, in combination with protocols for measuring the direct effects of environmental stressors, determining the precise role of a particular stressor on an individual's health and wellbeing.

d) how physical, chemical and social stress affects our bodies and minds

We lack detailed and verified knowledge of the mechanism by which various stresses affect human health adversely. For example, it is known that low-level chemical exposures can trigger many measurable symptoms in sensitive individuals. There is no consensus on the mechanism by which such individuals are affected. Immunology research is teaching us that many body parameters are affected by stress of all kinds, but more work is needed to unravel the remaining details.

e) how to help people heal, who are victims of environmental stress

There is limited experience in helping individuals who have been adversely affected by their environment, to heal the effects. Clinical work is being done with people who become hypersensitive to minute chemical exposures, but there is no consensus yet on appropriate therapy, other than reducing the stress load on the individual. This is a serious gap in light of the fact that many physical and social factors in present-day Canadian environments pose considerable health risks.

f) knowledge of the effects of cleaner environments

We are no longer in the position, in Canada, of having comparison populations who can give us a clear picture of what health could be in a much less polluted and much less stressful environment. Most people in Canada are exposed to complex mixtures of indoor and outdoor contaminants, albeit at trace levels. While lifestyles vary considerably, most Canadians also experience a complex environment of social and economic stress. Competent decisions about the value of less polluted or less stressful environments will require good information about the effects of alternatives. To gather this knowledge, experimental situations will be required.

3. Methodological Considerations

Much of what is needed in the field of environmental health will involve the extension of technology and research methods well beyond the present state-of-the-art. It will involve tackling physical and social stressors without an overlay of hopelessness which presumes, without testing the limits of our ingenuity, that nothing can be done. Some examples of new methodologies needed are as follows:

a) protocols for measuring chemical hazards

It has become important to know, in far more detail than before, how products behave, physically and chemically. We require efficient and economical laboratory protocols for determining what substances are given off into our air and onto our skins, by the materials we handle and the environments we live in, from day to day.

b) protocols for measuring effects of environment

We need well-controlled, scientifically-verified procedures for obtaining accurate answers about the effects of specific environmental exposures on individuals. New and promising technology, such as computerized brainwave analysis and other developments described in this review, may provide the means of doing this.

c) protocols for responding to physical environmental complaints

We need better protocols for responding to situations in which individuals feel that their immediate environment is making them ill. While industrial situations involving high-level exposures have been well addressed, situations where exposure levels are below accepted standards often are not satisfactorily resolved. These will require better detective work, a greater understanding of the possible effect of low level exposures on sensitive individuals, and more direct involvement of people who are affected.

d) protocols for measuring the effects of alternative environments

New procedures, facilities and technology are needed to open up a new field which might best be described by the term 'experimental environmental health'. Sometimes the best way of determining the effect of an environmental stressor is to remove it from the environment, and to measure how we feel without it. This would open the way to using individuals as their own scientific controls, and, in combination with protocols for measuring the direct effects of environmental stressors, determining the precise role of a particular stressor on an individual's health and wellbeing.

e) a means of interrupting prejudice and oppression

For years, we have relied on the idea that prejudice in Canada will naturally reduce with each new generation. Legislation has been used to guard against blatant violations of rights, and some focus has also been placed on acknowledging and welcoming multiculturalism in Canada. What has yet to be done well, is research to determine how we can deliberately reverse existing prejudices and eliminate oppressive institutional practices in a much shorter time frame. The key to doing so is making far more information available about the full humanity of devalued individuals.

f) methods for population-wide education about environmental health

Environment plays a far bigger role in health than has been previously accepted. Many people in Canada are both adversely affected by environment, and ignorant of the possibility that environmental factors may be contributing to their ill health. Those they consult about health often overlook environment as a factor. Since achieving healthy environments is a very individual affair, health for all cannot be achieved without widespread knowledge of the role of environment in health. Effective ways of spreading this knowledge must be chosen, tested, evaluated and implemented, if health for all is to be achieved.

g) methods for allowing greater individual environmental control

Both physically and socially, people have a better chance of maintaining their health and thriving in Canada, if they have control over their lives. Alternate living conditions, location, employment, education, and lifestyles are not available to those who are locked in, by lack of resources or by other oppressions that restrict their life choices. We need a more creative effort at every level (from building design to design of all our social and economic institutions) to allow individual diversity and control.

h) greater participation by individuals in physical and social design

A great deal of work has already been done on user participation in design. Methods appropriate to each new design situation need to be selected from the literature and put into practice.

4. Suggested Indicators

Canadian society is rich with indicators of environmental problems, in the form of specific physical health, mental health, and social problems that could be the result of unhealthy environments. At our present state of knowledge, however, it is difficult to link cause and effect. Enough practical experience has nonetheless been gained to guide the way in searching for environmental conditions which can and should be changed. Currently, we are often focusing solely on the indicator as a problem or disease in itself.

The first line of defence for understanding the connections between unhealthy environments and their effects, and therefore for finding reliable indicators, is direct contact with people who are experiencing ill health. In our zeal for obtaining objective and measurable indicators in the past, we have often ignored subjective information that provides clues which are both useful and necessary to carrying on the detective process. Once those clues have been processed and understood, there is an important role for scientifically determined indicators. The process requires both.

There are many problems in Canada that could have environmental factors as possible contributing causes. Rigorous investigation of these possible connections, including direct contact with individuals affected, will yield indicators which can be used with better confidence. Some possible health indicators are listed below along with the environmental connections suggested in the literature:

<u>environmental condition</u>	<u>possible indicators</u>
indoor air pollution	unexplained physical illness frequent medicinal drug use learning disabilities
outdoor air pollution	hospital admissions for respiratory illness mortality from respiratory and cardiovascular distress
contaminants in food & water	cancer incidence allergic reactions
sexism and oppression of women	female/male income disparity poverty among women sexual harrassment and assault battering of women
adultism	apathy among young people drug & alcohol abuse sexual and other child abuse youth suicide rate

unemployment	physical and mental illness poverty
prejudice	under-representation of groups in employment, educational and economic categories invisibility or closeting of certain characteristics
job stress	substance abuse mental health admissions unexplained physical illness
restricted information	unwanted pregnancy sexually transmitted diseases mortality from AIDS
undervaluing elderly or handicapped people	fear of aging loss of self-esteem inadequate health care poverty

For each type of environmental stressor, specific indicators are needed to signal potential problem levels, recognizing that this method will only alert us to the levels at which relatively large numbers of people may begin to be adversely affected, and will not be sufficient to warn of damage to those who are more sensitive than most to each factor.

For indoor air pollution, residential exposure guidelines have been developed and are being refined by a joint Federal/Provincial Advisory Committee on Environmental and Occupational Health, and contain some consideration of sensitive individuals. Environment Canada guidelines on outdoor pollutants provide some measure of problem levels, but do not yet address the problem of the more sensitive individuals.

Direct behaviour indicators for measuring our progress in reducing attitudes and actions which devalue other people, have yet to be systematically researched, refined and applied. We have not used a 'sexism scale', for example, to chart our progress in eliminating the oppression and undervaluing of women in Canada. Nor have we used a 'racism scale' or 'anti-gay scale' to measure the extent of prejudice that is still targeted against individual human beings who happen to have, among their diverse characteristics, specific origins, skin colour, sexual orientation, or family lifestyle that have been judged by others as undesirable, based on stereotypes.

We also need performance indicators, calibrated in clean, healthy environments, to be able to measure impairment of human beings due to environmental factors. The more ubiquitous the chemical contamination, the less we can call on present baselines as 'normal' performance.

5. Data Needs

It is sometimes appropriate to focus our full capabilities on a problem even if only a few people are affected (e.g. arranging a liver transplant for an ailing infant). However, the nature of the resources we allocate often depends on the magnitude of each problem we face.

We therefore need to know the number of people adversely affected by specific environments. For example, we need to know the number of children whose health and learning are actually now being impaired by indoor pollution in the schools. We need to know the number of industrial workers whose well-being has already been eroded by industrial chemical exposures. We need to know the number of women who are battered, the number of children who are abused, the number of people who are discriminated against, and so on.

Each area of study suggested in the previous sections will require its own type of data. Determining the list of 'numbers needed', however, is more appropriately left until basic, qualitative information about the problems we have discussed has been gathered and understood in the context of healthy environments. In this literature review, we do not pretend to have investigated environmental health problems in sufficient detail to more specific about data needs.

The next step after initial awareness of environmental health problems is more direct contact with individuals who have been affected. Even a count of the number of people who come forward with information, once invited, will be helpful in determining the minimum order of magnitude of each problem. The understanding gained from direct contact will help to determine exactly what, or who, should be counted.

6. Priority Questions for Research

General research needs have been described in the foregoing sections and in Part II: Analysis. The following list places a priority on certain subject areas over others:

a. experimental environmental health and treatment of environmental illness

Establishment of a field of research in which human beings are observed in alternative, cleaner, less stressful environments, and in which effective treatments are developed for people who are adversely affected by physical and social stressors;

b. pollution, health and performance:

Development of measurement devices and protocols for determining both obvious and subtle health and performance effects in individuals exposed to pollutants;

c. elimination of prejudice:

Development of practical ways of eliminating prejudice and discriminatory behaviour, individually and collectively;

d. adaptable physical design

Continued development of physical designs (e.g. buildings, public spaces), that accommodate a wider range of diversity;

e. health monitoring devices:

Development of devices that can be used by individuals at home, at school, or at work, for monitoring both environment and health conditions;

f. specific hazard investigations:

Further detailed investigation of newly recognized health hazards, e.g. airborne mycotoxins from household mould growth, and mixes of volatile organic chemicals in home and work environments;

g. safe product research:

Promotion of the design of safer, less-polluting products of all kinds; priority to building materials and interior finishings such as paints, and household combustion devices;

h. social environment research:

Investigation in more detail of the effects of various social environments on health;

i. internalized oppression:

Development of practical ways of addressing the effects of oppression in individuals who have been undervalued, and encouraging their full powers and capabilities;

j. alternative social structures:

Investigation of alternative income systems, social structures, institutions and legislation which will promote a society where all people with minority characteristics are fully valued as human beings, and not merely tolerated;

k. healthy lifestyles:

Investigation of alternative lifestyles or lifestyle factors which will contribute to better physical and mental health.

7. Political and Social Considerations

Sufficient new knowledge has accumulated over the last few years, concerning the connection between environmental factors and ill health, to have caused increased pressure on politicians, governments, organizations, employers, physicians, and health care institutions to provide or promote healthier environments in Canada.

Some of the important new political and social considerations arising from the present review are as follows:

a. 'Health For All' requires elimination of prejudice

One inescapable conclusion of the literature is that environments of prejudice and oppression contribute to ill health in many ways, both destroying self-esteem and locking some individuals into situations where they have less control than others, over income, lifestyle and surroundings, and hence their own health. The government cannot pretend to work toward 'Health For All' without taking an active role in achieving the complete elimination of various kinds of prejudice and oppression presently operative in Canada, including:

- sexism (oppression of women by men)
- ageism (devaluing of elderly people by younger people)
- adultism (devaluing of younger people by adults)
- racism (devaluing of one race or ethnic origin by another)
- prejudice against one religion by another
- gay oppression (prejudice on the basis of sexual orientation)
- devaluing of people who are unemployed
- devaluing of people in less advantaged economic positions
- devaluing of people who are handicapped
- devaluing of people who are physically or mentally ill

b. Healthy environments require individual freedom and control

If Canada is to move towards healthier environments for all, we must move in the direction of greater control by each person over his or her own surroundings and lifestyle. This has implications for building design, housing, income distribution, employment policy, income alternatives, human rights, health systems, pollution standards, and many other areas. The vision is one of environmental choice, respecting the vast diversity of human characteristics and situations in Canada. It leads to more individual empowerment, more diversity, and, ultimately, healthier and more effective human resources in Canada.

c. Structural, legislative and attitudinal changes may be required

In many ways, Canadian society from top to bottom has been paralyzed with a kind of hopelessness that says certain things can never be changed. In pollution control, in environmental design, in health care, in social

programs, the focus has been on meeting standards which accommodate the bulk of the population. The hidden assumption is that we cannot accommodate everyone. 'Health for All' contradicts this notion and suggests that we need flexible, clever designs in our physical environments and in our social structures, that will accommodate everyone well.

The literature did not reveal any real barriers to achieving this. It did suggest, however, that achieving healthy environments for everyone will require not only physical changes, but legislative, structural, and attitudinal changes throughout our society's major institutions as well.

d. Participation must be welcomed

Environments will be healthier, when those who are affected by them participate in their design and in their ongoing maintenance and development. This applies in legislative and social program design as well as in physical design. Many methods for user participation have already been developed and need only be selected and applied.

e. There is a worldwide market for expertise in healthy environments

Besides the obvious advantages for the health of people in Canada, a focus on making environments healthier will lead inevitably to economic spinoffs and exportable expertise of products and services. Becoming world leaders in the design and production of safe, non-polluting products and clean, healthy, supportive environments would be consistent with Canada's present world image as a clean, healthy, and peaceful country.

8. Implications for Action

There are four major areas of action that arise out of a commitment to seeing that healthy environments are available to everyone in Canada:

a) political vision and leadership

A new public stance at the highest level is needed to provide leadership in bringing about healthy environments, and to share the following visions:

- i) Environment is important in the health of Canadians;
- ii) Diversity is welcomed and valued in Canada;
- iii) Individuals deserve to control their own environments;
- iv) Prejudice and discriminatory behaviour in Canada must end;
- v) Canada can become a world leader in environmental health.

b) scientific and social research, and industrial incentive

Technologies and methodologies which allow better understanding of the effects of both physical and social environments on health need to be developed and supported. Industries need to be aware that Canadians want and demand cleaner, less damaging, and less risky products and materials. They may require some research support to do this product development well. Assistance in promoting clean, safe Canadian products in world markets might also be offered.

c) public education

We need much more detailed and widespread education about the effects of environment on health, and on ways of creating healthy environments for everyone. We also need to create a better understanding of the fact that there is a wide range of vulnerability to environmental stressors among the population. It is important that this range not be used as a basis for prejudice by people in one part of the range, towards people in the other.

d) legislation

Existing legislative powers need review, to determine the extent to which every person in Canada has adequate control over his or her own personal environment. Health and safety legislation for neglected settings and groups, for example, for schools and for tenants, needs to be encouraged at the appropriate government level. Any aspect of existing legislation, regulations, or institutions that may inadvertently devalue or restrict the opportunities of any Canadians on the basis of their inherent characteristics or situations, requires critical review, with participation by the persons who are affected.

As an initial focus, a complete re-examination of the ways in which sexism is institutionalized should be undertaken, and proposals should be prepared for accelerating the timetable for achieving full equality for women in Canada.

PART II: ANALYSIS

1. Introduction

Section 2 following places a structure on the information we discovered in the literature. Here we offer the reader a synopsis and analysis of the written knowledge about healthy environments. We have pieced it together in a logical sequence, to make clear the underlying messages being put out, in common, by the many people who have contributed their work to the literature.

Health is more than an absence of disease. To conceive of complete health, imagine even getting close to doing every day what is possible on our best day. The concept of health does not in itself include the highest functions of human beings, as typified by the Ghandis or the Mother Teresas of the world; we may not be able to deliberately create an environment which grows that type of individual. We do know, however, that by making the environment negative, we may prevent that type of individual from emerging (Gerdes, K., 1987). The following analysis will look at both the positive aspects, the resources, that are needed to create health, and the conditions which may deny it, for people in Canada.

The subject matter is very broad. It spans physical and social aspects of everything we do in Canada. To organize it into manageable pieces, we have acknowledged three main viewpoints: settings, human characteristics, and environmental resources. In simpler terms, these groups represent: WHERE, WHO and WHAT. Like a three-dimensional matrix, or like the "blind men and the elephant", they each represent one way of looking at a larger problem, and any one issue will have components of all three (e.g. pollution of children in schools, lack of social support for elderly women in retirement homes).

The common message in Section 2.1, which deals with settings for healthy environment issues, is that there is virtually no setting or arena of life in Canada where health and safety do not arise as major environmental issues. The fact that we often think of occupational health and safety when we hear these words, merely means that we have not yet finished the task of formalizing the attention also needed in other settings - school health and safety, tenant health and safety, etc.

One common message in Section 2.2, which deals with human characteristics and situations, is that we are collectively a nation of diversity. We are not all the same. Individually we are also each a model of diversity. We cannot be referred to by one of our characteristics alone, and be adequately considered by others. For each characteristic and situation, there are certain environmental features that are appropriate to health. For each individual, very unique combinations, very self-tailored environments, may be required before we can say his or her environment is truly healthy. This multiple diversity demands a new cleverness in design, to achieve flexible, common environments that accommodate more people well, more of the time, and adaptable, special environments, that suit some of our more unique needs, at least part of the time.

The second common message in Section 2.2 is that many people in Canada have been devalued, for one reason or another. This has usually been done on the basis of one or more characteristics that are part, but by no means all, of their inherent nature or human situation at any given time: age, gender, family type, race or culture, illness or handicap, or orientation and lifestyle.

Being looked down upon is, in itself, an unhealthy environment for those being devalued, leading to loss of self-esteem. The devaluing of one group by another also prejudices the latter's access to environments that are healthy for them. The system really has no winners; most people are part of some devalued group at one point or another in their lives (for example, by growing old, becoming ill, or becoming handicapped).

Section 2.3 describes the environmental and social resources, and hazards, that constitute healthy or unhealthy environments for people. The common message is that everyone in Canada has many kinds of basic physical and social needs. If we are to create truly healthy environments, we cannot just do well in one area of need and ignore the others. Each area discussed is a necessary, but not by itself sufficient, condition for a healthy environment.

For example, all the personal and social support in the world may not prevent us from becoming ill when exposed to toxic contaminants in our environment. All the clean air in Canada will not guarantee health if we have not enough money to place nutritious food on the table. All the good food and housing in the world will not guarantee good health for our young people, if we treat them without respect and deny them sufficient autonomy to plan their own lives.

The keys to creating healthier environments (or the HOW that is missing above), are woven into each viewpoint in the matrix of Section 2, and addressed in summary form in Section 3. The bad news in section 3 is that many things that we have previously thought impossible, and diligently ignored because we couldn't conceive of how to bring them about, are necessary if the concept of 'health for all' is to be achieved. The good news in section 3 is that not only has our knowledge and technology advanced sufficiently to make what we need possible, but we could flourish as a country, not just physically and socially, but politically and economically, if we pursue such goals.

Achieving healthy environments for all is possible, but will require a concerted political, social and intellectual effort to bring it about, and may require extensive institutional, structural and attitudinal changes throughout Canada. Section 3 paints the vision that can and must be brought to life if we are serious.

The analysis in Sections 2 and 3 is limited, for the most part, to what we found in the literature. If the emphasis is on the negative, it is because that is the focus in the literature. Where there are gaps, we have identified them in Part I: Overview.

Healthy Environments for Canadians: PART II: ANALYSIS

The structure in Section 2, the selection from the literature, and the visions in Section 3 are ours; we take full responsibility for their rationality and accuracy. Where our statements are specifically supported by the literature, references have been noted; the reader can follow up by referring to the Annotated Bibliography in Part III, which is ordered alphabetically by author's name.

For the reader who prefers to browse rather than read sequentially, we have reproduced the index for Part II below. We would also encourage you to consult the extensive subject index, found in Part IV, which gives ready access to specific topics in all parts of the report.

2. Healthy Environment Issues	A4
2.1 By Setting	A4
2.1.1 At Home	A4
2.1.2 At School	A8
2.1.3 In Workplaces	A12
2.1.4 In Public Places	A19
2.2 By Human Characteristic & Situation	A23
2.2.1 Age	A23
2.2.2 Gender	A32
2.2.3 Family Type	A36
2.2.4 Race & Culture	A38
2.2.5 Illness & Handicap	A40
2.2.6 Orientation and Lifestyle	A46
2.3 By Environmental Resource	A50
2.3.1 Safety	A50
2.3.2 Clean Air	A51
2.3.3 Food and Water	A56
2.3.4 Physical Factors	A60
2.3.5 Social Conditions	A64
2.3.6 Employment and Money	A69
2.3.7 Personal and Social Support	A72
2.3.8 Health Care and Social Services	A74
3. Bringing About Healthier Environments	A79
3.1 Ending Prejudice and Oppression in Canada	A79
3.2 Creating a New Vision of Environmental Choice	A81
3.3 Fostering Technology for Assessing Our Environment	A82
3.4 Accommodating Diversity in Physical and Social Design	A84
3.5 Becoming World Experts in Designing Healthy Environments	A85

2. Healthy Environment Issues

2.1 By Setting

The environmental literature discusses health issues in a number of settings — at home, at school, in the workplace, in public places, including the natural environment. While some of the problems and opportunities are common to many different settings, there are many aspects which are unique to each. In particular, the amount of control any one individual has over environment varies greatly from setting to setting, with the home often representing the arena of greatest control.

2.1.1 At Home

Accidents in the Home

The risk of death due to accident in the home is about one third to one half of the risk of death in automobile traffic, for those under 65 years of age (Alphey, R.S., 1974). Accidents claim the lives of more children each year than the next six leading pediatric disorders combined, and produce injuries that require medical attention for one in three children. In the preschool age group, 91 percent of these accidents and over one-half the resultant fatalities occur in the home (Dershewitz, R.A., 1977).

Accident types with high injury rates are not necessarily the same as those leading to death. The severity of injuries differs among age groups even for the same accident type. In some cases, building characteristics, such as stair design, may contribute to the accident (Kose, S., 1986).

Many conditions in the home that do not present a high risk to able-bodied people, may present considerable risk to handicapped persons, who must diligently avoid situations which may worsen their disability (Durlak, E., 1987). Sometimes several factors together will make seemingly innocuous home environments more hazardous (e.g. physical reactions to indoor pollutants or medicinal drugs, affecting co-ordination and balance, combined with poor physical design or dangerous habits such as using stairways for storage. (Blumenthal, M.D., 1980), (Small, B.M., 1982).

Additional research on household accidents and injury can be found in (Cassidy, M.W.A., 1970), (Waller, J.A., 1978), (Webber, G.M.B., 1979), (Poyner, B., 1980), (Pauls, J., 1982), (Planek, B., 1982), (Szymusiak, S.M., 1982) and (Kose, S., 1987).

Indoor Air Quality in the Home

A great deal of the research literature reviewed is concerned with the presence of gaseous and particulate contaminants in the air of people's homes. Some pollutants are found in greater quantity indoors than outdoors, indicating that there are indoor pollution sources, and that the pollution levels

are not solely due to infiltration of ambient outdoor pollutants into the home (Colome, S.D., 1982). Most often, the levels of exposure are sufficiently low that it appears that only more sensitive individuals respond adversely. No competent estimates have been made, however, of the numbers of people affected.

A Federal/Provincial Advisory Committee on Environmental and Occupational Health has outlined maximum acceptable residential concentrations, for short and long-term exposures, for nine substances or groups of substances (aldehydes, carbon dioxide, carbon monoxide, nitrogen dioxide, ozone, particulate matter, sulphur dioxide, and water vapour) (Consumer Federation of America, 1987).

Some conditions, such as the use of unvented combustion appliances (gas stoves, kerosene heaters), contribute to pollution levels sufficiently high that they exceed industrial exposure standards, and may present a risk to everyone. The use of gas for cooking has been associated with a significantly increased frequency of chronic cough and a significantly greater percentage of people with impaired ventilatory function (Helsing, K.J., 1982). Other studies giving further information on both the emissions and the health effects of home gas cooking appliances include: (Ferris, B.G. Jr., 1979), (Kelley, M.D., 1979), (Melia, R.J.W., 1980), (Speizer, F.E., 1980), (Traynor, G.W., 1981 and 1982), (Girman, J.R., 1982), (Good, B.W., 1982), (Nitta, H., 1982), (Ozkaynak, H., 1982), (Hosein, H.R., 1986).

Tobacco smoking in the home is another source of indoor air pollution, and has been associated with symptoms of cough, wheeze, and sputum production, in the children of smokers (Dodge, R., 1982). The carboxyhemoglobin content of the blood (from absorption of carbon monoxide) can be as high as 7.6% in the fetuses of smoking mothers, compared to less than 1% in nonsmokers, and 5% to 10% in smokers (Jaeger, R.J., 1981).

The increased interest in energy conservation, through tightening homes to prevent air leakage, has in many cases had the effect of increasing indoor contaminant levels and health risks (Berk, J.V., 1980), (Breysse, P.A., 1981), (Young, G. Stewart, 1981), (Turiel, I., 1985), (Tobin, R.S., 1987). Newer design methods address the goal of combining energy conservation with low-pollution interiors (Small, B.M., 1983).

Products commonly used in the home have been cited as potential contributors to levels of indoor air pollution which could affect health (Turiel, I., 1985). Examples include: scented cleaning and laundry products, treated fabrics and paper products (Imbus, H.R., 1982), refrigerants, insecticides, sponge rubber, plastics, and numerous home finishing materials (Randolph, T.G., 1976). Aerosol hair spray can have significant health effects (Friedman, M., 1977). Art and craft materials used in the home can also be hazardous, particularly when used without extra ventilation. Ceramics, printmaking, painting and sculpting are particularly hazardous. The elderly, children, and people with health problems are especially vulnerable to the effects (Harrison, J., 1983).

Some modern building products, such as urea formaldehyde (UF) resin-bonded particle boards, emit formaldehyde and other gases. A yearly release of as little as one to two percent of the formaldehyde contained in the resin into the air of moderately tight homes can cause the ambient concentrations to exceed the recommended 0.1 ppm limit for formaldehyde. A surplus of formaldehyde from the manufacturing process, and a slowly decreasing generation of formaldehyde from hydrolysis, result in an exponential emission decay with a half-life of about three to five years (i.e. the emission will decrease by half in this time) (Gaudert, P.C., 1985).

Some individuals who are exposed to gases released from such building materials then become sensitized to them (Sprague, D.E., 1982). Urea formaldehyde foam insulation (UFFI), banned in Canada in 1980, caused health problems, as well as fungal growth and structural deterioration of homes (Chown, G.A., 1981), (Campbell, J.S., 1981). Some people who were exposed to UFFI gases, and who had no previous indication of chemical susceptibility, appeared to become generally chemically susceptible as a result of the exposure (Small, B.M., 1982). Sensitivity problems related to housing are commonly the result of continuous, cumulative exposure (Wales, R., 1984).

Some formaldehyde exposure is common in conventional homes, but it is often found in elevated levels in mobile homes (Breysse, P.A., 1979). Symptoms in a new Saskatoon mobile home occupied by an elderly couple included nausea, vomiting, dizziness, headaches, sleep disturbances, shortness of breath, burning eyes, running nose, and chronic fatigue. An initial indoor formaldehyde concentration of 0.7 ppm, was found, far exceeding the recommended limit for residential buildings of 0.1 ppm (Marchant, R., 1985). Senior citizens apartments built with particle-board underlay have also been found to have elevated formaldehyde levels (Kalnins, R., 1985).

Home heating systems have also been cited in cases of building-related illness (Randolph, T.G., 1976), (Silberstein, S., 1979). Ventilation conditions and weather conditions sometimes combine to cause backdrafting of oil or gas furnaces, with accompanying spillage of exhaust products into the home (Small, B.M., 1983). Home humidifiers are sometimes associated with growth of bacteria, which enter the air of the home (Van Assendelft, A., 1979), (Brundrett, G.W., 1979).

Many recent investigations have suggested that radon gas and its radioactive daughter products are present in many Canadian dwellings, and might be a contributing cause of lung cancer in Canada (McGregor, R.G., 1980). The two most important sources for radon in buildings are building materials and infiltration of gases from soil surrounding a home (Hildingson, O., 1982). Radon is also present in water supplies, and may enter the home air while water is running, e.g. during a shower (Hess, C.T., 1982). Decreased ventilation can lead to radiation exposures which exceed present guidelines (Axelson, O., 1979), (Budnitz, R.J., 1979), (Moschandreas, D.J., 1982). Radon concentrations in dwellings vary by more than two orders of magnitude, depending on building materials used, ventilation, infiltration, and regional geological factors, including water supply and substrate (Sachs, H.M., 1982).

Creating Healthy Home Environments

Residential indoor air quality can be improved by attention to principles of low-pollution building design, careful selection of materials, and creative ventilation (Small, B.M., 1983). Particulate matter reduction can be achieved with a medium efficiency extended surface fabric filter, and continuous low-speed fan (Raab, K.H., 1985). Simple cleaning technologies and non-odorous products can also avoid indoor pollution problems.

Choice is important in people's interaction with all environments, and it is central with home environments. Some constraints which may limit and distort any person's choice of home environment include: lack of resources and market availability, inability to cope with or plan for finding a better environment, degree of willingness or ability to move, knowledge of alternatives, and prejudice and discrimination (Rapoport, A., 1985).

Feeling secure at home is also important. Being burglarized is often experienced as being defiled, and can have psychological impacts on a person's relationships with others (Korosec-Serfaty, P., 1985). Home is often more than just our surroundings — a home gains meaning from the psychological and interpersonal events that occur in it (Werner, C.M., 1985), and is treated differently in different cultures (Altman, I., 1985).

Everyone has the right to shelter, but there are still people in our society who are homeless (Greer, N.R., 1986a). When designing shelters, programs, and permanent homes for the homeless, it is most important to take into account their psychological as well as physical needs, including the need for a sense of dignity (Greer, N.R., 1986b). One of the biggest obstacles facing developers of shelter or low-income housing is community resistance (the 'not in my backyard' phenomenon).

2.1.2 At School

School represents the mandatory environment for most of Canada's young people, from about age 5 through their mid-teen years and beyond. Both in the literature and on school board agendas, there is a new interest in the effects of school environments on health and learning. One area in which attention appears to be concentrated at the moment, is that of indoor air pollution in schools.

Unhealthy Aspects of the Physical Environment in Schools

Staff and students in schools are exposed to many pollutants which originate both within their school buildings, and in the neighbouring communities. Many of these pollutants affect brain function, learning ability, behaviour, health, and education. The sources of pollution range from industrial emissions and automobile exhaust outside, to painting, cleaning chemicals, reproduction processes, art and science materials, tobacco smoke, and even chalk dust, inside (Dadd, D.Lynn, 1982). In many schools, ventilation is insufficient to adequately exhaust the pollutants being generated inside. Some students and staff are experiencing acute adverse effects from such exposures, while others do not appear to be suffering, at least in the short term, from the same exposures (Small, B.M., 1985).

Air pollution in schools is rarely diagnosed correctly as a contributing cause of poor scholastic performance of susceptible children, and of the dopiness and confusion of susceptible teachers (Randolph, T.G., 1976), (Faust, H.S., 1981). When exposed to pollutants to which they are sensitive, school-age children can become irritable, excited, depressed, unreasonable or antisocial (Blume, K.A., 1976). In the majority of cases, the brain and central nervous system are the major areas of hypersensitive reaction which is reflected in the child's behaviour and ability to learn (MacLennan, J.G., 1985). Many children who demonstrate an inability to learn, who are slow learners, who have reduced reading comprehension and speed, reduced memory, or who show mental confusion, may improve when placed in a cleaner environment (Rapp, D.J., 1987). There is also evidence that air with a high ambient concentration of negative air ions may contribute to a moderate improvement in short-term visual and auditory memory, compared to regular classroom air (Kershner, J., 1985).

Formaldehyde, ammonia, hydrogen sulfide, sulfuric acid, carbon monoxide, sodium hydroxide and chlorine, among others, are used in the manufacture of various types of paper used in schools. The handling of most newspapers, magazines, hardcover and paperback books, art paper, paper tissues, plates and cups and glossy paper readily produces dermatitis in formaldehyde-sensitive individuals. Carbonless paper can produce mild dermatitis and severe upper respiratory symptoms as well as eye irritation. Chemical agents in various types of photocopier papers also causes dermatitis in some people (Fisher, A.A., 1983).

Some of the volatile inhalants emitted by products used in schools pose a health problem because of their neurobehavioural toxicity, and share certain properties with classic central nervous system depressants. For example, toluene and 1,1,1-trichlorethane have some pentobarbital-like effects (Rees, D.C., 1987).

Methyl alcohol evaporating from spirit duplicators, and from the copies themselves (which students like to sniff), represent a very specific hazard in schools, although many of these machines have been replaced in light of the evidence of this hazard. Adverse health effects reported include blurred vision, headaches, burning of the nose, sluggishness, dizziness, sore throat, dermatitis, chest tightness, and depression, all characteristic of toxic exposure to methyl alcohol (Pryor, P., 1981). In two cases persons exposed to methanol fumes initially developed clinical symptoms of multiple sclerosis, including visual disturbances, reduced abdominal reflexes, impaired coordination and difficulties with walking (Henzi, H., 1984).

Many schools were built with friable asbestos insulation material in the ceilings. Asbestos fiber concentrations in the air in several schools studied were in the range 5-40 ng/M³ before removal of the insulation. The airborne asbestos concentrations approximately one week after removal showed reductions of 56 to 90 percent (Bozzelli, J.W., 1981).

Building-related illness ("the sick building syndrome") has been reported with increasing frequency in schools and kindergartens (Subcommittee on Formaldehyde and Air Contamination in Public Buildings, 1983), (Gravesen, S., 1986). The construction of the buildings, with flat roofs, often leads to water damage with subsequent microbial growth. Further, reduced cleaning budgets combined with wide use of needle-felt carpets, as well as improperly maintained ventilation systems, will lead to pollution by dust and microorganisms (Gravesen, S., 1986). Infection can also be spread among school-age children, by airborne viruses and bacteria circulated through the ventilating system, and from exposure in school buses (Riley, E.C., 1978) (Lidwell, O.M., 1979).

In Danish schools, significantly more numerous cases of allergy (asthma and hay-fever) were found in schools with textiles as floor coverings and significantly more numerous cases of skin conditions were found in those schools which were less than five years old or which had been renovated within the past five years. Some 31% of the pupils and 40% of the teaching staff stated that they took headache medicine at least once per month. Under one third of these pupils and over half of these teachers took headache medicine at least once weekly (Ibsen, K.K., 1981).

In 1973 a study was conducted by the Environmental Health and Light Research Institute in Sarasota Florida, showing dramatic reactions in hyperactive children. In two first-grade classrooms, the standard cool white fluorescent tubes and fixtures with solid plastic diffusers remained unchanged. The plastic diffusers stopped the transmission of any trace of long-wavelength ultraviolet. In two other classrooms, the tubes were replaced

with full-spectrum fluorescent tubes that more closely duplicated natural daylight. By means of hidden time-lapse cameras, student behaviour was observed and recorded.

One study reviewed showed that under standard cool white fluorescent lighting, some children in first-grade classrooms demonstrated nervous fatigue, irritability, lapses of attention, and hyperactive behaviour. Within a week of the new full-spectrum lights being installed, a marked improvement in their behaviour appeared and overall classroom performance improved (Ott, J.N., 1973). Others feel that there is insufficient data to support a connection between standard fluorescent lighting and hyperactive behaviour (Wotton, E., 1981). However, it was also suggested that students working under fluorescent lighting with very good colour rendering become less visually fatigued than those working under some other forms of fluorescent lighting.

Other Potentially Unhealthy Aspects of School Environments

Increased psychological stress in the school setting, specifically examination stress, will alter immune functioning and heighten psychological responses (Didriksen, N., 1986).

Parental smoking, socio-economic status and scholastic performance are all related to smoking in high school students. Of the three factors, poor scholastic performance was clearly the strongest correlation, with parental smoking next in strength and low socio-economic status a weak third (Borland, B.L., 1975).

A severe isolation from real-life experiences in the education system has led to a separation from the fact that learning is an activity which is natural to human beings. Children have ceased to be authentic participants in the life of society and would learn better in an environment free from compulsory schooling and rigid age segregation (Priesnitz, W., 1987).

Making School Environments Physically Healthier

A Working Group of Ontario School Boards is presently reviewing procedures for "total building performance" evaluations, as a means of addressing indoor environmental concerns as they arise, and in the design of new buildings (Gosnell, D., 1987). The procedure, developed by the Toronto Board of Education's Energy Conservation and Environment Department, recognizes that a variety of elements contribute to occupants' satisfaction with a building environment, and also recognizes the special needs for those who experience greater sensitivity to the local environment. The procedure encourages all occupants to participate in the planning, design, problem-solving and operating practices of their building. Part of the rationale for the procedure is to give individuals or groups the tools to solve building environment problems.

The same group has begun to make it known to manufacturers and suppliers of materials used in schools, that they want to know, when tenders are submitted, what chemicals each object or material will emit into the air of the schools, or what contaminants will be absorbed by staff or students handling the materials (Gosnell, D., 1987). This is part of a long term program to reduce indoor pollution in the schools, by using only well-researched, "school-safe" materials that do not pollute indoor air (Small, B.M., 1985).

The Toronto Board of Education's Pollution and Education Committee has expressed a parallel intention, to investigate applications for rezoning of land within approximately 1000 feet of school sites, in order to monitor any possible outdoor pollution sources that might pollute school grounds or school air intakes (Gosnell, D., 1987). The Toronto Board is also encouraging all schools in its system to establish 'pollution relief centres', or rooms which are kept particularly clear of contaminants and dust, for use by sensitive students and staff.

The first special low-pollution, or "ecology" classroom was established in Kitchener, Ontario, to address the needs of students who had developed severe chemical sensitivities. Teacher Brad Tucker, one of the driving forces behind the project, is particularly knowledgeable about dealing with the special problems encountered by sensitive students. Some students, who could at first only tolerate the special classroom, are now able to spend more time in the rest of the school, using the ecology classroom as an oasis if reactions flare (Toronto Star, 1985).

The York Region Roman Catholic Separate School Board, operating north of Toronto, Ontario, is also planning to build two special ecological classrooms for children who suffer from severe allergies. The classrooms are part of a strategy to decrease the number of allergens in school buildings, generally, while at the same time addressing specific needs of highly allergic children. New schools in the system are being designed to take into account the average child's allergies (Kleiman, C., 1987).

There are many practical solutions that can be applied easily at the individual school and classroom level, to indoor air pollution problems in schools. These include careful control of all materials entering the classroom, strict school rules prohibiting smoking in the building, and education of staff and students about the effects of perfumed personal hygiene products on sensitive individuals (Schreiter, A., 1984). Classrooms and academic offices are more suitable for chemically susceptible individuals when fresh air is introduced directly into the room, rather than through a central recirculation system which mixes fresh air with large volumes of polluted air (Tall, F.D., 1987).

There are also many basic design criteria which can be applied in the planning of new school facilities and upgrading of old ones, to ensure a healthy working environment. Some areas for particular attention include reduction of noise, storage of volatile substances, and ventilation of odours

in special purpose classrooms including woodworking shops, automotive shops, fine arts studios, music rooms, science laboratories, changing rooms, and various occupational training areas (Ball, R., undated).

Measurements of newly built preschools in Sweden show that organic compounds emanating from building materials decline in concentration mainly within the first six months of occupancy. During that time, symptoms such as dry throat, irritation of eyes and lips, hoarseness, hacking cough and itching are often reported. One way of assuring a healthier environment even in new schools is to allow the building materials to gas off during the first six months after construction, with no recirculation of return air allowed. During at least one to two additional years, the recirculation rate of return air can also be restricted, perhaps to 50% (Berglund, B., 1982).

There is a lack of safety standards for school science laboratories in Canada, and no systematic monitoring of safety conditions in some jurisdictions (Rieber, E.R., 1984). At the same time, the school is an ideal place to instill a lifelong awareness of potential hazards (Borrows, P., 1984). Development of consistent standards, procedures for monitoring, and programs for spreading information about school health and safety, would represent practical and useful steps toward healthier school environments.

2.1.3 In the Workplace

Physical and Chemical Hazards in Industrial Settings

In the past, many occupational diseases were named after the specific trades or occupations in which they were first observed. A few diseases still remain quite specific, but many biological, chemical and physical hazards are now so widespread that they may be encountered in jobs in a wide variety of industries (Proctor, N.H., 1978), (Medical Services Division staff, Workers' Compensation Board, 1984).

Chemicals of organic and inorganic nature, met as dusts, vapours, and fumes in the workplace, can cause respiratory allergic disorders. Widely different agents in different forms are implicated in asthmatic reactions (Pepys, J., 1982). It is also not unusual for industrial workers to experience deterioration in behavior or in mental function after exposure to common industrial chemicals. For example, a simple measurement of mercury exposure will give a reliable prediction of certain memory problems (Batts Young, B., 1981).

Neurotoxic volatile organic solvents used by house and car painters may lead to professional toxic encephalopathy after several years of exposure. The symptoms are memory impairment, fatigue, personality changes, headache and dizziness. Inner ear dysfunction is also common, and may be helpful in detecting early changes in exposed persons and in determining more accurate safety limits for harmful chemicals (Selikoff, I.J., 1975), (Arlien-Soborg, P., 1981). Peripheral neuropathy has been discovered among spray painters (Mallov, J.S., 1976). Central nervous system and psychological disturbances have been observed among workers exposed to styrene in glass fibre manufacturing operations (Lindstrom, K., 1976), (Axelson, O., 1978). Long-term exposure to solvents is also linked to chronic but nonspecific neuropsychiatric conditions (Axelson, O., 1976), (Hane, M., 1977).

Formaldehyde is often found in industrial environments, particularly where resin compounds which are employed as industrial laminates, binders and adhesives are produced or used. The potential for diseases of the skin has been well documented; it can induce dermatitis by irritation, delayed-type hyper-sensitivity and immediate urticarial reactions. Its capacity to irritate the eyes and upper respiratory tract is well established. Observations to date also incriminate formaldehyde in the induction of an inflammatory bronchitis as well as bronchial asthma (Bardana, E.J., Jr., 1980).

The welding process, common to many industries, produces potential respiratory hazards in the form of respirable metal oxide fumes and various gases, of which ozone and oxides of nitrogen are the most dangerous (Liss, G.M., 1987). The industrial setting can also include exposures to toxic heavy metals such as cadmium (e.g. in electro-plating plants). Cadmium can accumulate in tissues, and health effects of excess exposure can include lung damage and kidney damage or failure (Gibson, B., 1985).

Many industrial situations involve considerable exposure to diesel exhaust, which carries many health risks including possible carcinogenicity (Pepelko, W.E., 1980), (Unknown, 1982). Polycyclic aromatic hydrocarbon (PAH) compounds have been identified as the most potentially hazardous agents found in diesel engine exhaust. Although a properly maintained diesel engine would be likely to emit extremely small amounts of these agents, the possibility that they might be carcinogenic to humans singles them out for careful attention (Lassiter, D.V., 1978). Experiments with rats have suggested that diesel exhaust exposure during the development of an organism can lead to behavioural differences in adulthood (Laurie, R.D., 1980). Experiments with mice have shown an increased susceptibility to bacterial infection following diesel exhaust exposure, presumed related to the nitrogen dioxide and acrolein vapour components. (Campbell, K.I., 1980). Evaluation of the toxicity of the major components of diesel exhaust, with a view to developing add-on exhaust hardware, and to developing other techniques to reduce emissions from diesel engines, is of current interest (Dainty, E.D., 1986).

A variety of solvents and other potentially hazardous compounds are typically used in art workshops, including acetone, styrene, toluene, benzene, methyl cellulose acetate, xylene, and others (Wadden, R.A., 1985). Artists are among the highest sufferers of cancer, and other diseases, due to exposure to solvents, heavy metals, mineral dusts, gases, and other hazardous chemicals such as acids and alkalis. (Visual Arts Ontario, 1981).

Construction workers are exposed to a wide variety of toxic materials about which they know very little, and over which they have little control. The traditional solutions in fixed industry, using substitution, ventilation, process modification and administrative controls, are not always possible for construction sites (McVittie, D.J., 1986). Modification of building design to reduce indoor contamination for the sake of future building occupants could at the same time reduce some of the toxic exposures for construction workers.

Tobacco smoke, which contains over 50 known carcinogens and many other toxic agents, is a health hazard for smokers as well as nonsmokers who are regularly exposed to it. Serious acute health effects in nonsmokers, from short-term exposure, are probably limited to the one fifth of the population with pre-existing health conditions that are aggravated by exposure to tobacco smoke. Risks for everyone of long-term exposures include decreased lung function and lung cancer. The composition of tobacco smoke is such that there may not be a safe level for such exposure (Collishaw, N.E., 1984).

There are also hazards from the exposure to fungi in the workplace (Salvaggio, J.E., 1986). The diseases they cause may be so mild as to be asymptomatic, or may be extremely debilitating. Some buildings have humidifier and/or ventilation systems which are contaminated with fungi, and which may cause symptoms among the occupants. (Giddings, M.J., 1986).

The reader is referred to further descriptions of industrial environmental hazards in the literature, which are illustrative of the general points made above, for example: asthma produced by exposure to toluene di-isocyanate (Chester, E.H., 1979), (O'Brien, I.M., 1979); effects of chronic occupational

exposure to benzene (Fishbeck, W.A., 1978); reproductive hazards at work (Chenier, N.M., 1982); occupational hand eczema (Pedersen, N.B., 1980); and neurobehavioural toxicity (Wood, R.W., 1981).

The interaction of workplace conditions and other factors can increase the risk of injury on the job. For example, delayed effects of social alcohol intake have been observed up to 18 hours after ingestion. These included lengthened reaction time, poor motor performance, and decreased motor sensory skill, as well as inability to manipulate and position without tactile and/or visual facilitation (Wolkenberg, R.C., 1975). Abuse of other substances, including illicit as well as prescription drugs, can also increase risks. Close co-ordination will be required between management of alcohol and drug-related problems and the management of safety in the workplace (Shain, M., 1982).

Recognizing the possible hazards associated with one's chosen occupation, whether it be exposure to organic dusts, chemicals or dangerous machinery is of greatest importance for prevention of injury or infection (Ontario Federation of Labour, 1982), (Canadian Centre for Occupational Health and Safety, 1985). Persistent symptoms, no matter how small, should be regarded as important, and measures taken to locate their source. Care must be taken when identifying the cause of a worker's complaint, as these diseases may be misdiagnosed, which in turn may lead to irreversible physical damage (Giddings, M.J., 1986).

The true dimensions of industrial disease are not totally revealed by Workers Compensation Board statistics. Often the real causes of disease and injury on the job are ignored, and economic considerations prevail at the expense of the health and safety of the worker. There is widespread resistance of both senior and middle managers, to increased worker participation in both work organization and job design questions. Greater workplace democratization may be a necessary condition for bringing about adequate reform of workplace health and safety (Sass, R., 1986).

Physical and Chemical Hazards in Offices

Substances such as ammonia, asbestos, benzene, cadmium, carbon monoxide, ethanol, fiberglass, formaldehyde, methanol, nitropyrenes, ozone, PCBs, particulates, radon, tobacco smoke, toluene, trichlorethane, trichloroethylene, trinitrofluorenone and vinyl chloride are all present in the typical office environment, albeit in small quantities. Some are suspected or proven carcinogens, while a number, such as trichloroethylene, carbon monoxide and benzene have been implicated in central nervous system damage (Makower, J., 1981), (Nussbaum, K., 1981), (Konopinski, V.J., 1983). Office contaminants are usually present in very low concentrations relative to promulgated or recommended industrial hygiene exposure levels, but in high concentrations relative to outdoor air. Sources include new and aged building materials, wet-process photocopiers, tobacco smoke, and building maintenance products (Miksch, R.R., 1982).

Recent investigations suggest that employees who work in sealed office buildings make more mistakes, have more office accidents, take longer to do tasks and are sick more often (Cannon, M., 1987). Inadequate ventilation in such buildings can also bring on symptoms of headaches, drowsiness, tired eyes and upper respiratory irritation (Rand, G., 1979), (Hollands, J., 1984). Severe outbreaks of illness have also been traced to ventilation problems in sealed hospital buildings (Sterling, T.D., 1983).

In one investigation of health and environmental complaints in a large, modern, hermetically-sealed office complex, there was little indication that the type of symptom and type of environmental complaint were correlated in any meaningful way (Benard, J.M., 1985). Other investigations have shown under more controlled conditions that specific sources can produce certain symptoms. For example, exposure to airborne emissions of carbonless copy forms can cause acute nasal congestion (Camp, J.E., 1985). Handling such forms can produce urticaria on the hands, and changes in lung function characteristic of upper airway obstruction have also been observed (Marks, J.G., Jr., 1984). The offending agent in the paper which caused the allergic contact dermatitis proved to be a colour-forming coating composed of paratoluene sulfinate of Michler's hydrol (PTSMH), which has a low sensitizing capacity (Marks, J.G., 1981).

The current technologies used in photocopying machines involve a number of health concerns, including the refilling and disposal techniques for toner in dry machines, levels of isodecane produced by wet-process machines, the levels of ozone produced, heat and noise levels, and escape of ultraviolet light (Halton, D.M., 1983).

Most commercial settings experience carbon monoxide concentrations above zero indoors, because it tends to seep into buildings from vehicular emissions outside. The levels commonly found are usually below 5 ppm and seldom higher than ambient air quality standards. However, indoor garages and buildings with attached indoor parking areas are exceptions and can experience relatively high carbon monoxide concentrations (Ott. W., 1982).

Tobacco smoke exposure is greater, on average, in the workplace than it is in the home (Repace, J.L., 1983), and has often been assumed to be a major cause of building illness. While there is no doubt about the carcinogenicity and the allergenicity of tobacco smoke, (Repace, J.L., 1983), (Repace, J.L., 1984), there is still some question as to its relative importance, compared to other sources, as a cause of building illness (Sterling, T.D., 1987).

Other factors besides chemical sources and inadequate ventilation may also come into play in modern office environments. Artificial lighting, high noise levels, fluctuation in temperature and humidity, and low-level radiation and electromagnetic fields (e.g. from video display terminals) are also being investigated as potential health hazards capable of causing stress and illness (Makower, J., 1981), (Sterling, E.M., 1983), (Fleishman, J., 1984).

Creative office design and effective ventilation, along with strict attention to chemical sources and other physical factors that are capable of adversely affecting health, may be required before the now-common 'sick building syndrome' is under control. It is also becoming clear that more medical knowledge is required concerning the health of individuals affected, because of the potential for increased sensitization that may last well beyond remedial changes in the office environment (Small, B.M., 1984), (Small, B.M., 1985). Effective control strategies can be implemented that do not compromise energy efficiency (Miksch, R.R., 1982).

The Range of Vulnerability Within the Working Population

Some workers react adversely to certain substances at low levels, below the threshold concentration associated with injury or definite discomfort in the majority of workers. These particular workers are distinctly more susceptible than the majority, and may be inadequately protected by control procedures that appear to suffice quite well for most employees (Reinhardt, C.F., 1978). At the same time, some situations that demonstrably affect only susceptible workers, may still present longer-term dangers to the health of so-called nonsusceptible workers. Reducing total environmental chemical exposures may be the most appropriate solution to the risk problem for everyone (Wright, M., 1979).

In the past, episodes of building illness involving groups of people were often labelled as mass hysteria. The subjective symptoms documented are similar to symptoms experienced by persons being tested for susceptibility to various petrochemicals. Specific testing for chemical hypersusceptibility may remove the uncertainty in such cases (Miller, C.S., 1979).

Psychological Factors in the Workplace

Most people generally like their jobs, are interested in their work, and are comfortable with their supervisors and their fellow workers (Skrzycki, C., 1987). However, many conditions and attitudes within the workplace can be stressful, and need to be carefully examined if truly healthy environments are going to become available in workplaces throughout Canada.

Poor mental health is directly related to unpleasant work conditions, the necessity to work fast and to expend a lot of physical effort, and to excessive and inconvenient hours (Cooper, C.L., 1976), (Resin, T.S., 1978). Inability to adjust and control ventilation, lighting, temperature, and humidity to individual comfort needs represents a stress that has a significant influence on health and comfort of office workers. Health-related symptoms can also be aggravated when employees are unable to participate in the way time at work is allocated, when jobs are not secure, and when there are poor relations with supervisors (Sterling, E.M., 1983).

In the workplace, creating an environment of sensitivity, consideration and tolerance of differences is considered a positive mental health practice (Mental Health and the Workplace Committee, 1984). Helping people find work situations that allow them to reach personal goals they define for

themselves, can make an organization a dynamic, fulfilling place to work, and creates the type of climate conducive to effective performance. Worklife choices may include leaves of absence, job sharing, flexible part-time work, transfers, and other choices (Hopkirk, G., 1986). The organization of worktime must also respond to the specific needs of women, who now make up a significant part of the labour force (Simard, M., 1986).

With new technology now available, even more flexible work-options may become commonplace. For example, running a computer-based business from a home office is a work alternative that allows flexibility and control over the working environment. Many people are able to integrate family and work lives when they work at home. Higher efficiency, economy, reduction of stress, and the elimination of commuting time are other benefits. Telecommuting has possibilities for creating employment for the disabled (Chalupiak, S., 1987). Care will be required during the expansion of such alternative work-options; employment of home-based workers as independent contractors, for example, can present the risk of creating second-class corporate citizens without full employee rights and benefits (Christensen, K.E., 1986).

2.1.4 In Public Places

There are many healthy environment issues regarding public places in Canada. The literature concentrates more on urban-related health issues, and our discussion below reflects this, taking the opportunity to designate cities as a kind of public place. It is important, however, to remember that all Canadian settlements, of any size, need the public places within them, as well as their general environments, to be healthy for everyone. Designing for the diversity of people, and the diversity of uses, of public spaces, represents a considerable challenge.

At the same time as Canadians may be very diverse in some of their characteristics, we must remember that urban dwellers, suburbanites, people in small towns and settlements, and in rural or wilderness areas still have very many human characteristics, goals, and needs in common (Baldassare, M., 1975). Many of the details of the following discussion will therefore have some applications in all these settings.

Indoor Air Pollution and other Hazards in Public Buildings

In general, the move to increase energy conservation in public buildings has led to increased concern about health effects from indoor air that is being contaminated while being recirculated (Dimmick, R.L., 1980). Special purpose public buildings such as transportation terminals (e.g. train stations and bus depots), and medical facilities, also present special indoor air quality problems because of strong indoor pollution sources (Suess, M.J., 1984).

Specific indoor pollution problems are increasingly showing up in the literature. For example, high formaldehyde levels have been found in day care centres where extensive use has been made of particle-board (Olsen, J.H., 1982). Lack of ventilation can also lead to unacceptably high concentrations of carbon dioxide (Lundqvist, G.R., 1982). Dried detergent residue left in carpets after they were shampooed has caused respiratory and eye irritation among both day care staff and children (Kreiss, K., 1982).

In other situations we have taken for granted that traditional technology is benign, only to discover, upon measurement, that major health hazards may have existed for years. For example, episodes of illness have been reported in ice arenas, which have been traced to high levels of carbon monoxide and nitrogen dioxide from ice-resurfacing machines. Typical symptoms have been headache and nausea among children and headache among adults, as well as sore throat and tightness in the chest. Carbon monoxide concentrations up to 250 ppm have been found, which far exceed acceptable outdoor ambient air quality standards (Anderson, D.E., 1971).

Fortunately, more jurisdictions are moving towards severe restrictions and even total prohibitions on tobacco smoking in indoor public environments, including business offices. About one-third of Canadian adults smoke regularly. The primary obstacle to overcome is that smoking is a powerful addiction with failure rates after cessation as high or higher than those for heroin addicts or alcoholics (Wigle, D.T., 1982).

The physical design of public buildings and outdoor spaces is also important for ensuring safety for Canadians. Stairway accidents, for example, are largely caused by design shortcomings and can be largely prevented by thoughtful design features (Asher, J.K., 1977).

Air Pollution in Canadian Communities

Air in communities throughout Canada, from remote settlements to dense urban areas, is contaminated with low levels of pollutants emitted by industrial sources, municipal incinerators and utilities, motor vehicles, household heating systems, fireplaces and woodstoves, and local use of pesticides and herbicides. The literature is clear that air pollution is a health hazard, both to the general population and to many individuals who, for one reason or another, find themselves considerably more vulnerable than others to its effects. There is no doubt that aiming for the lowest air pollution levels that can be achieved, within our resources and our technology, is working in the direction of healthier environments for all Canadians.

In high sulphur dioxide and smoke pollution areas, adverse effects on ventilatory functions and an increased occurrence of acute respiratory diseases can be expected (Saric, M., 1981). Prevalence of respiratory disorders and allergies can also be a function of proximity to major roadways, because of exposure to automobile exhaust (Nagira, T., 1981).

The effects of carbon monoxide from automobile exhaust are particularly marked for those involved in vigorous exercise. The endurance performance of distance runners in urban environments, for example, is subject to interference from commonly encountered carbon monoxide concentrations for many hours after exposure to the gas has ended. While at rest, breathing air containing 50 ppm of carbon monoxide causes the concentration in the bloodstream (carboxyhemoglobin, or COHb) to rise to 5% in five hours. During vigorous exercise, this level of air pollution can produce a concentration of 5% in less than one hour. It is not metabolized but must be dissipated by being exhaled (Daniell, H., 1976). Additional Effects of low levels of carbon monoxide encountered on urban streets and in traffic tunnels have been documented in the literature (US National Research Council, 1969).

Absorption of auto exhaust fumes also leads to elevated whole blood lead levels in distance runners. While the elevations do not appear to constitute a short-term health hazard for adult runners, the longterm or chronic effects of these "above average" levels are not known (Van Rensburg, J.P., 1982). Lead from vehicles may not be the most significant contributor to blood lead levels of children living or being educated in high-traffic urban areas (Millar, I.B., 1982). Urban soils and settled dusts, both indoors and outdoors, could have a significant impact on human health, because they also carry heavy metal pollutants, including lead and cadmium, from industrial sources (Solomon, R.L., 1977). Urban atmospheres also receive gas and particulate emissions from automobile brakes and clutches, including asbestos particles (Jacko, M.G., 1973).

Waste incineration emissions are also receiving attention as potential health risks, since data on health effects of many chemicals are incomplete, particularly for long-term, low-level exposures. Technical and operational measures need to be implemented to reduce emissions of chemicals, and questions need to be asked about siting of new incinerators and about acceptable levels of health and environmental risk (Macpherson, A.S., 1987).

Are Cities Healthy?

Some may argue that because of the tendency for urban areas to have higher pollution concentrations than non-urban areas, cities themselves are, by nature, unhealthy. It is important, however, to consider both the specific design of a city, and the specific technologies being used within it, before being tempted to declare cities, in general, as unhealthy. The choice of the gasoline engine, for example, for personal transportation, has had an enormous adverse impact on pollution and health. It is more useful, in terms of actions we can undertake to improve environments, to conclude very specifically that gasoline engine technology creates unhealthy environments, than it is to conclude that cities are unhealthy. Cities with alternative, cleaner technologies can certainly be imagined and ultimately created.

There are, of course, many aspects of cities the way they exist now that are unhealthy: noise, air pollution, traffic congestion, crowding and lack of open areas of trees, lakes, grass, and rivers. Many people within our cities are subject to additional stresses, such as living in higher crime areas, working in jobs exposing them to toxic chemicals, or being denied an adequate education due to discrimination (Kennedy, D.A., 1977). The chronic, cumulative health effects of such stresses may be of considerable importance (Bakacs, T., 1972). At the same time, cities are high-energy places with many activities, which in itself can be very health-supportive to urban dwellers (Humiston, K., 1987).

In urban areas, health risks from physical problems were given a great deal of early attention (e.g. sanitation), while those from social conditions (e.g. overcrowding) were often ignored (Weinstein, M.S., 1987). Now, researchers are putting more attention on social structure and participation of the city dweller in ongoing environmental design. "Solutions must come from the grass roots level, and must deal with the underlying issues, not the symptoms" (Duhl, L.J., 1984). Isolated specialization in urban planning, without public input, can be destructive to the quality of urban life (Appleyard, D., 1976).

Further improvement of urban environments is required: to remove causes of danger; to maximize opportunities for contact between individuals at block, neighbourhood and town levels; and to enhance the visual and aesthetic qualities of the urban environment, as one means of promoting a favourable change in individual psychological behaviour (Pressman, N.E.P., 1982).

Natural Environments

The outdoor environment is an important factor in children's social and psychological development. High-rise buildings, traffic, long distances to attractive play spaces, and bad weather, may restrict children from staying outdoors, as will a lack of suitable activities for adults who accompany their children to the playgrounds. Allowing children opportunities for concrete action upon the environment (e.g. digging, building things) helps them to develop a sense of competence, and encourages them actively to bring about changes in the environment (Bjorklid, P., 1985). Creating more habitable environments requires understanding of natural settings — air, water, geology, plant and animal life (Whiston Spirn, A., 1984).

The United Nations Conference on the Human Environment (United Nations, 1973) declared that both the natural and "man-made" environments are essential to people's well-being and to the enjoyment of basic human rights, and that we must shape our actions with a more prudent care for their environmental consequences. The conference called for: protection of non-renewable resources; the halting of the discharge of toxic substances and the release of heat in such quantities as to exceed the capacity of the environment to render them harmless; economic and social development for ensuring favourable living and working environments; integrated and coordinated approaches to development planning so as to ensure that development is compatible with the need to protect and improve environments; education in environmental matters for both young and old; scientific research and development to solve environmental problems; and international cooperation to spare the earth the effects of nuclear weapons and all other means of mass destruction.

Many natural and synthetic compounds have entered the ecosystem in the past 40 years. The adverse effects of many of these were unanticipated, in that these compounds are highly resistant to degradation by natural processes and there is often a pervasive buildup on a worldwide basis. In the case of carcinogens, mutagens and teratogens, safe levels cannot be established because even extremely small doses must be assumed to result in an increase in the number of people who are adversely affected (Ray, S.M., 1980).

Acidic precipitation, due to the growing introduction of sulphur dioxide into the atmosphere, has been linked with detrimental effects on fish, vegetation, buildings and people (Kruus, P., 1979) (Kato, T., 1981). According to a new United Nations report, acid rain is now a health threat to more than half a billion city dwellers worldwide, and another billion people are exposed to high pollution levels that result from coal, wood, oil combustion and automobile traffic dust (Toronto Star, 1987). Studies on the indirect effects of acid depositions on water quality have shown that when acidified lake water is left standing in the plumbing system the quality of the water can deteriorate; levels of copper and lead found in it can exceed federal guidelines (Franklin, C.A., 1985). Environmentally-sensitive people may be the first to feel the effects of low-level environmental contaminants (Bradley, J., 1987).

Healthy Environment Issues (continued)**2.2 By Human Characteristic and Situation**

For each human characteristic, and for each unique combination of human characteristics and situations, "healthy" environment will take on a different meaning. What is healthy, or at least benign, for one person, may be unhealthy for someone with different characteristics. The key to understanding healthy environments is to understand people in considerable detail, and in particular, to understand that human beings are, by nature, a very diverse lot.

Many people are subjected to environments which are unhealthy. In many, if not all, such situations, the same people also happen to be subject to prejudice because of some specific characteristic (e.g. being old, being handicapped, being female). While there are specific physical or social conditions in these environments (e.g. pollution or crowding) that are the direct causes of ill health, the devaluation by others is the mechanism by which these people get stuck in, or stay trapped in, unhealthy conditions. The message conveyed is a simple one: "we don't care if your environment is unhealthy, because people like you aren't as valuable as people like us".

In each of the subsections following, we address a particular array of human characteristics, with which we often distinguish one person from another. Where we have found sufficient information in the literature for each characteristic, we have addressed these issues:

- 1) How people who happen to have this particular characteristic are sometimes devalued by people with different characteristics;
- 2) How people with this characteristic are mistreated; how they get stuck in unhealthy environments more than people with different characteristics;
- 3) What the real differences are that go along with having this particular characteristic; how these differences compare to common stereotype images;
- 4) How people with this characteristic differ in vulnerability to environmental factors;
- 5) What environments would be healthier for people who happen to have this particular characteristic.

The array of human characteristics discussed is not exhaustive. These particular ones were chosen because they illustrate well the manner in which the devaluing of one group of people by another may contribute to unhealthy environments.

2.2.1 Age

In Canada, we sometimes vary our treatment of people, on the basis of the age that they happen to be at the time, relative to our own age. People who are very young, and people who are very old, are often treated very differently from the rest of the population, who are temporarily inbetween these two extremes in age.

The rational basis of this variation in treatment has to do with distinct physical and other differences in human beings of different ages. Many features of the environment that a person who is under 10 years of age will need, to be healthy and to grow, will differ from those of the environment needed for someone who is, for example, between 40 years and 50 years of age.

The irrational part of treating people of various ages as differently as we sometimes do, stems from our use of stereotyped images of people who are a certain age, without reference to the considerable variability among people who happen to be in the same age group, and without reference to the many human characteristics we still all share, despite the difference in our ages.

The Devaluation of Young People

Young people have suffered innumerable injustices as a result of ingrained attitudes on the part of people older than themselves. Like racial and cultural minorities, like women, like those missing one or several of the "normal" human traits, children have been stereotyped as somehow inferior (Davies, F.L., 1988). Despite surveys showing that Canadians place the issue of child health high on the national agenda, people at the Canadian Institute of Child Health often feel that children are Canada's forgotten people (McCauley, G., 1987).

How Young People Differ in Vulnerability to Hazards

Many of the physical differences between younger people and older people are obvious, for example, size and strength. Internal physical differences are less visible, but just as important. For example, pre-school children have a relatively higher metabolic rate and ventilation at rest than older people, leaving them more vulnerable to inhaled pollutants (Stephens, R., 1981).

The embryo and foetus are highly vulnerable to environmental assault, and as a result, the prenatal period is considered a critical stage in human development. Avoidance of social, prescription and illicit drug use, as well as other harmful chemical exposures, may prevent many adverse outcomes for the baby (McCluskey-Fawcett, K.A., 1986). For some prenatal chemical exposures, such as carbon monoxide, some researchers conclude that no threshold can be established below which there is a margin of safety for the foetus (Waterman, F.K., 1984).

Environmental factors to which a young person is exposed in the womb can sometimes be contributing causes of birth defects or cancer. Some that have been implicated include radiation, chemical agents such as nitrosamines and organic solvents, and infectious agents such as the Epstein-Barr virus. Certain compounds may lead to birth defects when a prenatal exposure occurs early in pregnancy, and may act as carcinogens when the exposure occurs late in pregnancy (Gordis, L., 1986).

Young children appear more susceptible to exposure to heavy metal contaminants, particularly in combination, than are older people (Chisolm, J.J., 1974). Children absorb and retain a greater percentage of ingested lead than adults (Schauss, A., 1980). Pre-school children also have more contact with contaminated soil outdoors than older children or adults (Stephens, R., 1981).

Early exposure to lead can lead to subtle neurological deficits and behavioural impairments. These effects may be immediate or delayed (Finberg, L., 1974) and are often not recognized until the child enters school, when he or she exhibits a short attention span, hyperirritability and aggressiveness, sensory and motor impairments (Weiss, B., 1974). Marked reductions are also seen in general development of the hippocampus portion of the brain, following postnatal lead exposure (Petit, T.L., 1983).

Research on the effects of such contaminants on children has become highly specific. For example, cadmium has a significantly stronger effect on verbal IQ than does lead, and lead has a stronger effect on performance IQ than does cadmium. Hair cadmium and lead levels have been significantly correlated with both intelligence scores and school achievement scores, but not motor impairment scores (Thatcher, R.W., 1982). A strong case has been presented in the literature that lead at low doses is an important and widely distributed neurotoxin, and that particularly for children, its removal from the human environment is warranted (Needleman, H.L., 1983).

Long-term exposure of children to contaminants such as formaldehyde, from certain building materials, leads to a higher occurrence of abnormal findings in immunity indicators. Elimination of the exposure leads to a prompt normalization of the findings (Pfeifer, J., 1983).

Deaths and injuries due to road-crash involvement are a major health and safety problem among youth. Alcohol is one of a number of factors involved in the overrepresentation of young drivers in road crashes. Frequent and heavy alcohol consumption among teenagers and young adults is not unusual, although they are less likely than older age groups to drive after drinking. Nonetheless, those young people who drive after drinking have a greater risk of crash involvement than older drinking drivers at all blood alcohol concentrations (Mayhew, D.R., 1986).

How Young People Are Mistreated and Held in Unhealthy Environments

Most young people who find themselves in unhealthy environments have little, if any, power of their own to change the situation. Before birth, in early childhood, within the home, and at school, the kind of adult support available and the nature of the physical environment, are in most, if not all cases, non-negotiable.

Young people are sometimes victims of acts of physical, sexual or emotional abuse by adults, often by people they know and live closely with. Child abuse is any act by an adult, or the omission of any act, which results in harm to a child. Physical abuse involves any kind of injury or extreme

punishment of a child and also includes the failure to provide a child with the food, clothing, shelter, and health care needed for the child's optimum development. Sexual abuse is the exploitation of a child by an adult for sexual gratification and includes incest, sexual molestation, sexual assault, and the exploitation of the child for the purposes of pornography or prostitution. Emotional abuse or neglect is the constant ill-treatment of a child through the withholding of affection or through repeated humiliations. (Canadian Home and School & Parent-Teacher Federation, 1986).

Some physical mistreatments are only now beginning to be recognized. For example, children may be adversely affected by trace levels of contamination brought home from their parents' workplaces. A parent's workplace exposure to chlorinated solvents increases the child's risk of developing leukemia. Parental exposures to spray paint, cutting oil, methyl ethyl ketone, and dyes or pigments may increase this risk (Raloff, J., 1987). The use of incense and garden pesticides has also been associated with notably increased risks of leukemia during nursing and pregnancy.

Young people who live with people who smoke often have higher rates of respiratory illness than those in non-smoking households (Bonham, G.S., 1981). This shows up most clearly when the additional chemical burden of living close to a main highway is present (Kasuga, H., 1979). The effect of parental smoking on the pulmonary function of their children is independent of any direct use of cigarettes by the children (Tager, I.B., 1979).

Many young people are adversely affected by their immediate physical environment or by their diet. Environmental health information and accurate information about the possible effects of airborne allergens and diet are only now becoming widely available to adults in Canada, and have still not penetrated school curricula and nutritional training to any great extent (Daglish, S., 1987). Young people often lack awareness even of the possibility they may be impaired by what they breathe or what they eat, and if they become aware, are sometimes given little or no credence or support by family members or health professionals. Their power to make needed lifestyle, environmental or dietary adjustments is more limited than that of adults. Environmentally-related learning or behavioural problems often continue without appropriate diagnosis or treatment, with long-term adverse consequences for the young person in terms of personal relationships, skills development and self-esteem (Rapp, D.J., 1986).

Television has a major impact on children in at least four areas: aggressive behaviour; racial and sexrole stereotypes; decreased interest in reading and school activities; and poorer health habits and attitudes. The modest statistical associations between television viewing and children's cognitive or behavioural problems may be masking the more extreme responses of small groups of especially vulnerable children. The identification of a high-risk subsample of children has been virtually ignored in the research that has been conducted thus far (Zuckerman, D.M., 1985).

Healthy Environments for Canadians: PART II: ANALYSIS

Adults in Canada often do not take young people seriously, commonly referring to them as "kids" and not treating them like people in their own right (Ball, I., 1987). As a result, many young people feel superfluous or unimportant, feel that they don't fit in, and that it doesn't matter what they do. Adults often expect less than full humanness from young people, and do not treat them with respect (Turner, G., 1981).

Adults do have power over young people and can force them to do things (Ball, I., 1987). Many older people still feel that it is acceptable, perhaps even necessary, to treat children in ways that are usually not considered acceptable to treat adults, for example: to be humiliated, insulted, judged constantly, interrogated and compelled to obey (Davies, F.L., 1988). Such attitudes erode a young person's sense of importance, of confidence, and of power. Then when young people become older, they are more inclined to go along with oppressing others, and to give up following their own ideas. It is important that young people maintain pride in themselves and not collude with many false ideas about youth that are pushed at them by older people (Turner, G., 1981).

Adults may underestimate younger people's general environmental competence and ability to choose healthy environments, while at the same time overestimating specific areas of children's performance, for example, in traffic safety (Spencer, C., 1985). In many institutional settings for young people such as schools, control, authority and accompanying surveillance are based on the underlying assumption that in their absence, children will be out of control. The resulting environment (which does not always reflect the goals that teachers, administrators or designers said they were trying to achieve) often creates conformity and teaches children to be passive and powerless, rather than to be active creators of their own lives and experiences (Wolfe, M., 1986). Strong pressure on children through the educational system can eventually cause frustration and aggression (Toronto Star, 1987). This may result in their withdrawing totally from the school system at the earliest opportunity.

Creating Environments That Are Healthier for Young People

Environments that are healthier for young persons allow them to use their capabilities fully, to more actively shape their own lives. To achieve such environments, we must create new ways of designing and planning environments, which will involve children in the process. This involvement, as well as the environments created, will foster the healthy development of children (Wolfe, M., 1986). It is important for older people to remember that young people can have active, participatory roles in society, and make significant contributions to the world's work and well-being (Boulding, E., 1979).

Intellectual learning is facilitated best by allowing children to figure things out on their own, because doing so fosters remembrance and confidence. We should provide young people with acceptance and recognition for having performed an intellectual feat, rather than correct their errors. Our habit of correcting young people can make them self-conscious of their learning, and they may begin to see the world as a place of danger, from which they must protect themselves (Holt, J., 1987).

There is a growing community of people in Canada who view as desirable the full development of a child's capacity for independent reflection, judgment, decision-making, and action. It cannot be fostered in an atmosphere of coercion, and children suffer greatly from lack of respect and autonomy in schools. A non-compulsory, more broadly-based system of community education is proposed in which people of all ages would participate in an on-going, self-generated process (Priesnitz, W.K., 1987).

Young people's physical and emotional dependence on adults around them makes it very difficult for a young person to interrupt abusive adult behaviour, without considerable help from outside the household. The healthiest environment is one in which adults are not abusive; much work is required to determine how this circumstance can be brought about. In the meantime, a safer, healthier environment for young people would give them more support and more power to escape abuse. Agencies involved in aid to children have a primary responsibility to protect young people; public and professional awareness of all forms of child abuse is essential and must be encouraged, to increase the number of allies (and therefore personal power and control) available to children who are abused (Robinson, A.M., 1983).

To determine what environments are healthy for young people, an abundance of research involving children will be required. Those who conduct or sponsor such research must protect children from harm by limiting the risk to which they may be exposed as research subjects, and children should be empowered, as soon as they are capable of some degree of understanding, to decline participation in research if they so choose. The exclusion of children from the process of consent to involvement in research is one aspect of their more general condition of dependence on adults who are responsible for their care. (National Commission for the Protection of Human Subjects of Biomedical and Behavioral Research, 1977).

Research into child abuse may yield answers which may bear on other oppressive and unhealthy environments. Environmental factors and diet may be important (for example, by leading to hyperactivity in a child, whose behaviour in turn stresses an adult's ability to cope beyond its limit, or by impairing the adult's behaviour). Long term studies of abuse victims are necessary, including means of remediating adverse effects (Robinson, A.M., 1983).

The Devaluation of Older People

Like younger people, people who are elderly are often excluded from many aspects of life. They may encounter severe legal and economic handicaps, and social prejudices, that make it more difficult for them to continue to make important contributions to society (Boulding, E., 1979).

Seniors in Canada are starting to organize to help overcome the barriers that separate them from others, and to help governments and others understand their perspectives on current issues, including health, housing, economic and other pressing social policy issues. One Voice - The

Canadian Seniors Network is a newly-formed national, non-profit organization which operates under the belief that Canadians' attitudes towards aging can be changed for the better (Holland, J., 1987). Seniors are focussing on such issues as maintaining good health to increase the quality of life in the later years, maintaining independence in the home, creating change within the system, and developing a supportive social network (Hall, C., 1987).

Characteristics Especially Associated With Aging

Many elderly people are less able to adapt to changes in the surrounding environment than are younger people, and they have more difficulty coping with stress (Adelman, R.C., 1979). Place and possessions have a special significance for the elderly, providing continuity with the past as well as privacy. Many physical changes also accompany aging, such as hearing losses, visual changes, reductions in motor skill, changes in memory and learning, and increased illness (Drader, D., 1982). Elderly people who have smoked previously are even more likely to report poorer health and permanent disability (Hirdes, J.P., 1987).

Because of differences in coordination, balance, muscle and bone strength and other factors, falls are more common occurrences for elderly people than for younger adults, and they are more dangerous when they do occur, often leading to prolonged bed care, complications and sometimes death (Gordon, M., 1982). Chronic care environments involve special attention to handrails and grab bars for additional support (Canada Mortgage and Housing Corporation, 1983). Environmental factors such as indoor air quality can also combine with human factors and building design, to increase the risk of falling accidents (Small, B.M., 1982). The reader is also referred to a more extensive literature on accidents and the elderly: (Calabrese, E.J., 1979) (Blumenthal, M.D., 1980) (Prudham, D., 1981).

The elderly are considered a high-risk group in terms of sensitivity to indoor climatic conditions, including the effects of indoor pollutants such as carbon monoxide (Calabrese, E.J., 1978). There is concern among health specialists about a possible relationship between respiratory diseases and housing conditions such as reduced ambient air temperatures and ventilation rates. Indoor airborne viruses and bacteria are important causes of disabling illness (World Health Organization, 1984).

Healthy Environments for People Who are Older

Elderly people, like any other human beings, need above all to be treated as people of value to those around them and to society in general. Aging itself is a complex issue involving many special needs, and it is tempting to let solutions revolve around these characteristics alone. But truly healthy environments for older people cannot be achieved without consideration of the many other aspects of an aging person's life, including his or her values, language, culture and interests. It is important, for example, when considering health and social services for older people, and when researching the needs of the elderly, to look beyond the mainstream white

middle class (Lee, A.J., 1987). Aging gay people, as a specific case, have pointed out that they encounter a system that does not presently allow for lifelong same-gender lovers sharing facilities in an institutional setting (Hall, C., 1987).

For many seniors, some form of illness is a fact of life, and a cure is not always possible. Health management counselling may be required, and seniors should be encouraged to accept responsibility for themselves to the fullest extent possible, including responsibility for creating a lifestyle that is beneficial to their own health. Ideally health promotion education should be directed at the entire population. It should begin well before retirement, and should be presented in a manner that is relevant to today's seniors. A strategy for a healthy lifestyle for seniors would include: ways and means of preventing further illness; nutrition; exercise and activities; wise use of medications and alcohol; and encouragement to become socially active as volunteers and participate in one's community (St. Lawrence, I., 1986).

A substantial number of older people living in rural areas experience the compound problems of aging and coping with physical disabilities, yet they continue to reside independently or with spouses in the community. For these individuals, the social and physical environment plays a vital role in maintaining the balance between their needs for autonomy and their needs for security. Settings perceived as important in their lives vary widely and include many nonservice settings such as homes and social groups (Norris-Baker, C., 1985).

A healthy environment for an aging person whose personal capabilities are diminishing, is one in which the environmental characteristics are well fit to his or her growing needs, allowing the person to maintain independence and personal control in his or her life. If environmental characteristics in residential settings restrict their behaviour, elderly people are less likely to exercise control and independence, which may be vital to maintaining their health (Kalymun, M., 1985).

It is also important for people to have security, continuity, identity and stimulation in their lives. Environmental design that enables the elderly to more adequately meet these needs can be found in a variety of housing alternatives (Combs, E.R., 1985). Designs have been developed for housing units which accommodate health and mobility limitations, ensure affordability and ease of access to public facilities, allow for socialization, communication, and interaction, and provide a sense of home for residents (Canada Mortgage and Housing Corporation, 1983).

The elderly, living largely on low and fixed incomes, share certain housing and support needs with other low-income individuals, such as students. Imaginative options involving both groups may be available which would help elderly people remain in their current homes or neighbourhoods, achieving affordable, age-integrated housing that is secure and supportive of special needs (Wilde, V.L., 1985).

If older people move from homes in the community to congregate housing, their new housing environment may assume the role of the homes they left behind in the community. Older residents of congregate housing tend to personalize their apartments, and should be encouraged to do so, since residents with more decorative possessions tend to be more satisfied with their apartments. Personalization brings a sense of control, and reinforces self-identity while communicating values to others, enabling social ties to develop (Kinney, J.M., 1985). New, improved settings for elderly people have been shown to foster more favorable attitudes, better mental health and social life, diminished interpersonal frictions, greater emotional independence, and fewer health complaints (Carp, F.M., 1967).

2.2.2 Gender

It is abundantly clear that women have historically been discriminated against and exploited in many ways (Innes de Neufville, J., 1981). Despite many changes and improvements, the undervaluing or devaluing of women continues in Canada today. This is, in itself, an unhealthy environment for women. The devaluing of women has also had numerous direct and indirect effects on women's mental and physical health.

Many women are stuck in unhealthy environments, as a result of continuing prejudicial treatment, based purely on their gender. There is no evidence concerning the nature of women, compared to men, which would justify the kind of differential treatment that exists. Rather, we are dealing with an institutionalized oppression, which is historically rooted, but which now requires change.

How Women are Exposed to Unhealthy Environments

One of the unhealthiest environments women are subject to is economic exploitation; many women in Canada are living below the poverty line. Full equality between men and women will require equal pay, equal opportunity, equal valuing of the work of both sexes, equal access to all levels of employment, and equal distribution of all forms of work, including men and women sharing labour such as parenting and housework, which has been seen traditionally as belonging only to females (Balser, D., 1987).

As a result of their segregation into a few occupational categories, large numbers of women in the paid workforce are exposed to many types of conditions which can prove unhealthy to those who are sensitive to them (Gregory, J., 1981). These include clerical workers who are exposed to long hours at video display terminals, poor lighting, excessive noise, toxic substances, and poor ventilation, as well as poorly designed furniture (Stotsky, K., 1987).

Women in retail and service jobs often face chemical exposures such as dry-cleaning fumes or formaldehyde from freshly-unpacked synthetic clothing, as well as physical stresses such as bending, lifting and carrying. Hair-dressers working around tonics, dyes, pungent chemicals, and aerosol sprays on a daily basis are susceptible to respiratory problems and skin conditions. Teachers, child care workers and nurses are exposed to a variety of communicable diseases, often exacerbated by poorly ventilated buildings. Health care workers are also exposed to radiation, anaesthetic gases, and various toxic substances (Stotsky, K., 1987).

Regardless of their place of work, stress is a common problem for most women workers. Factors causing stress include heavy workload, lower pay than men for equal work, little job control, lack of recognition, monotonous work, unrealistic deadlines, and the added burden of home responsibilities (Davidson, M.J., 1980)(Stotsky, K., 1987). Many women also find sexism and difficulties with partners to be particular sources of stress (Clark, E.J., 1986).

The majority of women, young and old, learn to live every day with the fear of physical and emotional violence (Balser, D., 1987). Many have been beaten, or are still being beaten, by fathers, brothers, husbands and lovers. Many have been victims of sexual assault, and some are still being sexually abused on a regular basis. Many are regular victims of emotional abuse. Almost 1 million Canadian women from all walks of life have been abused (McLeod, L., 1987). The remainder fear it could happen to them next.

Battered women report a significantly higher level of physical complaints, anxiety, and depression than other women, and are at higher risk of developing pronounced mental health problems (Jaffe, P., 1986). Abused women may also feel isolated, and some attempt suicide. For their children it means terror, pain, behaviour problems, the likelihood that the cycle of violence will be repeated in their future families, and a higher risk of becoming violent outside the family. The typical battered woman is trapped in a cycle of poverty from which there has traditionally been little chance of escape (McLeod, L., 1987).

As health care consumers, women encounter medical procedures that they sometimes find degrading, and many have become concerned that some of the treatments, lab tests and surgical procedures they are advised to undergo are unnecessary and/or dangerous. Doctors may be unaware of negative feelings toward female patients, since medicine's condescending attitude toward women has been thoroughly institutionalized (Mendelsohn, R.S., 1981).

The majority of those using mental health services in Canada are women. However, relatively little attention has been paid to their special or unique needs (Canadian Mental Health Association, 1987). The incidence of depression in women is at least two times as high as in men, yet there is very little discussion in the psychoanalytic or psychiatric literature which addresses itself to this gender difference in any meaningful way (Symonds, A., 1986). During their childhood, women are often rejected because of gender, demeaned, trained to serve others, and taught to repress their own needs. They have been raised in an emotional climate which leads to low self-esteem, insecurity, and depression, and this produces lifelong effects in many women.

Women are living increasingly longer than men, yet older women report more acute and chronic illness and disability than men. They are disproportionately represented in nursing homes, since many women are alone; 25% aged 70 or over have no living children and over 60% of older women are widowed, divorced, or single. Older women have fewer personal financial resources for health care than men. They face age and sex discrimination on the part of many health care providers and are subject to a growing tendency to be seen as a burden to the health care system (Lewis, M., 1985).

Healthier Environments for Women in Canada

The fundamental key to achieving healthier environments for women in Canada is the complete elimination of sexism, and the recognition of women as a decisive force in determining Canada's future. Dismantling sexism need not require increased tension between men and women, if it is recognized during

the process that the fundamental cause of sexism is the longstanding institutionalization of oppressive relationships in society, and not the particular individuals today who have been conditioned to perpetuate it. Men are inherently good human beings, and are fully capable of eliminating sexist behaviour (Balser, D., 1987).

Actions directed to improving women's mental health need to be coordinated with efforts to improve women's social status. Significant improvements in women's health and well-being cannot be expected without general gains for women in education, employment, and representation in decision-making roles. Women should also be active participants in the process of developing mental health programs (Canadian Mental Health Association, 1987).

Specific actions which would provide a healthier environment for women in the short-term include promoting the awareness of sexual harassment, increased funding for rape crisis centres and shelters for battered women, the development of more comprehensive standards for occupational health and safety, assisting more women to seek training in the field of mental health planning and policy development, support of initiatives to improve women's social and economic status, and child care (Canadian Mental Health Association, 1987).

A major step forward in improving women's health would result from changing the conditions in our society which perpetuate violence, and particularly violence against women (Balser, D., 1987).

Many women now have two roles; they work outside the home as well as taking responsibility for domestic life. The public services that might help women manage are not given priority by suburbs, which seem to operate under the myth that all women are housewives. Higher density, mixed-use residential environments in city suburbs would give women more options, allowing an integrated life of domestic work, productive work, and leisure (Saegert, S., 1980).

Policy-makers have been slow to recognize the nature and extent of the changes that have taken place in the economic role of women. Policies and programs based on an outdated perception of "the family" and of the relationship between family life and work may be hampering a smooth transition to a new social structure. New patterns of worktime arrangements must make sure that women have access to a fair share of meaningful, dignifying paid work. We must value alternative forms of work such as family responsibilities and voluntarism, but also recognize that for too long women have been engaged in these activities without recognition or pay in addition to their paid employment (Townson, M., 1986).

Early training and environment for females also will require considerable rethinking. Girls and boys receive multiple reminders from adults and from peers as to what are and are not suitable activities for them. It seems that girls are practicing and being prepared for roles in the home, and boys for roles outside. Girls are not encouraged as much as boys, to freely

The majority of women, young and old, learn to live every day with the fear of physical and emotional violence (Balser, D., 1987). Many have been beaten, or are still being beaten, by fathers, brothers, husbands and lovers. Many have been victims of sexual assault, and some are still being sexually abused on a regular basis. Many are regular victims of emotional abuse. Almost 1 million Canadian women from all walks of life have been abused (McLeod, L., 1987). The remainder fear it could happen to them next.

Battered women report a significantly higher level of physical complaints, anxiety, and depression than other women, and are at higher risk of developing pronounced mental health problems (Jaffe, P., 1986). Abused women may also feel isolated, and some attempt suicide. For their children it means terror, pain, behaviour problems, the likelihood that the cycle of violence will be repeated in their future families, and a higher risk of becoming violent outside the family. The typical battered woman is trapped in a cycle of poverty from which there has traditionally been little chance of escape (McLeod, L., 1987).

As health care consumers, women encounter medical procedures that they sometimes find degrading, and many have become concerned that some of the treatments, lab tests and surgical procedures they are advised to undergo are unnecessary and/or dangerous. Doctors may be unaware of negative feelings toward female patients, since medicine's condescending attitude toward women has been thoroughly institutionalized (Mendelsohn, R.S., 1981).

The majority of those using mental health services in Canada are women. However, relatively little attention has been paid to their special or unique needs (Canadian Mental Health Association, 1987). The incidence of depression in women is at least two times as high as in men, yet there is very little discussion in the psychoanalytic or psychiatric literature which addresses itself to this gender difference in any meaningful way (Symonds, A., 1986). During their childhood, women are often rejected because of gender, demeaned, trained to serve others, and taught to repress their own needs. They have been raised in an emotional climate which leads to low self-esteem, insecurity, and depression, and this produces lifelong effects in many women.

Women are living increasingly longer than men, yet older women report more acute and chronic illness and disability than men. They are disproportionately represented in nursing homes, since many women are alone; 25% aged 70 or over have no living children and over 60% of older women are widowed, divorced, or single. Older women have fewer personal financial resources for health care than men. They face age and sex discrimination on the part of many health care providers and are subject to a growing tendency to be seen as a burden to the health care system (Lewis, M., 1985).

Healthier Environments for Women in Canada

The fundamental key to achieving healthier environments for women in Canada is the complete elimination of sexism, and the recognition of women as a decisive force in determining Canada's future. Dismantling sexism need not require increased tension between men and women, if it is recognized during

the process that the fundamental cause of sexism is the longstanding institutionalization of oppressive relationships in society, and not the particular individuals today who have been conditioned to perpetuate it. Men are inherently good human beings, and are fully capable of eliminating sexist behaviour (Balser, D., 1987).

Actions directed to improving women's mental health need to be coordinated with efforts to improve women's social status. Significant improvements in women's health and well-being cannot be expected without general gains for women in education, employment, and representation in decision-making roles. Women should also be active participants in the process of developing mental health programs (Canadian Mental Health Association, 1987).

Specific actions which would provide a healthier environment for women in the short-term include promoting the awareness of sexual harassment, increased funding for rape crisis centres and shelters for battered women, the development of more comprehensive standards for occupational health and safety, assisting more women to seek training in the field of mental health planning and policy development, support of initiatives to improve women's social and economic status, and child care (Canadian Mental Health Association, 1987).

A major step forward in improving women's health would result from changing the conditions in our society which perpetuate violence, and particularly violence against women (Balser, D., 1987).

Many women now have two roles; they work outside the home as well as taking responsibility for domestic life. The public services that might help women manage are not given priority by suburbs, which seem to operate under the myth that all women are housewives. Higher density, mixed-use residential environments in city suburbs would give women more options, allowing an integrated life of domestic work, productive work, and leisure (Saegert, S., 1980).

Policy-makers have been slow to recognize the nature and extent of the changes that have taken place in the economic role of women. Policies and programs based on an outdated perception of "the family" and of the relationship between family life and work may be hampering a smooth transition to a new social structure. New patterns of worktime arrangements must make sure that women have access to a fair share of meaningful, dignifying paid work. We must value alternative forms of work such as family responsibilities and voluntarism, but also recognize that for too long women have been engaged in these activities without recognition or pay in addition to their paid employment (Townson, M., 1986).

Early training and environment for females also will require considerable rethinking. Girls and boys receive multiple reminders from adults and from peers as to what are and are not suitable activities for them. It seems that girls are practicing and being prepared for roles in the home, and boys for roles outside. Girls are not encouraged as much as boys, to freely

manipulate the environment, and this can limit their spatial abilities and other problem-solving abilities. It is important that all children have the freedom, ability, and confidence to define and carry out their own goals (Hart, D.R., undated).

Unhealthy Environments for Men

Male stereotyping (the 'macho' breadwinning superman image) has led men to both devalue and exert power and control over women, often forcing women into situations that are in many ways unhealthy for them. Men's sexist actions and attitudes in turn create an unhealthy environment for themselves, (though they experience the drawbacks from a position of relative power and privilege). These attitudes cut off a great deal of the closeness, companionship, understanding, intelligence, co-operation and mutual support that can be derived from equal relationships with women. At the same time, men often feel restricted from enjoying the full range of human emotions and skills they are capable of, and from establishing close relationships with each other.

Men are also dehumanized, particularly by being treated as expendable economic and military resources. Exhaustion, poisoning, manipulation, and physical exploitation are common conditions for many working class men in Western society. Men are taught to feel ashamed of protesting and are encouraged to think that endurance is manly. They are discouraged from saying 'this hurts' or 'this is not right' and from exercising power to change unhealthy conditions (Kreiner, C., 1986).

Accepting this role is part of the stereotype, which includes being strong, aggressive, capable of violence, unfeeling, sexually compulsive, oppressive to others, and less able to establish close relationships than women. The threat of being called 'gay' or 'not a man' is used, from an early age, to enforce the achievement of this male image. The elimination of the pressure to emulate the macho stereotype must proceed in parallel with the elimination of sexist behaviour (oppression of women by men), for either process to be successful (Irwin, J., 1987).

Healthier Environments for Men in Canada

Recognition of the full diversity of behaviour that men are capable of, and acceptance of men exploring that full range, would constitute a much healthier environment for males (and for females) in Canada than the present. The steadily increasing involvement of fathers with their children is a positive sign of change in a healthy direction. A significant number of male workers are beginning to restore the balance in their family responsibilities and parenting relationships (Couchman, R., 1986).

Recognizing the rights of women to be equal partners, not only in domestic relationships, but in all other aspects of life, is seen as an important step in major world issues such as the elimination of nuclear arms and the threat of global war (Balser, D., 1987).

2.2.3 Family Type

Gradually changing lifestyles in Canada are creating more diverse forms of households and family types. Single-parent families are common, two-parent families where both parents work are common, and the traditional two-parent family with father-breadwinner and mother-housewife is getting more and more rare. There are both young and old couples living without children. There are young and old singles, living by themselves, or in shared households with other singles or families. There are also gay and lesbian households, with and without children. There are many children who have only one parent, or who split their time between two parents who live separately.

Each family environment may have both its healthy and its unhealthy aspects. Each is also affected by other conditions, such as inequitable rates of pay for women, which make it extremely difficult for single female parents to raise a family, or the availability of good day care for young children, which would make it possible for more than one adult in a household to work outside the home. The price and availability of goods and services, such as housing or transportation, may in turn influence the type of household arrangements that are formed.

The proportion of Canadians in poverty was 14.9 percent in 1986. The percentage of mother-led households among those below poverty-level incomes has increased from 9.9% to 12.7% since 1979. The poverty rate among single-parent families headed by women has risen from 55.4% to 56.1%, while the number of families has continued to grow (Shifrin, L., 1987).

A recent report illustrates the kind of hazards that may be inherent in a single-parent situation with insufficient money, and how other external conditions such as housing shortages make the situation worse: A single mother with an employment income of \$800 a month pays almost half of her income to live, with her two teenaged sons, in a tiny, poorly maintained, second-floor, one-bedroom apartment in Toronto. She recently broke some ribs falling down a steep staircase that has no protective rails, and one son's asthma is aggravated by poor ventilation. The basic services like heat and plumbing are inadequate. While municipal officials are aware of this and many similar rental properties, they fear pressuring landlords too hard in case scarce accommodation will be taken off the market. The tenant has been waiting more than a year for subsidized housing, but is relatively low on the priority points system that ranks the 6,000 families now on the waiting list and feels that she would be in a better situation if still on welfare assistance (English, K., 1987).

The struggle to correct the imbalance in work loads between husbands and wives in two-adult families continues. Women working full time often assume an imbalanced workload, involving work outside the home as well as most of the household tasks. An increasing number of men are beginning to take on more family responsibilities, by participating in part-time work, job sharing, and flexible work-time arrangements (Couchman, R., 1986).

Finding appropriate day-care for children has been a nightmare for many families, and there are many potential consequences of inadequate arrangements. For example, when a child's eligibility for a program is tied to his or her mother's participation in a training program or a particular job, frequent damage to the child's sense of stability and security can result from subsequent changes. Child care services that are child-centred, rather than focused on the needs of adults in the labour force, would provide a more stable environment (Bourne, P.G., 1971).

Up-to-date information is required about the diversity of ways in which people in Canada group themselves together in households for mutual support. The foregoing section is only sufficient to briefly illustrate the kinds of problems which may be discovered. What is needed now is direct contact with, and participation by, people in many different family and household situations, in order to obtain an accurate picture of environmental needs and problems from the point of view of family or household type.

2.2.4 Race and Culture

Canada's multicultural nature allows a wide variety of environments responding to the specific nature and origins of its people. At the same time, many who are not the same race, religion, or cultural background as Canada's dominant population groups have sometimes experienced lingering prejudice, which has had the effect of imposing a less healthy environment upon them than should otherwise be possible in as resource-rich a country as this.

Discriminatory conduct (for example, restricting employment, education or housing opportunities on the basis of race or ethnic origin) itself causes emotional suffering for those who are the object of it (Griffith, E.E.H., 1986). Its secondary effects may impose lower economic status, poor nutrition, unhealthy working conditions, inferior housing, high pollution levels, inadequate education, and other less-than-healthy conditions upon its victims.

Examples of Mistreatment of Cultural and Racial Minorities

Some community colleges discriminate against ethnic minority groups in Canada by severely restricting access to the training and education they need. Problems with admission requirements, lack of bridging programs, failure to acknowledge skills obtained in other countries, too few courses combining learning English with studying a trade, little attempt to reach out to minority groups and invite them to take advantage of educational programs, lack of child-care facilities, and insensitivity of college staff and course content to cultural diversity, have all contributed to decreasing education opportunities for people whose race or culture is in the minority in Canada (Contenta, S., 1987).

Civil suits are pending, even in 1987, against a major urban police force for alleged racism, brutality and false arrest. Misunderstanding of cultural differences, for example, ignorance of what might constitute an offensive remark or action, also contributes to feelings of tension between people of ethnic minority background and law enforcement officials or other authorities (Farber, M., 1987).

Canadian Native people living on reserves have higher morbidity and mortality rates than the rest of Canada's population. Native people who migrated to urban centres have high rates of mental health problems, specific diseases, injuries, infant death and hospital admissions. The low socio-economic status, cultural differences, and discrimination that they find in cities are identified as the primary blocks to good health and adequate health care (Shah, C.P., 1985).

The Union of Ontario Indians has charged that the attitudes taken by the federal government and certain corporations towards the Ojibway people (Anishinabek) and the pollution of their land, have contributed to an unhealthy environment for Ontario Native people (Roy, A.W., 1987). A sulphuric acid factory built in the 1950s within the Serpent River Indian Reserve was

abandoned in 1963 for economic reasons, leaving a toxic waste dump containing 120,000 tons of sulphur, calcine and pyrite, which eventually contaminated soils over 100 acres of the reserve (Stokes, P.M., 1985). (The site is only one of several pollution concerns on the Reserve. The Serpent River itself is contaminated with radioactive material from Elliot Lake, and the Aird Bay fishery has been tainted by pulp and paper mill effluent from Espanola.) The Union spokesman suggests that the original decision to establish such an environmentally deleterious operation in that location was indicative of unhealthy attitudes towards Ontario's aboriginal people, and speculates that if the same situation existed in Southern Ontario, with a constituency of the dominant society middle class Ontarians, the site would have been rehabilitated before 30 years had passed.

Healthy Environments for People from non-Majority Cultures

The healthiest environment for people who happen to be of a race, religion, ethnic or cultural background different from the dominant groups in Canada, is to be treated as first class citizens, in every way welcome to enjoy life and privileges in Canada on a par with everyone else. This includes the choice of living among, working with, and participating alongside everyone else, regardless of origin (Winkel, G.H., 1974).

Every opportunity needs to be taken to directly involve the people who will be affected by an environment, in its design. The unique needs or viewpoints of people in non-majority cultures must be taken into account, if the environment is to be a healthy one for them. For example, the Regina Health Department has hired native Canadian health workers, to help motivate native people to avail themselves of the department's programs, to adopt healthy lifestyles, and to facilitate liaison with the native community and organizations (Hutchison, P.A., 1986).

Cultural differences can create barriers to the use of health care facilities (Shah, C.P., 1985). Some immigrant families find that the social organization and ideology of Canadian health care services are not compatible with their own customs for managing illness. This often leads to non-compliance and ineffective treatment. Practitioners need to include patients and families in decisions that affect their lives, and to negotiate culturally acceptable care with them (Anderson, J.M., 1986).

2.2.5 Illness and Handicap

Many Canadians experience some form of illness or handicap which makes their environmental requirements slightly different from those of others. Sometimes, those with illness or handicap have experienced either overt or subtle discrimination, which leaves them feeling both devalued, and stuck in situations which are unhealthy for them.

An accurate understanding of the physical or mental capabilities and needs of people who are handicapped, by those who are at least temporarily able-bodied, would go a long way towards making healthy environments possible for everyone. An attitude of valuing people who may happen to have either temporary or permanent physical or mental disabilities, would also provide a healthier environment for both the handicapped and the able-bodied.

Variations in Vulnerability to Environmental Factors

It is becoming increasingly clear that there are wide variations among the population in the ability to withstand pollution exposures without impairment. For example, lung function was significantly affected by ambient air pollution in only a portion of a studied population (Kagawa, J., 1980). People who are asthmatics respond differently to sulphur dioxide exposures than non-asthmatics (Witek, T.J., 1984). Exposures to ozone can also produce adverse responses in some asthmatics (Silverman, F., 1979). Nutritional status can also modify the toxicity of environmental pollutants, including that of pesticides and heavy metals (Shakman, R.A., 1974) (Mahaffey, K.R., 1979). Exposure to various pollutants in combination may lead to hazardous impairment of performance in older subjects (Ettema, J.H., 1975).

Children with bronchiolitis, with cystic fibrosis, children who have had asthma during childhood, or who have developed hyperlucency in the radiograph after childhood infection, are all those in whom exposure to industrial irritants or to an inclement environment may be serious. Tobacco smoking also increases the susceptibility of an individual to pollutants (Reid, L.M., 1979).

Indoor air pollution exposures are of concern for populations at higher than average risk, including infants and the elderly, pregnant women and their fetuses, anemic and asthmatic persons, those with compromised cardiovascular or pulmonary function, and siblings of sudden infant death syndrome victims (Ammann, H.M., 1987). When populations with common risk factors under exposure to various pollutants are added up, greater than a quarter of the population is seen to be at elevated risk (Small, B.M., 1984).

While it is clear that not everyone in a given population may be affected adversely by an environmental stressor, it is not clear whether all those affected have any predisposing factors which target them for being the first affected. For example, within the last decade there have been hundreds of reported outbreaks of illness among occupants of new or recently remodelled offices, schools, and other public access buildings. These outbreaks have

characteristically affected a large number of individuals, in some cases up to 30-40% of the building's population (Godish, T., 1984). The diversity of different effects among those who appear vulnerable is also important. In outdoor air pollution episodes, a subgroup of people exposed may experience long-term lung damage, but this subgroup may not overlap with those who experienced the most acute symptoms over the short-term. (Stebbing, J.H., Jr., 1979).

The identification and the quantification of the numbers of individuals at high risk in the population is still in its rudimentary stages. This leaves in doubt the percentage of the population that is actually being protected from the toxic or carcinogenic activities of a pollutant, by clean air standards (Calabrese, E.J., 1978).

There are a variety of syndromes which lead to hypersusceptibility, including exposure to high levels of chemicals, nutritional deficiency, and genetic predisposal to environmental chemical sensitivities. In setting standards for environmental contaminants, there must be the awareness that high risk groups are not a small portion of the population, but include virtually everyone from time to time, due to differences in susceptibility for each pollutant, nutritional factors, etc. (Plumlee, L., 1979).

Unhealthy Situations for People With Physical Illness or Handicap

Inadequate knowledge, or warnings, concerning products which may adversely affect a vulnerable portion of the population, present a health hazard to particularly sensitive individuals. For example, the in-home use of paint removers containing methylene chloride results in the absorption of this solvent, which is metabolized to carbon monoxide. Exposure for two to three hours can result in the elevation of carboxyhemoglobin (COHb) in the blood to levels that stress the cardiovascular system. The metabolic formation of COHb continues following the paint-remover exposure, doubling the duration of the cardiovascular stress produced by a comparable COHb level after exposure to CO. Patients with diseased cardiovascular systems may not be able to tolerate this unexpected stress (Stewart, R.D., 1976).

For those who are ill or handicapped, it can be a continual battle to limit the amount of impairment, because many environmental conditions may present hazards to them, that are not as harmful to the general population (Durlak, E., 1987). For example, air pollution can aggravate existing disease conditions or put at higher risk those who are predisposed to ill health (Severs, R.K., 1980). Even very low levels of nitrogen dioxide can cause a significant increase in specific airway resistance for an asthmatic person (Orehek, J., 1976).

People with physical handicaps may also be at considerably increased risk of acquiring further disability, or dying, from environmental conditions that are not recognized as dangerous by temporarily able-bodied people. For example, a recent inquest into the death of a disabled woman was told that there were many things wrong with the highrise where she had lived. The legless widow died as a result of infection from injuries due to falls. Only

two of four main entrances were usable by those in wheelchairs, while one ramp was dangerously steep, too narrow, and lacked handrails, according to a study of the building. There were many more access-oriented problems in the building, which was originally not designed with the disabled in mind (Sutton, R., 1987).

Non-ambulant and semi-ambulant people are at particular risk in emergencies such as building fires. Knowledge of techniques of preparing, lifting, and carrying persons with various disabilities is required in emergency evacuations of buildings, as it is often difficult for untrained rescuers to motivate certain people to evacuate (Johnson, B., 1983).

When a child with severe chronic mobility problems becomes too heavy or too cumbersome to lift and to carry, parents must consider removing architectural barriers in the home. Practical and psychological difficulties frequently cause them to postpone making such changes until a crisis occurs. Practical difficulties include lack of resources, gaps in services, and a general disregard by medical personnel of how environments affect behavior. Psychological difficulties, which have so far received little attention, mainly stem from conflict between the parents' desire for an idealized house and their child's functional requirements, as well as from the parents' desire to appear as normal as possible to their neighbors. Adapting the home means accepting the permanence of the disability, and making the family's disability public — stigmatizing the house, and thus its occupants (Lewis, B.E., 1985). For further discussion about building accessibility for the handicapped, the reader is referred to (Margulis, S.T., 1981) and to the report index.

New environmental hazards are also being discovered within common household environments, for people who have impaired or suppressed immune systems. For example, types of fungus that once only affected foods such as tomatoes and peaches are turning up as the cause of some diseases in immune-suppressed individuals (Dunlop, M., 1987).

There is also a complex relationship between stress and other emotional factors, and physical disease. For example, it is known that emotional factors can trigger or worsen asthma in a physiologically or biochemically predisposed individual. The disease process also has a very definite emotional toll on the individual with asthma, as well as on other members of the family. Achieving an optimum environment and optimum functioning while subject to illness may require complex psychological adaptations, alterations in lifestyle, and in family and personal interactions. Emotional factors and psychological problems which accompany or result from having asthma can be a major impediment to progress toward health (Nieder, J., 1986).

For people who are severely ill, the question also arises as to whether modern technology has made it possible to keep people alive past the point at which life has ceased to be worth living. Some argue that everyone should have the right to control the ending of their own lives (Seguin, M., 1987). What constitutes a 'healthy' environment for those who are dying is not as clearcut a question as it can be at earlier stages of life, yet it may be of concern to everyone in Canada, at some point in their lives.

Unhealthy Situations for People With Mental Illness

Present mental health care systems, while endeavouring within available resources to help and cure people with mental illnesses, do fall short, in many areas, of providing truly health-promoting environments for people who have been categorized as emotionally disturbed. Mental health system consumers have expressed feelings of cynicism and powerlessness resulting from the overwhelming paternalism directed towards them, as clients of the system. Traditional psychiatric labels make it almost impossible for them to be seen as human beings (Hutchison, P., 1985).

Many people who have been mentally ill need independence but also require a great deal of support to become re-integrated into community life. They may periodically need some form of shelter or refuge, but they believe the imposing controls and practices of institutions and hospitals are generally inappropriate. The rehabilitation and welfare systems tend to trap many into a life of poverty and dependence, which result in feelings of low self-worth. Lack of appropriate housing conditions and work opportunities contribute to lack of dignity, income, and security (Hutchison, P., 1985).

Mental health services cannot adequately respond to the needs of the mentally ill without recognizing the harmful effect that a variety of social, environmental, physical, psychological and biological factors can have (Insel, P.M., 1983). For example, some pesticides, heavy metals, pollutants, foods, food additives, and drugs have been linked with unexpected behavioral disturbances, including paranoia, hyperactivity, anxiety, and hallucinations (Philpott, W.H., 1980) (Johnson, K.A., 1982). Careful control of a person's environment has commonly been ignored as a potential benefit in the treatment of mental disorders.

Healthier Environments for People Who Are Mentally Ill

The majority of those using mental health services are women. It would be appropriate to assist more women to seek training in the field of mental health planning and policy development (Canadian Mental Health Association, 1987).

In designing for the developmentally disabled, we also must design for those parts of people or their daily lives that are not disabled. Otherwise, the environments we design, both socially and physically, can help to create more disabilities (Wolfe, M., 1976). The media have portrayed mental illness and the mentally ill in a disparaging manner (Day, D.M., 1986), preventing others from seeing them as human beings. The hopes of people disabled by mental health problems are not unlike those of other Canadians; their use of formal mental health services does not mean that other things in life, like making a contribution to family, friends, work and neighbourhood, are any less important (Hutchinson, P., 1986). The stigma of mental illness can present many problems for the chronic psychiatric patient, for example, reduced job availability (Kirsh, S., 1983). It is important to involve and support individuals with a disability, to be part of the everyday things which many of us take for granted, including community activities.

Often, mentally disabled persons are either segregated into expensive formal health care and rehabilitation systems, or discharged, with limited support, into unprepared communities. Increasing concern has been raised about the fate of the de-institutionalized patient in the community, including reports of inadequate housing, life skills, activity and work preparation programs, as well as staff overload and frustration at lack of resources (Toews, J., 1982). A healthier social environment for mentally disabled persons will require more community investment, so that at some point, they can live with maximum reliance on the resources found or created within their local environment (Pape, B., 1987). Community-based rehabilitation offers a vision of citizenship to the mentally disabled, rather than one of control and exile (McKnight, J.L., 1983) (Trainor, J., 1984).

New initiatives in mental health advocacy have focused on helping former and present users of mental health services participate in planning for a better mental health system (Wolfe M., 1976) (Church, K., 1986). The reader is also referred to additional references for further discussion of environments suitable for those who have been diagnosed as mentally ill: (Stewart, D.W., 1982) (Campbell, D., 1986).

Appropriate Environments for the Chemically Sensitive

Over the past decade, the phenomenon of environmental sensitivity has been receiving increased attention. Sensitized individuals become reactive to a wide variety of low-level air contaminants in their immediate environment (Dickey, L.D., 1976) (Rea, W.J., 1979). Once this process has begun, common indoor and outdoor environments appear to exacerbate symptoms, and chemically less-contaminated environments appear to be therapeutic (Selner, J.C., 1986) (Daglish, S., 1987). There is some indication that the proportion of hypersensitive individuals in the population is increasing. The World Health Organization has recommended that special environmental requirements for the protection of the affected groups should be assessed (World Health Organization, 1984).

One of the most difficult aspects of dealing with hypersensitivity is the prevalence of negative attitudes among the less-sensitive population, towards individuals who find themselves highly reactive to physical environmental factors. In the words of one individual coping with the illness: "The most demoralizing and frustrating experience ... is that, while being seriously ill, weak, exhausted, confused, in pain, and isolated, one must also fight spouse, children, extended family, neighbors, friends, community, all levels of government, doctors, hospitals, social services, OHIP, OMA, psychiatrists, employers, insurance companies, pension and/or welfare agencies — in fact, the whole world — in order to survive" (Hall, M., 1987).

Many seemingly innocent practices present unexpected hazards to people who have become temporarily hypersensitive, including the use of perfumes built into stationery, advertising and periodicals, and the common use of scented cosmetics, body, and clothing care products by both men and women (Hall, M., 1987). Unannounced pesticing of apartment buildings and offices can render a previously safe location totally intolerable.

The present atmosphere of general ignorance about hypersensitivity diseases, and even about allergy, which has been studied for a longer time, is an unhealthy and a dangerous one for both allergic and hypersensitive individuals. It leads to feelings of hopelessness at being misunderstood, and often to extreme isolation, as friends and families withdraw, not knowing how to deal with the environmental needs of the afflicted person (Hall, M., 1987). Health care environments have often proven to be particularly hazardous to the asthmatic and/or chemically sensitive individual seeking emergency treatment (Durlak, E., 1987). For example, many care-givers and hospital workers are unfamiliar with the potential effects on sensitive individuals of volatile chemicals commonly used in hospitals.

The food and chemically sensitive individual will sometimes have difficulty in maintaining a healthy and supportive social network for himself or herself. Some will have childhood histories of learning problems and deficits in appropriate social skills, possibly relating to early malfunction or specific physiological systems. Behaviours commonly exhibited include increased activation, an increased probability of frustration, decreased attention span, perceptual problems, and memory deficits (O'Banion, D.R., 1981). Social skills may be required, however, to negotiate environmental changes that represent, to date, the best means of reducing the individual's hyper-reactivity.

2.2.6 Orientation and Lifestyle

There are adverse mental, physical and social consequences to people who are lesbian, gay, or bisexual, from living in an environment of prejudice against their orientation and lifestyle. The literature is clear that it is the oppressive environment and its consequences, and not the choice of loving someone of the same gender, that is the causative factor in their particular health problems.

Homosexuals are probably the most frequent victims of hate-motivated violence, and are targeted for assault, verbal intimidation, and vandalism more often than blacks, Hispanics, Southeast Asians, and Jews (Finn, P., 1987). The coping strategies of lesbians and gay men in reaction to such oppression in everyday life, and the damages suffered by them, are in many ways similar to those of Jews and blacks in response to anti-Semitism and racism (Adam, B.D., 1978). People survive domination through resistance, accommodation and compliance.

The Devaluation of Lesbians and Gay Men

Gay oppression is an integral part of society's pressure on all men and all women to conform to rigid stereotyped images. We are told, as we grow up, what men and women should be like. If a boy or girl strays from the stereotypes and makes independent choices, he or she faces rejection, sometimes violence, and often the accusation that he or she is gay, "not a man", or "not a woman" (Irwin, J., 1987).

To maintain the threat of prejudice against everyone, overt prejudice and even violence has been directed against individuals who choose to engage in loving relationships with people of the same gender and/or in same-gender sexual practices (Finn, P., 1987). All other human characteristics, skills, goals, and needs of gay-oriented individuals, which they share with heterosexually-oriented individuals, are often completely ignored in favour of focussing attention on the differences in their choices of lifestyle.

How Lesbians, Gay Men and Bisexual People are Mistreated

A homosexual adolescent is often placed at a greater distance from his or her parents than his heterosexual counterpart. For those who reveal their sexual orientation, the family response is often not supportive. In the absence of a support group, the adolescent feels alone and isolated (Paroski, P.A., 1987). Negative reactions of family and peers toward a young person's homosexuality, damaged self-esteem, the desire to be with gay-identified peers, and exposure to street life are important factors predisposing gay men to a higher risk of involvement with prostitution, alcohol abuse and drug abuse. There is also potential for intense family discord, and physical and/or emotional abuse when homosexuality is first revealed (Remafedi, G.J., 1985).

Many men who ultimately decide that they are homosexual or bisexual have at some point in their lives married or established continuing relationships with women. Many of these men have also fathered children. The life of the gay father is often fraught with fears and anxiety involving marriage breakup, access to children, discrimination against himself or his family, and the process of 'coming out' as gay (Heath, L., 1981).

The current epidemic of acquired immune deficiency syndrome (AIDS) has led to many instances of overt discrimination against gays, including social ostracism, and even the deprivation of various rights such as housing, employment, transportation, and funeral services (Douglas, C.J., 1985). It has also highlighted the need for equitable legislation regarding spousal rights and distribution of property upon death, for gay couples (Adam, B.D., 1987).

Groups fighting for gay rights stress that public policies often reflect and reinforce homophobic attitudes, particularly in such areas as education, family law, police/community relations and the administration of health (Coalition for Gay Rights in Ontario, 1986). Other human rights issues for lesbian and gay men and women include child custody, tax laws, municipal zoning regulations, social welfare, insurance and employment. Quebec, Ontario and the Yukon Territory have added "sexual orientation" protection to their human rights codes. A number of municipalities have also instituted limited forms of protection and many unions have won such protection in their contracts. At the same time, some governments, social agencies, businesses, and landlords continue to deny lesbian and gay individuals basic civil and human rights (Kinsman, G., 1987).

Stigmatization of homosexuality has also contributed to the invisibility of this aspect of their lives, for many men and women in Canada who choose a gay, lesbian, or bisexual orientation (Adam, B., 1987). There is an immense cost suffered by those in hiding, in terms of psychological suffocation, and in terms of the fear that public identification of their orientation will invite prejudice and loss of those freedoms, privileges and opportunities normally taken for granted by the heterosexual population.

Since the gay and bisexual population is under these circumstances neither easily identifiable nor enumerable, information-gathering and information dissemination about health and other issues is difficult (Soskolene, C.L., 1986). At the same time, it has been a matter of life and death for gay and bisexual men, as a high-risk group, to obtain accurate information about the transmission of Acquired Immune Deficiency Syndrome (AIDS) (Allen, M., 1987). Prejudices against homosexuality have led to obstacles being presented against distribution of literature about 'safer sex', a term used to describe sexual practices which minimize the risk of transmission of sexually-transmitted diseases such as AIDS (Adam, B.D., 1987). Information about sexual decision-making, personal hygiene, prevention of sexually-transmitted diseases, and substance use, can be life-saving for the young homosexual in particular (Remafedi, G. J., 1985).

How Lesbians, Gay Men and Bisexuals Differ from Heterosexuals

The differences between gay-oriented and non-gay-oriented people are not highly visible. Unless you know someone fairly well, it is difficult to know for sure if he or she is lesbian or gay, bisexual or heterosexual in orientation. Lesbians and gay men are conscious of erotic and affectional attractions to people of the same gender. They may or may not also have attractions to, and sexual relations with, people of the opposite gender, but their preference is for closeness and sexual relations with persons of the same gender (Clark, D., 1987).

Bisexual people have erotic and affectional attractions to people of both genders, and prefer to have, during their lifetime, closeness and/or sexual relations with individuals of both genders. Heterosexual people are conscious of erotic and affectional attractions to people of the opposite gender. They may or may not also have attractions to, and sexual relations with, people of the same gender, but their preference is for closeness and sexual relations with persons of the opposite gender.

With the exception of having a different gender-preference, lesbians, gay men and bisexuals all have the same kinds of needs and desires for everything in life (family, food and nutrition, clean air and water, shelter, clothing, companionship and support, achievement and self-worth, life goals, self-expression, sexuality, parenting, community, communication, information, relaxation, excitement, health care, and so on), as every other human being. At the same time, there is as wide a diversity among lesbians, gay men and bisexuals in Canada as there is among heterosexuals, in their individual natures and their lifestyles.

Difference In Vulnerability to Environmental Factors

The AIDS crisis at first made it appear that homosexuals and bisexual men were more vulnerable than others to infection. The predominance of gay and bisexual men among infected individuals turned out to be due to the virus having been first introduced to North America into the gay population, and to the rapid transmission of the virus through the exchange of bodily fluids via anal intercourse among gay men.

Once the AIDS virus began spreading into the heterosexual population, its transmission through exchange of fluids in heterosexual sexual activity also became apparent. Recent shifts in the nature of sexual activities among homosexual and bisexual men have begun to reduce the risk of transmission of the disease among them, through restriction and modification of those practices which would have transferred bodily fluids, and substitution of "safer sex" practices (Adam, B.D., 1987).

The specific life experiences and the nature of oppressions experienced by lesbians, gay men and bisexuals may have a strong bearing on the suitability or health of particular environments for them, in the absence of any changes in the overall incidence of prejudice in our society. The design of gay-supportive housing, for example, would differ from present

designs (which do not take lesbian and gay orientations into account) since gay individuals do not yet enjoy the same freedoms of association or affection in public spaces without fear of abusive behaviour by non-gay people (Bradley, E., 1987).

A Healthier Environment for Lesbians, Gay Men and Bisexual People

The healthiest environment for those in Canada who choose a lesbian, gay or bisexual orientation and lifestyle is one of acceptance and respect, protected by explicit legislation guaranteeing individual rights and freedoms.

Critical in the termination of unhealthy conditions is the interruption of overt prejudice against homosexuality in Canada. Prejudice breeds on ignorance and inaccurate, stereotypical images of people. Gay oppression has remained intact in Canada because lesbians, gay men, and bisexuals, for reasons of survival, are largely invisible in our society. Non-gay people therefore have little or no accurate information to counter the stereotyped and false images of homosexuality with which they were brought up (Clark, D., 1987).

Protection of the civil rights of lesbians and gay men is a necessary precondition for lesbians and gay men to have equitable access to both civic rights (the rights to the impartial administration of civil and criminal laws in defense of property and person) and political rights (including freedom of speech, of press, of assembly, to petition for the redress of grievances, and to join with and be identified with other persons for common political goals). Exercising all these rights requires public actions, which are impossible for a person who must remain invisible, hidden and secreted, in respect of his or her minority status, as a condition for maintaining his or her livelihood and housing (Mohr, R.D., 1985).

Little attention is given in medical school curricula to providing appropriate health care for the 5-10% of the population whose orientation is homosexual (Bauman, K.A., 1985). Health care providers need to be aware of the sense of isolation, the process of hiding one's homosexuality, the conflicts that homosexual adolescents have regarding their lifestyle, and the need of lesbians and gay men for non-judgmental health care (Paroski, P.A., 1987).

2. Healthy Environment Issues (continued)

2.3 By Environmental Resource and Social Condition

The eight subsections following give a partial sampling of environmental factors, both physical and social, that affect health. Depending on what was available to us in the literature, we have tried to include both the positive and the negative: the environmental resources and social conditions that seem to foster good health, as well as the environmental and social hazards that contribute to ill health.

One message from the literature is that while many of the conditions we discuss may be necessary for good health, none is by itself sufficient. Human beings are very adaptable, but not so adaptable that we can get away with forgetting major needs, whether they be physical, social, or psychological, without paying some price in health and wellbeing.

2.3.1 Safety

Injury is among the most important public health problems, both in terms of numbers of people affected, and in terms of the potential effects, which include permanent disability and death. Preventing injury extends far beyond awareness of precautions which will reduce the risk of accidents within existing environments. It requires that we take a conscious look at selecting the technologies we use, taking potential risk of injury into account. High injury rates are associated with road vehicles, guns, farm machinery, sports equipment, cigarettes, and many other consumer products (Robertson, L.S., 1986).

Children are particularly vulnerable to accidental injuries, often being unfamiliar with the extent of hazard involved in situations adults have learned to live with relatively safely (Garling, T., 1985). Understanding the child's perceptions of hazards in familiar environments may be critical to making better progress both in product design and in safety education (Sheehy, N.P., 1985).

Egocentric thinking on the part of able-bodied adults who have maximum power to structure their environment, may leave other people, such as younger people, the elderly, or the disabled, at more risk. A failure to recognize the different perceptual capabilities, decision processes, motivations, etc. of a particular group can lead to the provision of environments which present no great risks to a healthy adult, but which can be extremely hazardous to others (Stratton, P., 1985).

Careful thinking may be required to better anticipate physical and chemical hazards which may only arise under emergency conditions. For example, an electrical fire in a large commercial building caused oil containing PCBs to spill into the blaze. The resulting toxic gases included dioxin, which is extremely hazardous. Contaminated soot was dispersed throughout the building by the ventilation system (Yulsman, T., 1985).

2.3.2 Clean Air

One important resource which is an indispensable part of a healthy environment is clean air. The literature contains a great deal of discussion of the contamination of air, both indoors and outdoors, by various chemical pollutants. The evidence is very clear that contaminated air can present some risk to the entire population, and a much higher risk to a subset of the population, who appear to be more susceptible than others.

Information about the health effects of low-level contamination of air has not yet been widely distributed, though the effects are well documented. Testing for the effects of contaminated air on people is not yet a widely accepted practice within the health care systems in Canada.

Much of the basic information about the adverse health effects of polluted air, and of the therapeutic and diagnostic role of clean air, has been presented and discussed in earlier sections, from the point of view of a particular setting, such as the home, the school or the workplace. The information below summarizes additional facts about the nature of the contaminants in air, the sources of these contaminants, and their health effects, independent of the setting.

Contaminants in Outdoor Air and Their Effects on Health

Both particulate and gaseous contaminants are now routinely measured in outdoor air at various points across Canada. Pollutants that predominate outdoors include sulphur oxides, ozone and related oxidants, trace metals, halogen compounds, petrochemical compounds, and pesticides (Lebowitz, M.D., 1983). There are many sources for the pollutants found in outdoor air, including industrial activity, transportation, utilities, municipal incineration, home heating, and agricultural activities.

Various studies, in Canada and beyond, have established a link between exposure to common outdoor pollution and increased incidence of health problems. For example, one Ontario study found highly significant associations between excess admissions to hospitals for respiratory disorders, and the levels of sulphur dioxide and ozone, during the months of July and August over several years in the mid-to-late 1970s (Bates, D.V., 1983).

In patients with asthma and hyperreactivity, small concentrations of a pollutant like sulphur dioxide can cause substantial and clinically important respiratory effects. In acute severe episodes of air pollution, predominantly people with pre-existing heart and lung disease show an increase in mortality (Simonsson, B.G., 1986).

There is a close association between air pollution and various respiratory indices in children, and smoke appears to have a greater effect on health than sulphur dioxide. It is likely that initial effects on children's respiratory indices will show up in areas with an annual mean concentration of sulphur dioxide and smoke of 50-200 $\mu\text{g}/\text{m}^3$ (Colley, J.R.T., 1980). Children living in highly polluted areas show a higher prevalence of cough and other

chronic respiratory symptoms, and decreased lung measures, compared to children in less polluted areas (Kerrebijn, K.F., 1975), (Sugita, K., 1981).

The immune system may be considered to be a series of mechanisms for adapting to an ever-changing and hostile environment. Any teratogenic effect early in embryonic life can have profound consequences on the immune system of the developing infant. A number of chemical pollutants are known to injure the immune response of the fully developed organism. The human fetus and newborn are especially susceptible to substances like lead, cadmium, nitrogen dioxide, and sulphur dioxide (Bellanti, J.A., 1974).

There is little data on the health effects of mixtures of pollutants. Some experiments indicate there may be synergistic effects. For example, repeated inhalation of mixtures of nitrogen dioxide and ozone (at relatively high levels) was more effective than the single gases, in reducing resistance of animals to infection (Ehrlich, R., 1977).

Many industrial processes utilizing or producing inorganic materials release metals (usually in the form of oxides or salts) into the environment. Significant quantities of metals are also released from incinerators burning industrial and municipal waste. Metals are also released from combustion sources such as fuel burning for heat and power (including cars) (Yocom, J.E., 1983). Of particular concern is the effect of lead, since lead levels formerly considered safe have recently been shown to increase abnormal behaviour, cause learning difficulties and reduce intelligence. Children can absorb enough lead to impair their performance on tests of reasoning, coordination, intelligence and reading. Some symptoms of lead toxicity are: hyperactivity, perceptual disorders, mental retardation, fatigue, irritability, temper tantrums, learning disabilities, speech disturbances, perceptual motor dysfunctions, and emotional or behavioural problems (Schauss, A., 1980).

Another review emphasizes that particulate air pollution, even at current levels, could be of concern for public health and that new research is needed to assess the effects of acidic aerosols and oxidants involved in the formation of acid precipitation (acid rain), on human health in Canada (Ozkaynak, H., 1985). Health and environmental organizations are also becoming very concerned about potential health effects of municipal projects such as garbage incinerators (Poole, R., 1987).

Various additional researchers report significant effects of inhaled air pollution on health:

- effects of ozone (Jaffe, L.S., 1967).
- effects on body and mind functions (Laverne, A.A., 1970)
- effects of carbon monoxide on mortality (Hexter, A.C., 1971)
- effects on cardiovascular and other systems (Finkel, A.S., 1976)
- effects on height and bones in children (Thielebeule, U., 1980)
- effects on the respiratory tract (Phalen, R.F., 1981)
- effects of low levels of lead intake (King, E., 1982).
- aluminum, iron, & lead in respirable particles (Tosteson, T.D., 1982).

Contaminants in Indoor Air and Their Effects on Health

Urban residents typically spend more than 90 percent of their time indoors. The indoor environment is likely to include exposure to radon daughter nuclides, formaldehyde, carbon monoxide, nitrogen dioxide, respirable particulates and asbestos, tobacco smoke, microorganisms, and aeroallergens, organic vapours, mineral and synthetic fibers (Lebowitz, M.D., 1983), (Spengler, J.D., 1983). Indoor exposure may constitute 80-95% of the total exposure for some pollutants (Dudney, C.S., 1981). Recent investigations have also shown that as much as 10% of the pollutants exhausted from a building may commonly reenter the building rather than be dispersed to the atmosphere (Reible, D.D., 1985).

The health effects of formaldehyde include: irritation to the skin, eye, lung and mucosal surfaces; sensitization; alteration in irritancy threshold; mutagenicity; and potential carcinogenicity (Gupta, K.C., 1982), (Nethercott, J.R., 1982), (Wartew, G.A., 1983). Indoor levels of formaldehyde, due primarily to off-gassing of building materials, may range from less than 0.01 ppm to approximately 3 ppm (Ulsamer, A.G., 1982). Canadian guidelines propose 0.1 ppm as the upper limit in residential situations. Particle board, a common building construction material made of wood shavings held together with a urea-formaldehyde glue, continuously emanates formaldehyde (Andersen, I., 1975). Formaldehyde levels indoors increase with temperature, and decrease with the age of the source (Moschandreas, D., 1986). (See also section 2.1.1 "At Home").

The respirable suspended particulate levels generated by tobacco smoking overwhelm the effects of ventilation and inflict significant air pollution burdens (Repace, J.L., 1980). In rooms with even moderate smoking, a great deal more fresh air is required, even for nonsensitive occupants to judge the comfort level acceptable, than in nonsmoking conditions (Cain, W.S., 1982).

Statistically significant dose-response relationships between exposure to tobacco smoke and loss of life expectancy have been calculated. Even a limited amount of exposure to tobacco smoke carries a greatly increased risk of premature mortality (Repace, J.L., 1981). More than 30,000 Canadians die each year from smoking-related illnesses, including cancer, cardiovascular disease, and pulmonary disease. While overall tobacco use has declined in Canada, a number of disturbing trends persist: smoking onset among 12 to 14 year olds has remained high; the prevalence of smoking among young women (20 to 29) is increasing; and regional and socioeconomic differences in smoking rates continue to exist (National Program to Reduce Tobacco Use, Consultation, Planning and Implementation Committee, 1987).

In the presence of tobacco smoke, many normal nonsmokers experience eye and throat irritation, headache, rhinitis and coughing. Allergic persons report wheezing, sneezing and nausea as well. Particularly acute symptoms may be found in infants, children, persons with cardiovascular or respiratory disease and wearers of contact lenses (Repace, J.L., 1980). Passive exposure to tobacco smoke can increase heart rate, systolic and diastolic blood pressure, and venous carboxyhemoglobin. People who are prone to angina may experience typical symptoms, including a reduction in heart rate and systolic

blood pressure at onset of angina (Aronow, W.S., 1978). Nonsmokers breathing ambient tobacco smoke are at significantly increased risk of illness and mortality than those not exposed (Repace, J.L., 1982).

Houses causing environmental illness problems have higher concentrations in the air of Volatile Organic Compounds (VOC) than houses without problems. Below 0.6 mg/m³ no irritation by VOC can be expected. Above 2 mg/m³ VOC are expected to cause irritation. Above 5 mg/m³ irritation and decreased mental performance may occur (Molhave, L., 1982), (Molhave, L., 1986).

Indoor air, whether in commercial, industrial, or residential buildings, typically contains levels of polychlorinated biphenyls (PCBs) at least 1 order of magnitude higher than outdoor levels. Defective fluorescent light ballasts (from an era in which PCBs were used) can emit PCBs and are an important source of indoor atmospheric contamination (MacLeod, K.E., 1981).

A very broad range of concentrations of radon is observed inside buildings, particularly in homes, and even the average concentration has an estimated risk that is large by comparison with most environmental risks. (Nero, A.V., 1986). Further discussion of the problem of radon gas and radon daughter products in homes is found in section 2.1.1. "At Home". Additional references dealing with radon include: (Jonassen, N., 1981), (Letourneau, E.G., 1985).

Bacterial and viral disease can sometimes be transmitted from person to person on indoor air, particularly when poor ventilation allows indoor concentrations to build up. Devices such as humidifiers may be sites of culturing of bacteria, as can fan coil units (Hambraeus, A., 1986), (Johnson-Lussenburg, C.M., 1986). The most serious consequences on human health are usually observed in especially susceptible populations, but unexpected allergic responses ranging from trivial to life-threatening can also result among the general population (Tobin, R.S., 1986). The growth of *Legionella* bacteria, which can cause life-threatening illness, is amplified in such things as cooling towers, humidifiers, and hot water systems. It sometimes incubates indoors in warm and/or standing water found in humidifiers, shower heads, and air conditioning ductwork (Besch, E.L., 1981).

Indoor air always contains viable mould spores, but in buildings that do not generate their own sources of mould, the numbers are usually low and the species found are similar to those found in outdoor air (Hambraeus, A., 1986). Building materials of high cellulose and low nitrogen content that become moist and are subjected to temperature fluctuations can provide ideal conditions for types of fungal growth that produce macrocyclic trichothecene toxins, which can cause illness and can be immunosuppressive (Jarvis, B.B., 1986). Mycotoxins have been associated in particular with certain types of thermal insulation (Scott, P.M., 1986). In buildings with flat roofs, water damage can lead to subsequent microbial growth, which may be related to complaints about indoor climate (Gravesen, S., 1986).

Sensations of odour and irritation (the effects experienced by people due to stimulation of the olfactory and trigeminal nerves) may be the first warning signals a person receives that the atmosphere is contaminated. While scientific and objective measurement of the environment seems generally

favoured over potentially subjective information received from the people living in the environment, human senses still make a crucial contribution to the analysis of indoor air. It is still the effect of indoor air on human beings, as they experience it, which provides most of the norms for indoor air research (Engen, T., 1986).

One neglected source of both indoor air pollution and personal intake of pollutants is clothing. The additives and contaminants which occur in textile fibers vary widely. For example, synthetic fibers such as nylon and polyester contain trace amounts of contaminants such as catalysts and deactivators which remain after the synthesis of basic polymers. In addition, there are frequently a number of materials which are added to perform specific functions, such as traces of metals or metal salts, and antistatic agents and flame retardants. After the fibers are converted into fabric form, other substances are applied to act as lubricants, sizing agents, antistats, bleaches, wetting agents, dyes, and durable-press treatments (Barker, R.H., 1975), (Finkel, J.M., 1979).

The effects of these agents on human health has not been well studied, although allergic skin reactions, particularly due to formaldehyde in new clothing, have been reported (O'Quinn, Silas E., 1965). Recent research suggests that cardiovascular effects such as premature ventricular contractions may be more frequent in people wearing synthetic clothing, compared to those wearing cotton (Seyal, A.R., 1987).

Additional references dealing in detail with indoor air pollutants and their health effects include: (Bridbord, K., 1975), (Adkinson, N.F., 1977), (Muittari, A., 1978), (Schaumburg, H., 1978) (Andersen, I., 1979), (Molhave, L., 1979), (World Health Organization, 1979), (Hollowell, C.D., 1980), (Beall, J.R., 1981), (Hollowell, C.D., 1981), (Committee on Indoor Pollutants, National Academy of Science, 1982), (Wigle, D.T., 1982), (Rittfeldt, L., 1984).

Effects of Low-Level Contaminants on Sensitive Individuals

Overexposure to commonly used environmental chemicals appears to have been responsible for the development, in some individuals, of recurrent signs and symptoms of inflammatory type diseases. Repeated episodes of illness are triggered by ambient chemical fumes in the outdoor air and home environments. Once an individual is sensitized to a solitary chemical it is apparent that continued exposures result in a spreading phenomenon, involving sensitivity to other types of chemical exposures. Once this spreading occurs, reactions then proceed upon minute exposures (Rea, W.J., 1978). Exposure to environmental pollutants has also been clinically reported to provoke cognitive, emotional, and behavioural reactions in sensitive individuals (King, D.S., 1981).

A period of time in a relatively fume-free and particle-free environment has been used successfully to clear the majority of symptoms and signs in chemically sensitive individuals without the use of medication. Double-blind rechallenge with ambient dose levels of synthetic chemicals is used to reproduce and verify most of the symptomatology (Rea, W.J., 1978).

Additional discussion of chemical hypersensitivity can be found in Section 2.2.5 "Illness and Handicap".

2.3.3 Food and Water

Achieving health for all in Canada cannot be achieved without acknowledging and addressing the reality of Canada's food and water supplies. Food and water are far more complex than Canada's Food Guide would have our children believe. Nutritional needs vary widely from person to person, as do individual sensitivities to different diets and to various deliberate additives as well as inadvertent contaminants in most of Canada's food supplies. Water in some areas is highly contaminated and in others, though treated and deemed safe, may contain hundreds of trace levels of various toxic chemicals, the combined and long-term effects of which, at these levels, are totally unknown.

The health of our food and water supplies is dependent in complex ways on what is happening in the ecosystem. For example, high levels of aluminum in soil, and acid rain, which can leach aluminum from the soil, tend to increase the level of aluminum, and decrease the amount of selenium, in locally grown plants (Boegman, R.J., 1984).

Individual Differences

Recent evidence shows a complex interaction between nutrition and toxicity of ingested or inhaled pollutants. Vitamins, minerals and other nutrients can have both positive and negative effects on a person's sensitivity to the adverse effects of pollutants (Calabrese, E.J., 1980). For example, nutritional deficiencies of calcium and magnesium can enhance the absorption of aluminum, a neurotoxic metal (Boegman, R.J., 1984). People's nutritional intake and ability to absorb and utilize nutrients varies widely, depending both on lifestyle, diet, age and many other factors.

The variation in people's ability to tolerate individual foods is only recently becoming known and being utilized in diagnosing symptoms which are not responsive to other medical treatments (O'Banion, D.R., 1981). Even inflammatory cardiovascular diseases, including spastic vascular phenomena such as migraines and other vascular headaches, angina due to coronary spasm and Raynaud's disease, are responding in some cases to dietary and environmental management (Rea, W.J., 1987). For some people, dietary changes also appear to be able to remediate emotional distress (King, D.S., 1981), (Christensen, L., 1985).

The human immune system can tolerate and adapt to chemical exposures to a point, but that point is often exceeded, and the immune system can fail, resulting in symptoms and disease (Maulfair, C.G., 1987). The degree of adaptability will depend on many personal factors and individual circumstances.

Deliberate Food Additives: Not For Everyone?

Information on individual reactivity to various food additives is making it very clear that safety of additives cannot be universally guaranteed (Kon, S.H., 1978), (Miller, J.B., 1978). This makes labelling critical, and for some people, of life-and-death importance.

Healthy Environments for Canadians: PART II: ANALYSIS

British research underway indicates that perhaps as many as 15% of people feel they may be adversely affected by food additives. Actual incidence of reactions to food additives is being investigated. One way to confirm a suspected reaction to a food additive is to eliminate it from the diet for a few months (Daglish, S., 1987).

Sulfites have been widely used as preservatives in the food and pharmaceutical industries. In the United States more than 250 cases of sulfite-related adverse reactions, including anaphylactic shock, asthmatic attacks, urticaria and angioedema, nausea, abdominal pain and diarrhea, seizures and death, have been reported, including 6 deaths allegedly associated with restaurant food containing sulfites. In Canada 10 sulfite-related adverse reactions have been documented, and 1 death suspected to be sulfite-related has occurred (Yang, W.H., 1985). The American Food and Drug Administration has banned the use of sulfites as preservatives in fresh fruits and vegetables, citing growing number of adverse reactions to the compounds. The ban applies to retail sales of fresh produce by food stores and restaurants. In addition, the FDA requires new labelling of processed foods that contain detectable levels of the compounds (Raymer, W.J., 1986).

Inadvertent Additives to Food Supplies

Environmental additives are chemicals which inadvertently contaminate our food. Such substances as chlordane (a pesticide residue in milk), and sulfonamide (a drug residue in pork) can contaminate the food supply and present health hazards (Pim, L.R., 1981).

Breast milk samples from 5 different regions across Canada showed the continued presence of PCBs and a number of other chlorinated hydrocarbon compounds. There were minimal regional differences in residue levels, which were also similar to those found in other industrial nations (Mes, J., 1986).

Canadian consumers may not be able to fully protect themselves from harmful chemical substances in their food supply, even though the chemicals may be banned in this country. Central America has become a lethal dumping yard for pesticides that North American and European chemical companies are banned from selling or producing. Many foods which are imported to Canada from Central America contain residues of chemicals which are proven to cause cancer and/or central nervous system damage, such as DDT, paraquat, Phosvel, DBCP and others (Todd, D., 1987).

Not all foods are benign, or remain benign during storage. Mycotoxins such as aflatoxins, fusarium toxins, and penicillium toxins can occur naturally in some foods or agricultural commodities due to fungal infection, and have been related to cases of both human and animal disease (Scott, P.M., 1983). Some foods contain natural carcinogens or anticarcinogens (Ames, B., 1983).

Potential Health Effects of Pesticide Exposures

In an Australian study, long term exposures to organophosphorus insecticides were associated with schizophrenic and depressive reactions, along with severe memory impairment and concentration difficulty. A field survey gave some support to the possibility that psychiatric disorders might be less frequent in towns than in fruitgrowing regions of Australia where insecticides were used (Gershon, S., 1961).

Pesticide exposures at various levels have been associated with cancers, deleterious reproductive outcomes, and subtle neurologic changes. The nervous system has been recognized as a target organ for pesticide toxicity for several decades (Shearer, R.W., 1983), (Sharp, D.S., 1986). The delayed health hazards of pesticide use have been difficult to detect.

Environmental chemical exposure was postulated as a possible cause of Behcet's disease, a mucocutaneous-ocular syndrome leading to blindness, ulcers, and sometimes death. Most of the patients studied had a history of contamination with organochlorine compounds and/or organophosphorus pesticides. The number of patients in Japan with this disease declined following the banning of certain strong toxic chemicals, including DDT, BHC (benzene hexachloride) and parathion, with some time lag. The researchers postulated that Behcet's disease may be caused by environmental chemicals, which may cause supersensitivity of the tissues, possibly leading to abnormality in the immune system (Ishikawa, S., 1986).

Water Contamination

Federal researchers have been reported to have written that there is strong evidence to link Great Lakes pollution with health problems such as cancer and birth defects. Cancer rates and rates of birth defects were found to increase from west to east along the lakes, and are heaviest near highly-contaminated sections such as the Niagara and St. Lawrence Rivers. People living along the lakes suffer higher rates of heart disease and strokes than other Canadians. Followup research is planned (Israelson, D., 1987).

Many unexpected compounds can show up in drinking water or other beverages. Cadmium, for example, can enter the water supply from water pipes (Schroeder, H.A., 1973), as can copper when acidified lake water is used (Franklin, C.A., 1985). Asbestos was at one time widely used for filters in the chemical, food and beverage industries, resulting in asbestos fibres in most beers, wines and liquors, soft drinks, sugar and lard (Kruss, P., 1979). Plastic food and beverage containers can also allow various compounds to pass into food and water (Kailin, E.W., 1963).

A meeting of the Working Group was convened by the WHO Regional Office for Europe in collaboration with the Austrian Government. Legionnaires' disease, a multisystem disease with pneumonia as the principal clinical feature, acquired its name following the mysterious outbreak among people attending an American legion convention in Philadelphia in July 1976. In January 1977, a previously unrecognized bacterium was shown to be the agent

responsible for the Philadelphia outbreak. The bacterium was named *Legionella pneumophila* and is now known to be a member of a large family of *Legionellaceae*.

Dramatic common-source outbreaks of Legionnaire's disease have occurred in hotels, hospitals, and other establishments in association with contaminated water systems, in which the *Legionella* bacterium has incubated in warm standing water, for example, in shower heads (World Health Organization, 1982).

2.3.4 Physical Factors

There is an extensive literature on the effects of physical factors on people's health. These include many aspects of sound, light and other portions of the electromagnetic spectrum. It is not within the scope of the present study to review this literature completely, other than to sample it and bring it to the attention of the reader, who can follow up through bibliographies in the references that are cited. Where physical factors, or some particular aspect of them, have been recognized only recently as possible contributors to either ill health or good health, we have tried to include a more complete selection, to indicate the scope of any scientific debates still underway. Where information was available, we have included both negative effects of the physical factors, and some comment on benign or positive levels of the same factor.

Effects of Sound on Human Health

High sound levels can induce responses in the human body that are not specifically related to the auditory system. In most instances, a physiological change is evident only during, or for a short time following, the noise exposure. Noise can affect blood circulation, the resistance of the skin to electrical potentials, skeletal-muscle tension, hearing, breathing, and sleep. There is also evidence that noise can affect the gastrointestinal tract, change the size of the pupils of the eyes, and change the rate of saliva and gastric secretions (White, F.A., 1983).

Psychological and sociological response to noise is difficult to quantify. Many such reactions to noise seem to be related more to personality factors than to specific noise conditions. A person's psychological response will also be conditioned by whether he or she feels that the noise is a personal infringement of a basic right to acoustic privacy. Other influencing factors are whether the noise is essential, the relationship of the noise to personal activity, the noise's predictability or unpredictability, and frequency of occurrence (White, F.A., 1983).

In a U.S. study, a positive, but not statistically significant, association was found between aircraft noise exposure and the risk of high-frequency hearing loss (Green, K.B., 1982). A comparison made by others, between a sound-proofed and a non sound-proofed school, both exposed to aircraft noise, noted that children did not express any feeling of disturbances as to the frequency of airplane passage. However, after prolonged exposure to the auditory stimulus, perturbations of certain intellectual, psychomotor and personality aspects were observed. Noise-triggered attentional troubles were suggested as an explanation for most of the negative effects observed (Moch-Sibony, A., 1981). Another study indicated that the percent of elementary school students reading below grade level increased as the noise exposure level from aircraft increased (Green, K.B., 1982).

Effects of Exposure to Different Kinds of Light

Possible connections are being investigated between skin cancer incidence and occupational ultraviolet light exposure from fluorescent lights as well as from lamps used in printing and dyeline copying (Elwood, J.M., 1986).

Recent research indicates that there are a great many non-visual effects of light on people. Of these effects, the suntan effect and the control of rickets are two that are well understood. Other important but less well known effects include synchronization of a number of physiological rhythms, and prevention or control of infantile jaundice (Hathaway, W.E., 1983). Human muscle strength appears to be weakened under exposure to light that lacks part of the natural spectrum (McGee, C.T., 1979). A study comparing classroom behaviour in windowless rooms with standard type fluorescent cool-white tubes, with that in rooms with full spectrum, radiation-shielded fluorescent fixtures, showed increased hyperactive behaviour (Ott, J.R., 1976).

Windowless environments, especially in deep-core buildings, have a number of disadvantages that are reviewed in the literature (Health and Welfare, 1980). Teachers like the absence of windows in teaching spaces because they prevent the students from being distracted by outside happenings and extra wall space can be put to good instructional use. Others contend that the curious child in a well-conducted classroom already has an information overload. Some professional educators, however, have questioned whether the elimination of outside distractions is always something to be desired — an exterior happening may frequently provide a fruitful stimulus to educational activity (University of Michigan, 1965).

Effects of Exposure to Low-Frequency Electromagnetic Fields

A growing body of research indicates that electric and magnetic field exposures can have biological effects and may cause stress to human beings (Becker, R.O., 1982). However, there is not yet a clear consensus among researchers as to whether these effects constitute a long-term risk to human health, and, if so, the magnitude of that risk (Budd, R.A., 1985), (Ontario Ministry of Health, 1987).

Earlier studies of electrical power transmission and distribution workers, as well as railway high-voltage substation workers, did not yield any significant correlations of health with the subjects' measured or estimated exposure to electromagnetic fields (Broadbent, D.E., 1985), (Baroncelli, P., 1986). Later evidence has raised the possibility that an increased incidence of certain cancers, including leukemia, might be related to exposures to power frequency fields, including fields generated within homes (Edwards, D.D., 1987), (McDowall, M.E., 1986), (Savitz, D.A., 1987). This work has been reviewed by others and is not sufficiently conclusive or convincing to warrant the conclusion, at this time, that extra low frequency electromagnetic fields pose a significant health risk (Ontario Ministry of Health, 1987).

There is some evidence that the nervous system may be affected by ELF electric fields that are far too weak to have a direct effect on brain synaptic function or membrane excitability through field influence on ionic motion through the cell membrane. Animal studies have also shown effects on calcium exchange from various brain tissues. However, none of the evidence indicates that even strong electric fields have effects that compromise a person's ability to function, or have long-lasting or permanent effects on neuro-physiological health (Jammot, H., 1985).

There are preliminary reports from the United States (Rea, W.J., 1987) and from Great Britain (Smith, C.W., 1987) that a small segment of the population may exhibit a sensitivity to electric and magnetic fields, and may in fact be particularly sensitive to certain specific frequencies of field oscillation, including those in the power frequency range of 50-60 Hz. These individuals tend to exhibit immediate or chronic physical and/or psychological symptoms when in the presence of various electromagnetic fields. Massive pesticide exposures were suggested as the possible cause for some individuals being rendered both chemically sensitive and electromagnetically sensitive (Rea, W.J., 1987).

The possible effects of home and office electromagnetic exposures is now becoming of some concern to individuals seeking healthier lifestyles. Typical magnetic field exposures from hair dryers, motors, and other home appliances far exceed the magnetic field exposure even directly beneath high-voltage transmission lines (Epstein, S.S., 1974), (Becker, R.O., 1982). Video display terminals can expose operators to both electric and magnetic fields, and new designs have been suggested to incorporate more shielding (Marha, K., 1982), (Marha, K., 1983).

Although there are no Canadian standards for electromagnetic field exposure, Ontario Hydro notes that strict Soviet standards would still permit unlimited exposure by the public, of the fields under Ontario's 500 kilovolt lines (Ontario Hydro, 1983).

Exposure to Ionizing Radiation

There are a number of occupational and health care environments in which people in Canada may be exposed to ionizing radiation, which is known to have adverse health effects at certain exposure levels. Some radioactive materials emit high-frequency radiation which can penetrate the body even though the materials remain outside (gamma rays and X-rays). Some radiation is emitted in the form of beta particles, which are able to penetrate the outer skin layer and do some internal damage to humans. When radioactive particles are taken within the body through inhalation or ingestion, they can do more severe local biological damage to the cells immediately surrounding the bone, organ or tissue in which they lodge (Bertell, R., 1984).

Other Physical Factors

The percent utilization of total blood sulfhydryl group shows changes under stress from heat, cold, hypoxia, head-tilt, acceleration and vibration. Changes varied from a fall of 61% from normal to a rise of 43% (Iyer, E.M., 1985).

Studies have been undertaken of the effect on respiratory diseases of increasing indoor relative humidity. Kindergarten children, school-children, soldiers and office workers were studied, and the results showed significantly lower absenteeism and/or occurrences of respiratory illnesses in the spaces with the higher humidities (Green, G.H., 1979).

Breathing an increased concentration of ambient negative air ions was shown in one study to improve incidental visual memory in children. The action of negative ions on the neurotransmitter, serotonin, is suggested as the possible mechanism by which negative ions could produce such effects (Kershner, J.R., 1984).

2.3.5 Social Conditions

It is becoming increasingly clear that many social conditions are stressful to people, and may have an adverse impact on health (Selye, H., 1976). The literature has many suggestions as to what may be detrimental, but not much solid proof to link the suggested causes with the effects claimed. Some medical research, however, is demonstrating very specific stress effects within the body, in particular, on immune function, which lends some credence to the more general claims.

To the extent that researchers have suggested positive social conditions that will promote health, we have included discussion of them. Participation of individuals in decisions that affect them stands out as one of the most concrete suggestions. Personal situations such as poverty and unemployment, which may arise out of more general social conditions, are touched on in this section, but will be dealt with more specifically in Section 2.3.6 "Employment and Money".

The Specific Health Effects of Social Stresses

Psychological and social stressors have been found to affect the immune response and disease susceptibility. Different individuals can respond quite differently to the same environmental stressor, and patterns of immediate and delayed responses to stressors may differ markedly (Ciaranello, R., 1983). Interactions between different classes of stressors can exert markedly more severe adverse effects than can a single stressor in itself (Bell, I.R., 1982). There are stressful levels of environmental stimulation which can represent an overload. Conversely, social isolation and restricted environmental stimulation can lead to stimulus underload, which can also be unhealthy (Suedfeld, P., 1983). It is possible that high-energy places, people and activities can strengthen one's immune system (Humiston, K., 1987). Although environmental factors may contribute to stress, the specific behaviour which results will relate to each individual's own way of perceiving his or her environment (Keating, J.P., 1983).

Increased psychological stress (as typified by examination stress in graduate students) will alter immune functioning and heighten psychological responses (Didriksen, N., 1986). Humoral immune functioning, as measured by immunoglobulin levels, shows an increase during periods of stress. Anxiety can be related to external events, and this can increase under stress. Anxiety is associated with lower emotional stability and coping skills, and is coincident with tension, an increased number of physical complaints, and obsessive-compulsive trends (Didriksen, N.A., 1987).

Of course, not all environmental factors are negative. Seeking positive influences in life, including good family and spiritual life, food, clothing, housing, reading, music, art, and socializing, can have healing effects (Humiston, K., 1987).

Social Conditions Which Adversely Affect Health

Both physical factors and social factors in our environment contribute to our pace of life. Physically threatening, noisy, or polluted environments can be stressful. Similarly, information overload or decision overload can be part of a stressful environment. Environments which are unsuited to the user also produce stress (crowded freeways, prohibition of play in buildings containing children). Some environments carry psychologically and socially meaningful messages that affect people within them (e.g. people experiencing increased status become healthier, people whose educational background exceeds their current status are less healthy than others in the same circumstances) (Saegert, S., 1975). Uprooting, dehumanization of people by institutions, side-effects of innovations, and politics which constrains environmental health programs, are also seen as stressful (Levi, L., 1983).

Environmental stressors interact with numerous psychological variables to produce adverse effects on health and well-being (Campbell, J., 1983). Psychological factors such as perceived control, information about the stressor, coping resources, and certain personality traits all may mediate the influence of physical environmental stressors on human well-being. Global, chronic phenomena such as air pollution, community noise, and crowding can be viewed as a distinct class of stressors, namely ambient stressors. Such stressors represent noxious stimulation and place demands upon us to cope. Ambient stressors over the long run are expected to affect several dimensions of human well-being, including motivation, emotions, attention, somatic health, and behaviour (Campbell, J., 1983).

Stressors in the family and job environments have been proposed to play a role in the modulation of pain. One study indicated that environmental stressors, including family conflict, family control, and general stress, were greater in the group with chronic low back pain than in the group without. The relationship turns out to be a complicated one which illustrates how difficult it is to make competent generalizations about the effects of stress. For example, increased family conflict was associated with increased psychological distress and increased incidence of pain, but increased family independence was correlated with less distress and increased pain. Less peer cohesion, less physical comfort, and less job clarity were correlated with increased pain, but not distress. Work pressure was associated with decreased depression and less pain, possibly decreasing pain by distraction (Feuerstein, M., 1985).

In the design and organization of many physical settings, the human properties of the individual are ignored or oversimplified. Spaces and places are often improperly designed in physical terms. Human needs for privacy, territoriality and freedom of choice are often overlooked as well. The danger is that we will adjust at the price of a continuing erosion of the properties that make us distinctively human (Proshansky, H.M., 1975).

Different socio-economic groups deal with assaults on the territory adjoining their homes in different ways. For instance, residents on a high-crime, low-income street have little material or political means to use as a leverage against possible threats. Co-resident distrust is likely,

thereby precluding effective group social action against a threat. When people experience less control over this territory than is desired, the discrepancy is viewed as stressful (Taylor, R.B., 1985).

The pursuit of environmental security in an urban context becomes an unending process for urban dwellers. Physical settings increasingly make demands on people because of their frequently growing complexity and accelerated change. Having control, understanding, and competence in any setting are matters of critical importance to a person's self-identity and related feelings of self-esteem (Sarason, I.G., 1983). Fear of crime is greatest in communities that either lack the power to regulate themselves or perceive that they lack such power (Nietzel, M.T., 1986).

In cities, the nature and intensity of environmental stimulation, including its predictability and the extent to which it can be controlled, assumes considerable importance (Kaminoff, R.D., 1981). The quality of urban life is rooted in factors such as freedom from intergroup conflicts, full employment, maintaining democratic freedoms, and integrity in public life (Proshansky, H.M., 1986).

In local environments such as homes for the elderly, different modes of operation generate different stresses. In one case, problems arose when alert patients were housed together with confused patients, and antagonized each other. When confused patients were housed with frail patients, the former found their environment sad and depressing. This environment tended also to injure the confused resident's self-esteem, reduced functional competence, and increased the potential for aggression. As a result, the home was reorganized around four primary patient needs: independence, emotional support, safety and security, and physical care (O'Connor, D., 1987).

Poverty, lack of good housing, violence in the media, child abuse, parents who engage in antisocial behavior and alcohol abuse are cited as risk factors for children developing behavioural and psychological disturbances (Gelfand, D.M., 1986). Children are also aware of the arms race much earlier than adults suspect. This awareness and the perception that few adults are doing anything to prevent a nuclear war result in a profound distrust of adults and adult values (Goldberg, S., 1985).

Widespread abuse of illicit drugs and alcohol, the use of physical violence in domestic crises, and extensive use of cigarettes and caffeinated beverages such as coffee are all signs of a stressed population. Each of these practices in turn has health and injury consequences and poses a danger to public health (Meggs, W., 1987).

The uncertainties offered by both military and civilian uses of new, complex technologies may represent considerable stress, though this has not been well researched. Actual episodes of environmental contamination also represent sources of stress, through fear of passive exposure to toxic sources (Smith, L.F., 1987). Nuclear technology is one example; many people are concerned both by the potential for nuclear war, and by the use of nuclear power to produce electricity. Some are influenced by the psychological

association between nuclear power and nuclear weapons, while others are concerned about the gaseous and liquid radioactive wastes discharged during normal operation, the potential dangers from the radioactive clouds that could escape in the event of a nuclear accident, the difficulty of disposing of radioactive waste, the toxicity of plutonium, or the possibility that plutonium might be diverted clandestinely into nuclear weapon production (Eisenbud, M., 1978).

Some researchers feel that dense urban neighborhoods may be related to low neighboring among residents. Crowded urbanites may withdraw from local contacts with neighbors as a specific adaptation to immediate high levels of stimulation (Baldassare, M., 1975), (Saegert, S. undated). Others feel that density, though perceived as unpleasant, does not appear to have definite and consistent detrimental effects (Fischer, C.S., 1975).

Participating in Design of Our Environments

Most of us believe we have no opportunity to design or shape our environment. Professionals often make design decisions in isolation, although the environment has an awesome impact on our lives (Abel, C.B., 1981). Involvement in planning one's own environment is vital to self-esteem and good mental health. The probability of achieving an environment that will prevent mental illness is increased when planning of the environment is conducted by those who must function within the environment (Insel, P.M., 1983b). People have an active need for control over their own environment. Having control reduces the stress generated by an aversive event by providing individuals with an guaranteed upper limit, or by enabling them to match their internal state with external events (Miller, S.M., 1980).

Citizen participation in environmental decisions is critically important, because this is the way in which people can become identified with a new environmental action — the way in which they can possess and feel responsible for it — thus reducing their alienation. Identity, recognition, and some sense of power are human needs. Because an environment becomes a social symbol when it is perceived as representative of someone or some social group, physical planning decisions can threaten the identity and status of certain groups, while enlarging the powers of others (Appleyard, D., 1979).

Health promotion and disease prevention involving participation by many different sectors, as well as the public, is seen as necessary by one researcher. Community development is offered as the proper approach to many health problems, including: appropriate technology in waste management, transportation, food production, energy use, and manufacturing process; quality food, shelter, work, safety and education with the emphasis on self-reliance rather than dependence; integrated caring neighbourhoods, built to human scale; support services for those with special needs through a strong social support network; a wellness system rather than an illness system; and involvement of the whole community (Layton, J., 1987).

Many of our common environments could be more accessible and more adaptive than they are. An adaptive environment is possible when a) the users are skilled enough to use basic design tools to make small changes as their individual and group needs change; b) the environment supports specific activities that occur in the space for all users regardless of age; size; mental, physical or emotional state; and c) users and designers together assess what works and what fails to meet users' requirements, with successes and failures documented for future reference in designing similar settings (Abel, C.B., 1981).

The Need for Change in Social Conditions

A number of researchers expressed frustration that in overall health strategies in Canada, individual self-care is stressed without recognition of the overall physical and structural problems in our society. This emphasis may deflect attention away from environment and socioeconomic causes of ill health. It also pushes people to assume impossible levels of individual responsibility without recognizing that they have decreasing control over the environment in which they live (McKillop Farlow, D., 1987). Little attention has been paid to deliberately designing stress-reducing environments (Saegert, S., 1975).

Honesty in communication is proposed by one researcher as an essential stress-reducing ingredient, using the example of potential environmental contamination, such as would be involved in PCB spills or nuclear accidents. A steady flow of information must be maintained, even if it means admitting the temporary absence of data (Smith, L.F., 1987).

Another researcher proposes a new 'paradigm', or set of assumptions, to guide public health policy: that an individual becomes ill when he or she cannot resist or avoid biological organisms and their by-products, pollutants, or safety hazards; that resistance is lowered by stress; that the type of health problems attributable to pollutants will depend on the community's economic base and natural setting; and that health problems attributable to safety hazards are predictable from a community's economic base. The paradigm would expect that as stress levels rise in a given community, those with genetic predispositions will become ill before those with average or above average tolerance. If public health is to contend with the illnesses precipitated by stress, pollutants, and safety hazards, it must become more familiar with the economic and social processes which shape the human community (Catalano, 1987).

Yet another researcher suggests various ways of changing work to make it less stressful: increasing a worker's control of work arrangements; providing mechanisms for worker participation in decision making on the organization of work; avoiding monotonous, machine-paced, and short but frequent work actions; optimizing automation; helping workers see their specific task in relation to the total product; avoiding qualitative work overload or underload; facilitating communication and support systems among work mates and others (Levi, L., 1983).

2.3.6 Employment and Money

The Price of Unemployment

What has never really been faced and adequately dealt with, about unemployment, is the physical and emotional toll it takes from those affected. This may be due as much to attitudes toward employment in Canada, as it is to the financial difficulties individuals and families face when income drops significantly. Full employment in the traditional sense may be an economic fantasy; but finding more creative ways of helping everyone feel valuable, cope financially, and keep his or her life goals on track, fully employed or not, may not be. We are going through an era of post-industrial development which may involve considerable dislocation and adjustment. Better attention to the human consequences could bring Canadians the message that their health and wellbeing is at the top of the national agenda, and not just secondary to economic goals.

Substantial health differences have been observed between employed and unemployed individuals. The unemployed show significantly higher levels of distress, show greater short-term and long-term disability, report a larger number of health problems, have been patients more often, and use proportionately more health services. Unemployed individuals appear to be more vulnerable to serious physical ailments such as heart trouble, pain in heart and chest, high blood pressure, spells of faint-dizziness, bone-joint problems and hypertension. While these health differences persist across socio-economic and demographic conditions, females and older unemployed individuals report more health problems and physician visits whereas people under forty report more psychological distress. The blue-collar unemployed are considerably more vulnerable to physical illness, whereas the unemployed with professional background report more psychological distress. The low-income unemployed who are also the principal family earners are the most psychologically distressed (D'Arcy, C., 1985).

Evidence linking unemployment to poor mental health does exist and is stronger than that linking it to poor physical health (Hertzman, C., 1986). Mental hospital admission rates vary according to economic changes (Kirsh, S., 1983). People who are unable to work undergo not only economic hardship, but also experience physical, psychological, and spiritual distress. Preventive mental health care will require assessment, anticipation, and response to, the social impacts of employment patterns (Canadian Mental Health Association, 1984).

When personal and financial resources are stretched to the breaking point, women as a group are affected particularly severely, as they continue to try to provide mothering, support and household comforts in an atmosphere of tension and strain. Feelings of guilt, anger and despair, turned inward or outward, are a powerful part of the chronic unemployment picture (Canadian Advisory Council on the Status of Women, 1987).

For the mentally disabled, finding employment presents a special set of problems including the stigma of mental illness, the effects of some medications, time gaps in resumes, lack of work incentives, and an entrenched sheltered-workshop mentality (Kirsh, S., 1983).

The Price of Poverty

A special study of Ontario's health care system concluded that children in families living on welfare get sick because of poor nutrition and hostile living environments (Spasoff, R., 1987). Children from families living on welfare are three times more likely than their schoolmates to end up repeating a grade or being sent to remedial classes for slow learners. Only 46 per cent of the children of the poor end up in advanced high school programs needed for university entrance, compared with 88 per cent of children of middle-income-earning families (Crawford, T., 1987).

Health researchers from the National Centers for Disease Control in the United States determined that many children under five years of age from low-income families are shorter or fatter than most other youngsters, possibly because of chronic under-nutrition (Toronto Star, 1987). Other sources also confirm that children living in poverty suffer adverse health consequences (Miller, C.A., 1985). One million Canadian children live in families with incomes below the poverty line (Crawford, T., 1987).

The prevalence of risk factors for cardiovascular disease tends to be higher among men and women with a low level of education, and hence among people with generally lower income. (The risk factors considered include cigarette smoking, obesity, elevated diastolic blood pressure, physical inactivity, excessive alcohol consumption, elevated serum cholesterol level, diabetes mellitus and the joint use of oral contraceptives and cigarettes.) Both men and women in lower socioeconomic groups are more likely than higher-income people to die from cardiovascular disease (Millar, W.J., 1986).

Research in the United States supports the hypothesis that properties of the sociophysical environment in federally-designated poverty areas may be important contributors to the association between low socioeconomic status and excess mortality, and that this contribution is independent of individual behaviours. Residents of a federally designated poverty area experienced higher age-, race-, and sex-adjusted mortality over the follow-up period compared with residents of nonpoverty areas. This increased risk of death persisted when there was multivariate adjustment for baseline health status, race, income, employment status, access to medical care, health insurance coverage, smoking, alcohol consumption, physical activity, body mass index, sleep patterns, social isolation, marital status, depression, and personal uncertainty (Haan, M., 1987).

Resolving the Dilemma

New 'work agendas' may be required, in which some of the ideas people in Canada presently hold about work can be updated to take into account economic realities (Novick, M., 1986), (O'Hara, B., 1987). Access to decent jobs remains a primary life aspiration for the vast majority of the adult population in Canada, yet full employment in the traditional sense may be virtually impossible to achieve, particularly with the present pace of dislocation and change arising from both local and world adjustments in trade and technology. The present distributions of work among Canadians are also very rigid, and may require change, for example, to even out the extremes of the overworked and unemployed (O'Hara, B., 1987).

Without some form of resolution which allows both sufficient income, and sufficient dignity and respect, for those who are employed less than full time, or at less challenging work than they would like, we can expect adverse health impacts and growing stress for many people. Some suggestions include specific changes such as special tax rates to cushion annual job hour reductions, shelter policies to stabilize housing costs, national income credits for parents, universal leave programs, enriched pension entitlements, and tuition rebates and interest free loans for students of all ages (Novick, M., 1986).

Social, economic and collective determinants of lifestyles have tended to be ignored in recent approaches to health promotion. A focus on changing personal habits and lifestyles has been more effective among the higher educated and higher income groups. There is a danger that isolating lifestyle and personal responsibility as major determinants of health, without attention to the structural changes needed, will lead to frustration on the part of those who see a broader change as necessary (Angus, D.E., 1986).

2.3.7 Personal and Social Support

Stressful life events are associated with the onset, incidence, and prevalence of a wide range of physical and psychiatric disorders. Social support networks have been studied as possible buffers or mediators of such stress (Dean, A., 1977), (Fischer, C.S., 1982). Considerable work is still required, however, to fully understand both the role of life events and the role of support. The connection between life events and health is influenced by many factors, including a person's social characteristics, his or her stage of life, personality traits and coping repertoires, as well as the supportiveness of social networks (Dohrenwend, B., 1983).

Is having friends enough? Not always, because the concept itself means different things to different people. One study found that this label is likely to be applied: to an overwhelming majority of non-relatives in a largely unsystematic way; to associates lacking other, specialized role-relations; to people of the same age; to people known a long time; and to people with whom respondents had primarily sociable, rather than intimate or material, involvements (Fischer, C.S., 1982).

People may vary in their need for, and ability to sustain, social relationships. Some evidence indicates that elderly people may have fewer social relationships than younger adults, but are more content with what they do have. Elderly women who were studied had more affectional ties than elderly men. The presence of offspring in the same town increased the number of close ties and of social relationships, but were more important for men than for women. Elderly people with cognitive impairment or dementia had less social interaction than they would like. Elderly people who were depressed reported having markedly less social interaction than the mentally healthy elderly, but did not complain that it was too little (Henderson, A.S., 1986).

Stress associated with organizational settings, such as work or school, constitutes a major part of the total stress experienced in people's lives. Organizational settings place demands on people to perform and to relate to a broad range of others in specified ways. At the same time, organizations present opportunities for much human contact and could be valuable as part of an effort to reduce stress or improve health (Kahn, R., 1983).

The physical and social environment does influence whether we are well-connected with others, or largely alienated. Many researchers have claimed that modern urban life can destroy community, and some evidence tends to support this. For example, people with below-average household incomes and living in semirural communities were found to have more local non-kin associates, bound up in denser networks, than did low-income respondents residing in other places. The semirural people tended also to report a slightly more positive sense of well-being than their counterparts elsewhere (Fischer, C., 1982).

It is clear that our social networks do influence our behaviour, and in turn, that behaviour can affect our health. For example, drinking and smoking by both male and female study subjects have been directly associated with that of male and female friends, with congruent gender relationships being strongest. Activity level for both genders was positively associated with exercise by male friends. Prevention programs may need to incorporate strategies to maximize peer support for healthful behavior, and to counteract the effects of unhealthful behavior modeled by peers (Gottlieb, N.H., 1986).

Our perspective on how to achieve individual health may have to widen to include community environment, if we are to be successful. One researcher emphasizes that the process of healing "involves not only the individual, but the community, tribe, and family as part of the larger social and physical environment" (Duhl, L.J., 1984). This may be another way of saying that because we are influenced by our larger environment, we cannot achieve individual health without taking action to improve the larger environment.

2.3.8 Health Care and Social Services

Incorporating Environmental Awareness into Health Care

A competent, efficient and accessible health care system is a necessary component of a healthy environment. Its approach to health promotion and to healing may set the upper and lower limits to health in Canada. The message from the literature, over and over, is that the approach to health care in North America does not adequately recognize the role of environmental factors in health. This is part of a larger issue of general adaptation to many changing needs, including those of an aging population, a post-industrial society, and changing human service needs (Vayda, E., 1986).

The revolution in understanding of the health effects of polluted indoor environments arose in non-medical professions. Architects, engineers, building managers, chemists, ventilating experts, educators, learning specialists, and many others, have gained and spread information about low-level chemical exposures and their effects on health. One of the difficulties many of them face is a shortage of knowledgeable medical support for the environmental health problems they are now dealing with daily. Diagnosis and treatment of environmentally-related health problems are still concentrated in the hands of a small number of specialists in Canada, and have not been incorporated well into general practice. Multi-disciplinary programs are beginning to be developed to respond to indoor air quality problems (Marchant, R., 1985), (Gosnell, D., 1987).

As the health care system begins to assimilate new information about the effects of environment, we will see it applied more often by the general physician in diagnosing the role of physical factors in disease. A detailed knowledge of the effects of primary outdoor and indoor air pollutants, on both healthy and impaired people will be required (Guidotti, T.L., 1983). Scientists working on health complaints associated with energy-efficient, tightly-constructed, mechanically-ventilated office and residential buildings predict that physicians are likely to see increasing numbers of patients who work and live in 'sick' buildings (Riesenberg, D.E., 1986).

Considerable clinical experience has accumulated, primarily in other countries, with the effects of environmental contaminants on health. A spectrum of disorders affecting smooth muscles, mucous membranes, and collagen in the respiratory, gastrointestinal, genitourinary, and vascular systems have been described. The variety of symptoms presented are often mistaken for hypochondriasis, but actually can be triggered by foods and chemicals found in the patient's home and work environments (Rea, W.J., 1979a).

Ear, nose and throat involvement with environmental triggering factors appears to be the main early warning sign of environmentally triggered disease. Sensitivity to odours seems also to be common, many persons being intolerant of the odours of such substances as car exhaust, perfume, cigarette smoke, aerosol sprays, formaldehyde, alcohol, phenol, food, mould and dusts. Often rhinitis occurs. Recurrent sinusitis as well as severe otitis, various

forms of vertigo (including Meniere's disease), and laryngeal edema are frequent presentations of the problem. Any portion of the respiratory system can be involved, resulting in such inflammatory diseases as recurrent bronchitis, bronchiectasis, and asthma (Rea, W.J., 1979b).

Since some human exposure to environmental contaminants appears inevitable, despite the best application of environmental laws and protection technologies, the health care system must also consider the need to develop adequate treatment methods (Selner, J.C., 1986) as well as the potential of reducing levels of contamination within the human body once they have accumulated (Schnare, D.W., 1984).

Hospitals as Healthy Environments

Just as the whole health care system requires greater focus on the effects of environment on health, a greater understanding of environmental effects within hospitals and other health care facilities is also important. For example, a well-designed physical environment can play a role in the prevention and reduction of psychological and social problems encountered by patients in acute care and psychiatric institutions. Factors that are important include spatial layout and design of hospital environments, privacy problems, personal control and independence, information interventions, hospital social relationships, and levels of environmental stimulation (Winkel, G.H., 1985).

For children, hospitalization is often a frightening and stressful experience. The child may feel insignificant, anonymous, overpowered, isolated, and intimidated by the scale and procedures of the hospital. The provision of familiar environmental cues through attempts to make certain areas of hospitals more home-like can help. For adolescents, greater opportunity to exercise choice and control in the physical environment will most likely result in less time spent resisting authority. Design also needs to take into consideration the need for family participation in the child's hospital stay (Ferguson, R., undated).

Greater Responsiveness to Hazards in the Workplace

While occupational disease is not a newly uncovered problem, its potential magnitude has only recently been widely recognized. The increasing awareness has led to pressure for regulating industrial health hazards and to efforts for cleaning up the workplace to minimize adverse effects of physical agents, biological hazards, chemical agents, ergonomic factors, and psychosocial stress. Many work-related diseases are not likely to be diagnosed because some physicians lack awareness of occupational hazards and neglect to take a comprehensive occupational history. Claims for many potentially work-related diseases are almost systematically rejected in the absence of strong supportive research (Yassi, A., 1982).

Measuring Behaviour and Performance as a Function of Environment

The role of hypersusceptibility to both chemical exposures and food in behaviour problems has also received greater attention in non-medical settings, such as school boards, than it has within the health care system (Small, B.M., 1985). Poor nutrition, toxic metal and other chemical exposures, and food allergies are among the adverse factors now considered as possible causes of brain malfunction in young people (O'Banion, D.R., 1981), (Rimland, B., 1981), (Davis, K., 1985). Maladaptive central nervous system reactions to chemical exposures or to otherwise nutritious food can occur in susceptible individuals (Philpott, W.H., 1981). Introduction of new knowledge in this area into clinical situations will require a broader repertoire of assessment and treatment techniques than is now commonly used (Strickland, B.R., 1982).

At the same time, behavioural toxicology is now firmly established as a legitimate component of the environmental health sciences. The adverse health impact of environmental chemicals can be gauged not only by death or overt damage, but by how people feel and function (Weiss, B., 1983). For example, exposure to carbon monoxide has caused performance deficits in experimental subjects (Laties, V.G., 1969). Behaviour and performance measures are increasingly being investigated as a first sign of impairment from low-level chemical exposures (Valciukas, J.A., 1980).

Greater Emphasis on Prevention Through Environmental and Dietary Control

Some criticism has been levelled at Canada's health care systems because they are often treatment-oriented rather than prevention-oriented. For example, some researchers maintain that lifestyle and environmental factors cause or promote at least 70% of cancer cases, and that many of these factors are controllable at the personal level (tobacco smoking, diet, occupational choices, alcohol consumption, viruses, excess sunshine, medicine and medical procedures, food additives, and environmental pollution). Motivating the public and the medical community to make changes in their lifestyles and professional practices is suggested as the next advance required to prevent cancer (Cullen, J., 1986).

The clinical spectrum of conditions in which environmental factors are of importance is being expanded. Studies are appearing which confirm that environmental control is of value in the management of many diseases (Finn, R., 1986). Similarly, there are some diseases that new research is finding responsive to dietary controls. For example, in a blind, placebo-controlled study of dietary manipulation therapy in outpatients with rheumatoid arthritis, there was significant objective improvement during periods of dietary therapy compared with periods of placebo treatment (Darlington, L.G., 1986).

New Research Into Environmental Hazards

As environmental health research expands, information on more new developments will require rapid dissemination to health professionals and assimilation into everyday investigations within the health care systems in Canada.

For example, the role of mould growth in Canadian residences is getting new attention. Potentially toxigenic fungi can be isolated from house dust. These produce mycotoxins, either volatile or absorbed onto particulates, which can have subtle effects on the immune system. These effects are often masked by the more obvious clinical and pathological findings of opportunistic infections, thus allowing the underlying cause to escape the attention of clinicians (Schiefer, H.B., 1985). Some researchers are suggesting that identification of all moulds in a household is in order, when the home contains individuals who complain of multiple symptoms for which no medical explanation can be found by other means (Day, J., 1986).

Very little information is available in the medical literature about the health effects of moulds and yeasts, compared to the clinical knowledge available. Sensitivities to moulds can cause a variety of symptoms such as headaches, ear and hearing problems, respiratory problems like bronchitis and asthma, gastrointestinal difficulties, weight gain and loss, food cravings, as well as dermatologic, muscular, urologic, and neuropsychiatric symptoms (Calabrese, D.V., 1986).

New mechanisms are needed to further clinical research on the identification and treatment of ill health that is triggered by, or aggravated by, environmental conditions. The rapid dissemination of knowledge about new diagnostic and therapeutic techniques is critically important in an era when environments change rapidly. For example, within a decade, the combination of new building materials and an emphasis on energy conservation led to a need for understanding the effects of exposures to complex mixtures of volatile organic contaminants, with each component at a relatively low concentration, over extended periods of time (for example, in residential exposures).

Long-Term Visions of Health Care in Canada

Many of the long-term visions for health care in Canada include an increased emphasis on health promotion and disease prevention. Strengthening the role of the individual and achieving a community focus for health have been suggested as two actions which will contribute to a system which leads to greater health promotion (Evans, J.R., 1987).

One researcher suggests a synthesis of environmental health, community-based public health, the primary health care concept, the holistic health movement and the self-care movement. It is compatible with and leads towards a sane, humane and ecological future (Hancock, T., 1980).

Greater focus on environmental health has been suggested, including: improving and organizing the data base on environmental health, especially in terms of linkages between morbidity/mortality data and environmental exposure; developing national environmental health policies aimed at prevention; developing more comprehensive and standardized risk assessment processes which take into account economic, social, and other factors; conducting research in the biological sciences with emphasis on individual sensitivities and the identification of defence mechanisms against toxic substances that contribute to ill health; and strengthening environmental health in medical training. Public agencies should be encouraged to inform the public about comparative risks and the risks of individual hazards (World Health Organization, 1983).

Health for all cannot be achieved by the health sector alone; it requires contributions from, and a new kind of co-operation with, many other sectors, including agriculture, animal husbandry, food, industry, education, housing, public works, and communications. At the same time, the health sector also needs to participate more actively in support of other sectors, for example, by participating more in surveillance and in the development of safeguards in the area of industrially-generated health hazards (World Health Organization, 1986). Massive changes will have to occur that involve public policies in many areas. Regional and social class inequities, as well as health problems due to substance abuse, industrial hazards, environmental pollution, unemployment and social isolation, must be addressed. The health care systems must allow and encourage physicians to practise health promotion (McWhinney, I.R., 1987).

Physicians are not the only care-givers that have a major influence on, or participate in, social issues which contribute to health. Nurses, for example, have always been a strong social force, and continue to be involved in economic, energy, environmental and social welfare problems. Other major issues affecting health are population growth, poverty, education, clean water supply, and family planning. Individual nurses and nurses' associations can exercise their influence and power in promoting health at local, national, and international levels (Labelle, H., 1986). The fate of humane, family-centered health care may rest on the ability of different caregivers to respond well to complex needs of families in crisis. The health social worker can play a major role in negotiating between families and health care professionals during illness (Dillon, C., 1985).

In a new vision of our health care systems, medical education would be up-to-date and would realistically reflect the actual health needs of the Canadian people. We all will have advanced beyond the concept that health is merely the absence of disease and that the physician's goal is to recognize and treat disease. Preventive medicine and health promotion will play an important part in the curricula. Clinical programs will allow more time in settings where students can acquire a more realistic view of the nature and extent of the major health problems that afflict people in Canada. In this vision, physicians carry some community responsibility for spreading information about healthy lifestyles (Squires, B.P., 1987).

Whatever vision takes hold, it is clear that the health care systems of the future must welcome the active participation of many more people, so that the full diversity of needs in Canada is taken into account, and so that the system stays more in touch with the realities of the day.

3. Bringing About Healthier Environments

This final analysis section focuses specifically on broad principles and ideas that emerge from the literature review, upon which strategies for change can be based. Five important components have surfaced during the review:

- ending prejudice and oppression
- creating a new vision of environmental choice
- fostering technology for assessing our environment
- accommodating diversity through participation in design
- becoming world experts in designing healthy environments

Each section following will explain briefly how these ideas are connected with achieving healthy environments for everyone in Canada.

3.1 Ending Prejudice and Oppression in Canada

Throughout the discussions in previous sections, we have pointed out the coincidence of unhealthy environments with conditions of prejudice. It is becoming clear that people who are held in high value in our society do not appear to get stuck in environments recognized as unhealthy, as often as those whom we value less.

While major strides have been made in human rights legislation, personal attitudes, and day-to-day behaviour in Canada, both overt and subtle forms of discrimination still persist. We are now understanding and acknowledging that even seemingly minor forms of devaluation of one group of people by another can lead to unhealthy conditions for the former. The effects include decreased self-esteem, decreased physical and mental health, higher risk of health impairment, and decreased resources and independence, which in turn limit an individual's control over health and environment.

The irony is that our tendency to rank and devalue others (which may have, at one time, had some rationale) is self-defeating in the mixed society of Canada today. In fact, about one in four Canadians feel they have been discriminated against in one way or another (Toronto Star, 1987). It is unlikely that anyone in Canada can go through life today without becoming, at least temporarily, part of a group whose characteristics have been devalued in some way. For example, everyone starts out young, and ends up old; most go through periods of economic upheaval and many experience times of unemployment and poverty; and many people go through periods of physical disability and illness.

The kind of attitude that ensures that everyone can create a healthy environment goes beyond mere tolerance of differences between people. Simply transforming an inferiorized group from outcasts to a minority, making acceptance conditional upon assimilating and hiding differences, or providing token power without full equality, will not accomplish healthy environments for everyone (Adam, B.D., 1978).

What we must aim for is the idea that everyone in Canada is a person of value, no matter what age, gender, race, religion, ethnic origin, culture, physical shape, mental ability, sexual orientation, employment, financial level, lifestyle, or any other characteristic that has been used to categorize people. When we can see value in someone, we are more likely to be outraged if he or she is in any way subjected to an unhealthy environment. When everyone is considered valuable, no one is ignored.

Prejudice feeds on ignorance; in order to keep your prejudice intact you must be sheltered from too much information about the people who are the targets of your prejudice (Clark, D., 1987). To curtail the political processes that protect people who are in the minority, you need only make them invisible (Mohr, R.D., 1985). What is different about the present era is that larger numbers of people who are experiencing oppressive behaviour by others (e.g. women, elderly people, ill people, handicapped people, gay people) are beginning to organize and become more visible, and they are insisting on the rights that are now more accessible to them because of legislation.

The key to ending a cycle in which people get forced into, or stuck in, unhealthy environments because they happen to be young, or old, or women, or physically disabled, or gay, or anything else, is to spread detailed, accurate information about them to others (Clark, D., 1987). In all cases we discover for ourselves they are really human beings with much in common.

In the past, it was considered sufficient to enact human rights legislation and to hope that discriminatory behaviour would slowly decrease as new generations grew up. Governments at all levels assumed the role of protectors of minority rights, when resources allowed and when there was sufficient political support.

If 'Health for All' is to be achieved, a more active stance against prejudice will be required by every individual and every organization in the country. We must base this stance on the assumption that everyone, given detailed enough information and strong enough incentive, is capable of changing personal attitudes that are not in anyone's best interests. There is no reason why the Canadian population itself could not now begin to assume a more active role in protecting the rights of others, particularly if such behaviour is more likely to lead to a healthier environment for themselves.

3.2 Creating a New Vision of Environmental Choice

Many references in the literature offer new visions involving greater environmental control and choice for individuals (Duhl, L.J., 1969), (Hooker, C.A., 1982), (Repace, J.L., 1983), (Schrecker, T.F., 1984), (Bradley, J., 1987), (Rea, W., 1987), (World Commission on Environment and Development, 1987).

Some emphasize an individual's right to know what he or she is being exposed to (Johnston, A., 1983), (Twigge-Molecey, C., 1987), and the idea of educated consumer choice leading to less-polluting materials (Andersen, I., 1982). Others offer specific designs for alternative, healthier environments (McNall, P.E., 1975), (Sandia National Labs, 1982), (Small, B.M., 1985).

A common message in much of the literature is that being in control of our lives contributes to good health (Miller, S.M., 1980), (Insel, P.M., 1983b). Another message is that Canada is a nation of diversity, and that new approaches will be required to accommodate this diversity in a way that allows everyone a healthy environment (see also Section 2.3.4 "Accommodating Diversity Through Participation").

The vision required to implement 'Health for All', and in particular, healthy environments for all, is one in which, individually and collectively, we regain control over our external environments. This means empowering individuals and entrenching their rights to environmental control in many settings where they are now lacking (e.g. school health and safety, tenant health and safety), and improving their degree of control in other situations (e.g. industrial and commercial settings). On a wider scale, it means regaining control, as a society, over the introduction of new materials and new technologies into our lives, with the power to accept or reject based on solid information about their effects, both positive and negative.

The bottom line in a vision of environmental choice is information. Everyone in Canada has the right to better access to detailed information about his or her environments and the potential effects of those environments on health. Citizens in every setting are in a better position to take control and responsibility for ensuring that their own environments and those of others they affect are healthy ones, when they are knowledgeable about the 'ecology of health'. When citizens are better empowered environmentally, they tend to create for themselves the environments that suit their particular needs.

The answer to the question "what is a healthy environment for Canadians?" is that we need a great deal of flexibility and choice in our environments, because our needs and wants are diverse. Our common environments require more clever design, to accommodate a greater range of needs, a wider range of vulnerability to hazards, and a richer mix of human characteristics than we have acknowledged before. Our specialized environments need to be responsive to the specific characteristics of those who need them most, but must not ignore the full humanity of those whom they serve.

3.3 Fostering Technology for Assessing our Environment

We are fortunate that our era of rapid technological change, which has contributed to some of our environmental hazards, has also supplied us with new and better means to solve environmental health problems. There are a number of new technologies, all maturing over the last ten years, that together revolutionize the field of physical environmental health. They provide quantifiable information about the connection between environmental contamination and health, that was never available previously. They are beginning to resolve the decades-long debate as to whether low-level pollution can affect physical and mental health, behaviour and performance. In fact, they remove any doubt that we are affected by our surroundings.

The first breakthrough was the refinement of technology for measuring contaminants in air and water down to ultra-trace levels, confirming what kind of environment we are exposed to (Dravnieks, A., 1971), (Berglund, B., 1982), (Hijazi, N., 1983), (Meyer, B., 1983), (Nagda, N.L., 1983), (Wadden, R.A., 1983). Manufacturers are developing many new forms of this technology in response to growing demand. There is still an overwhelming need to extend this technology by devising economical, standardized, calibrated instruments that can be used by people with relatively little training, to measure the full array of contaminants associated with the sick building syndrome (Akimenko, V.V., 1986).

The second breakthrough was the improvement of technology for measuring contaminants in body tissues and fluids, using sensitive and selective techniques of high-resolution gas chromatography and high-resolution gas chromatography-mass spectrometry, confirming that we do indeed absorb contaminants from our surroundings (Laseter, J.L., 1983), (Root, D.E., 1986). The method has been used to test for the presence of different arrays or families of chemical contaminants, including volatile hydrocarbon solvents, and chlorinated hydrocarbon pesticides and their common metabolites. In a screening series involving patients who exhibited chemical hypersensitivity, 99% had pesticide residues at or above the .05 ppb level in their sera (Laseter, J.L., 1983). The 'footprint', or array, of contaminants can be compared to similar measurements of the individual's environments, to determine possible sources of the absorbed contaminants.

The third breakthrough was the enhancement of brainwave measurement technology, with the use of high-powered computers, to the point that we can measure subtle brainwave changes of individuals in response to different environments (Bokina, A.I., 1976), (Haider, M., 1976), (Thatcher, R.W., 1985), (Kaye, H., 1987). This computerized diagnostic system provides physicians with a window into the brain, providing a high resolution colour image of the brain's electrical activity as well as complete quantification and statistical analysis of all electrical measurements. This form of testing can be used to evaluate persons who have been exposed to environmental chemicals. Preliminary testing has provided support for the contention that there are specific changes in brain function, that accompany cognitive or affective shifts, in patients who are sensitive to exposure to common foods and odours (Kaye, H., 1987).

The above triad of methods can be used together, to confirm the effects of specific chemical exposures on specific individuals: we can find out what a person is exposed to; we can obtain proof of exposure in the form of absorption; and we can determine how he or she responds to the exposures.

A fourth technology was invented a great many years ago, but is nevertheless proving to be a potent diagnostic and therapeutic tool in cases of illness suspected to be related to environmental exposures. It involves simply placing a person who is suspected of being adversely affected by his or her environment, into an alternative cleaner and healthier environment, to observe whether any differences in response can be measured (Rea, W.J., 1978), (Rea, W.J., 1984). In one study using this approach, patients with non-arteriosclerotic cardiac arrhythmias and/or chest pain refractory to medication, having various associated symptoms relating to smooth muscle sensitization, were studied in a rigidly controlled, relatively fume-free and particle-free environment. The majority of signs and symptoms were cleared in most of the patients, without medication, while under environmental control. Arrhythmias were reproduced in most of the patients, with controlled, repeated, individual blind and double-blind, incitant challenges (Rea, W.J., 1978). Expertise for creating suitable alternative environments is being developed in Canada (Small, B.M., 1983).

A fifth technology involves using an array of objective neurobehavioural and performance measurements to ascertain individual human reactions, other than brain electrophysiology, in response to environmental stressors including chemical pollutants. A large number of chemicals are known to have direct toxic effects on the central nervous system (Pryor, G.T., 1983), (Anger, W.K., 1984). Testing of people being exposed to a mixture of common indoor air pollutants indicated that mental performance can be impaired by indoor air pollution, and that the effects of low concentrations of organic gases and vapours can be measured objectively (Bach, B., 1985).

Finally, analytical methods in immunology hold a great deal of promise for further clarifying the effect of different environments on human beings. For example, certain chemical compounds such as polycyclic aromatic hydrocarbons (PAHs) have been shown to cause immune suppression in mice within two days after exposure. Animal studies have shown profoundly suppressed antibody response after exposures which produced no mortality or other obvious clinical symptoms of general toxicity (Dean, J.H., 1983).

Such new and powerful techniques bring us into an era in which it is finally possible to determine specific acute and chronic effects of the environment on individuals. Previously, major issues of chemical contamination and other environmental insults were examined primarily in terms of epidemiological comparisons, using surveillance of possibly-related, but not conclusively-related symptoms (Elinson, L., 1984), (Spitzer, W.O., 1984).

It is time, therefore, to stop arguing over whether environment is an important factor in health. Our new task is to further develop and apply these and other similar technologies, to quantify environmental effects and test alternative environments, until we find, and verify, healthier ways of living.

3.4 Accommodating Diversity Through Participation in Design

Many of our surroundings were designed without taking into account the full diversity of the population that would use them. One of the reasons that many people find themselves in unhealthy situations is that they belong to the group of people who were not considered when the situation was created. Many public and private environments were designed and built to standards which do not take into account even modest differences in people's age, physical capabilities, lifestyles, tolerance for pollution, and many other characteristics (Rand, G., 1985), (Bradley, E., 1986), (Bradley, E., 1987).

In fact, people do vary widely in their response to environmental stressors (Cooper, W. Clark, 1973), (Calabrese, D.V., 1986), and even different organ systems in one individual may have markedly different responses to the same stressor (Voronova, B.Z., 1980). The mix of characteristics within the population (demographics) also changes with time, for example, as the number of older people increases (Last, J.M., 1987). All these variations are important for long range health planning.

Diversity tends to be ignored through homogenization and invisibility. Lumping people with one common characteristic together may simplify design of physical or social environments, but it ignores their remaining characteristics, which may vary widely (Bradley, E., 1987). Ignoring the existence of people who are in the minority, or hoping that the number of exceptions to the norm is small, helps to keep diversity invisible. Prejudicing against individuals on the basis of one or more of their characteristics also virtually guarantees that those characteristics will be suppressed or hidden, as much as possible, to the detriment of the individual.

A number of researchers have suggested, instead, that the full variability of response of people to environmental factors can be used to advantage, in establishing better health and safety standards, and in detecting in advance those environmental conditions which will prove, over time, to have adverse consequences for many people (Calabrese, E.J., 1978), (Plumlee, L., 1979), (Ostapovich, I.K., 1975).

One consistent theme in the literature is the need for participation in design of both physical and social environments, by people who will be using them (Wolfe, M., 1976), (Johnston, A., 1983), (Innes de Neufville, J., 1985), (Church, K., 1986), (Lecuyer, G., 1987), (Twigge-Molecy, C., 1987).

The antidote to ignoring diversity, similar to the antidote to prejudice, is more and better information. Direct involvement of people who are affected by any development, whether it be physical or social, will shatter the stereotypes used in design and guarantee that all human characteristics that must be taken into account are highly visible. To achieve this, we discard the assumption that professionals who design the various pieces of our society know the people they design for well enough to proceed without further discussion or information. We must further assume that people want to become, and are fully qualified to become, active partners in the design of environments that will affect them.

3.5 Becoming World Experts in Designing Healthy Environments

Recent reviews emphasized that economy and environment must be integrated, since long term economic growth depends on a healthy environment, and in turn affects our environments in many ways (Lecuyer, G., 1987), (Mustard, F.J., 1987), (World Commission on Environment and Development, 1987). Other reviews indicate that there is a growing international awareness of the need for new approaches to health (World Health Organization, 1986), (Hancock, T., 1987).

If Canada is to achieve 'Health for All' and healthy environments for all, it must be prepared for physical, social, and institutional changes that take into account the actual connection between the physical and social environments and health. These changes will involve new types of social design which take into account everyone's full value, including previously devalued or merely tolerated minorities. They will require more flexible physical designs and safer, less-polluting products and materials, which accommodate the full range of diversity of people, and the latest information on the effects of environments on health.

A nation cannot accomplish such a feat without learning a great deal. What we want in terms of healthy environments, the rest of the world also wants — we are all fully human and our similarities outnumber our differences. Every healthy physical design, every healthy social structure, every non-polluting product or material, every well-thought-out device that is built not to harm its users, every method for involving users in design, every treatment for the adverse effects of unhealthy environments, every piece of information about achieving healthier environments, will be in demand worldwide.

In an age of intense industrial competition, in an era when advanced technologies enable many countries to produce sophisticated and valuable goods, each country seeks a niche, a way of distinguishing its products and services from those of others. Canada has always been associated with clean, peaceful living and competent technical design. Learning the 'technology' of healthy environments will add one more reason for other countries to look toward Canada for products and services to fill their needs. The incentive for Canadian manufacturers to create healthier products, for example, might be government assistance in finding export markets, under a 'Canadian Means Healthy' campaign.

The bottom line is therefore one of opportunity. Achieving healthy environments is, in the long run, not a cost but an investment, both in our domestic resources (everyone in Canada) and in our potential for exporting goods, services and knowledge to the rest of the world. If the job must be done, Canadians can do it well and become leaders in the field.

PART III

DETAILED BIBLIOGRAPHY AND ABSTRACTS

This appendix lists and reviews the references that are cited in PART II: ANALYSIS, as well as additional references that were consulted during the study and which had a bearing on the analysis and overview.

Most of the references are available by inter-library loan through the services of local public and university libraries. The Canada Institute for Scientific and Technical Information (CISTI), Ottawa, Canada, the National Technical Information Service (NTIS), Washington, D.C., USA and the National Library of Medicine (NLM), Bethesda, Maryland, USA will have copies of most sources.

In most instances, further bibliographic information, including ISBN numbers, special report numbers, and addresses of author or publisher organizations, is available from the computer data bank of the Foundation for Independent Research on Technology and Health, c/o Sunnyhill Research Centre, R.R.#1, Goodwood, Ontario L0C 1A0, Tel: (416)-294-3531.

References have been listed alphabetically by the first author. All authors are indexed in the "Name Index" section, and all keywords as well as other principal words and phrases in the abstract have been indexed in the "Subject Index" section of this report.

Abel, Cora Beth (1981). What Is An Adaptive Environment?. Adaptive Environments, Massachusetts College of Art, Boston MA; 1981.

Most of us believe we have no opportunity to design or shape our environment. Professionals often make design decisions in isolation, although the environment has an awesome impact on our lives. When people work in an environment where outside noise affects their ability to concentrate, they usually react by cutting back their sensitivity to the environment rather than boldly acting to change it, muffling the noise. An adaptive environment is defined as a positive relationship between the physical setting and the people who use that setting. It is possible when a) the users are skilled enough to use basic design tools to make small changes as their individual and group needs change; b) the environment supports specific activities that occur in the space for all users regardless of age; size; mental, physical or emotional state; and c) users and designers together assess what works and what fails to meet users' requirements, with successes and failures documented for future reference in designing similar settings. An adaptive environment is accessible and barrier-free to a much greater degree than normal. It is also free from the effects of attitudinal barriers that lead to insensitive and elitist design and supports all people's daily activities on the physical, psychological, and emotional levels. It allows all users to participate to the fullest extent possible in activities that occur within the space.

Adam, Barry D. (1978). *The Survival of Domination: Inferiorization and Everyday Life.* Elsevier North-Holland, Inc., New York, 1978.

Dominated peoples develop a range of behaviour patterns to cope with their recalcitrant social environment. The author describes the reactions to oppression in everyday life, of individuals who are subordinated, by concentrating on three groups: Jews, blacks and gay people. He notes that how people survive domination through resistance, accommodation and compliance tells us much about how domination survives and how inequitable social order is reproduced. He comments on the difference between liberal tolerance and true acceptance, the former merely transforming an inferiorized group from criminals to minorities or deviants. Assimilation promises a 'deal', wherein the identity, culture, and values of the inferiorized are to be negated, or at least concealed, in return for the promise or opportunity of improved life chances.

Adam, Barry D. (1987). *The Rise of a Gay and Lesbian Movement.* Twayne Publishers, Boston, MA, 1987.

The author reviews human rights issues for lesbian and gay people, including child custody, tax and family laws, municipal zoning regulations, social welfare, health care, insurance and employment. The present AIDS crisis has also highlighted the issue of 'spousal rights' and distribution of property upon death. He points out that prejudices against homosexuality have led to obstacles being presented against distribution of literature about 'safer sex', a term used to describe sexual practices which minimize the risk of transmission of sexually transmitted diseases such as AIDS. He comments on the 'immense cost' to individuals of the psychological suffocation and fear, suffered by those in hiding.

Adelman, Richard C. (1979). *Loss of Adaptive Mechanisms During Aging: from the Symposium, Overview of the Biology of Aging.* Federation Proceedings, Vol. 38, No. 6; May 1979; Federation of American Societies for Experimental Biology.

One general feature of all aging populations is the progressively modified ability to adapt to changes in the surrounding environment. In this study, factors contributing to modifications in the capability for enzyme adaptation as an expression of aging are reviewed. Specific examples of altered enzyme adaptation include the responses of hepatic glucokinase activity to glucose and of hepatic tyrosine aminotransferase activity to starvation in rats. These reflect disturbances in hormonal regulatory mechanisms.

Adkinson, N. Franklin Jr. (1977). Environmental Influences on the Immune System and Allergic Reactions. *Environmental Health Perspectives*, Vol. 20, pp. 97-103; 1977.

Environmental interactions with the immune system may result in two types of adverse outcomes: immunodeficiency and immunopathology. Serious immunodeficiency most commonly results from ionizing radiation or as a recognized side effect of iatrogenic drug therapy, usually cancer chemotherapy. Environmentally-triggered immunopathology is a source of considerable morbidity and mortality. Additional research is needed, particularly in the area of identification of risk factors which predispose to immunopathological outcomes when individuals are exposed to sensitizing chemicals or other natural allergens.

Adkinson, N. Franklin (1977). Environmental Influences on the Immune System and Allergic Reactions. *Environmental Health Perspectives*, Vol. 20, pp 97-103; 1977.

Environmental interactions with the immune system may result in two types of adverse outcomes: immunodeficiency and immunopathology. Serious immunodeficiency most commonly results from ionizing radiation or as a recognized side effect of iatrogenic drug therapy, usually cancer chemotherapy. At present there is little basis for believing that biologically significant suppression of immune competence results from more subtle interactions with environmental agents. On the other hand, environmentally-triggered immunopathology is a source of considerable morbidity and mortality. Additional research is needed.

Akimenko, V.V.; Andersen, L; Lebowitz, M.D.; Lindvall, T. (1986). The 'Sick Building' Syndrome. Evaluations and Conclusions for Health Sciences and Technology; Indoor Air, Volume 6; pp 87 - 97; Swedish Council for Building Research; 1986.

There have been some advances in our knowledge of sick buildings during the last few years. The syndrome can only be diagnosed by building user complaints and specific causes of complainant symptoms have usually not been identified. Guidelines for future research needs are presented. These include an overwhelming need to devise better, standardized, calibrated instruments to measure health effects associated with the sick building syndrome. There is also a need for technical studies of the relative importance of the various chemical sources of pollutants in different types of buildings in order for preventive measures to be introduced. There is a need for simple and inexpensive methods for the study of ventilation rates and efficiency. Threshold limit values developed to protect workers in the industrial setting should not be used in the non-industrial setting.

Allen, Max; Armstrong, Russell; McPhee, James (1987). Censorship Bulletin #6. Canadian Committee Against Customs Censorship; June 1987.

The Canadian Committee Against Customs Censorship was formed to fight seizures of lesbian and gay books by Canada Customs. The committee makes the argument in this publication that customs guidelines are being used to discriminate against gay-related reading materials, and that to do so will deny many people accurate information on homosexuality, lesbianism, and sexuality in general. They suggest that the provision of detailed information on 'safer' sex for all individuals, and especially for young people, is a necessary component in the fight against Acquired Immune Deficiency Syndrome (AIDS).

Alphey, R.S.; Leach, S.J. (1974). Accidental Death in the Home: Building Research Establishment Current Paper. Building Research Station, Dept. of the Environment, Great Britain; 1974.

6,500 accidental deaths in homes in England and Wales were studied for the period 1950 to 1972. The deaths in 11 categories, ranging from the largest which was falls, to categories with small numbers, such as excessive cold or electrocution, are plotted and trends examined. Marked seasonal fluctuations of up to 100 percent between winter and summer occur, suggesting that dwellings are not adequately performing their function of isolating the occupants from the effects of climate. The risk to an individual from accidental death in the home is between a third and a half of that from death in a road traffic accident for those under 65.

Altman, Irwin; Werner, Carol M. (1985). Home Environments: from Human Behavior and Environment; Advances in Theory and Research; Volume 8. Plenum Press, New York; 1985.

This is an intensive treatment of the home environment. Some chapters span many cultures, others focus on homes in industrialized nations or developing nations, some deal with the indoor environment exclusively, while others focus on the neighborhood and country in relation to homes. Also analyzed are the broad societal forces such as housing services and political pressures.

Ames, Bruce N. (1983). Dietary Carcinogens and Anticarcinogens: Oxygen Radicals and Degenerative Diseases. Science, Vol. 221, No. 4617, Sept. 23 1983; American Assoc. for the Advancement of Science, 1515 Massachusetts Ave., NW, Washington DC 20005.

Ammann, Harriet M. (1987). Effects of Indoor Air Pollution on Sensitive Populations. *Clinical Ecology*, Volume V, No. 1, pp. 15-21; 1987.

This paper, prepared by a physiologist and health scientist at the U.S. Environmental Protection Agency, reviews health hazards and physiological effects of indoor air pollution on populations at higher than average risk, including infants and the elderly, pregnant women and their fetuses, anemic and asthmatic persons, those with compromised cardiovascular or pulmonary function, and siblings of sudden infant death syndrome victims. In the aggregate, these persons make up a considerable portion of the public. Considering time exposure, their risk may be considerably greater than that calculated using workplace exposure factors.

Mimicry of common illnesses, including influenza, food poisoning, gastrointestinal disorders, Alzheimer's disease, angina or brain deterioration, results in misdiagnosis of intoxication from indoor air pollution and the underestimation of its occurrence.

Andersen, Ib; Lundqvist, G.R.; Molhave, L. (1975). Indoor Air Pollution Due to Chipboard Used as a Construction Material. *Atmospheric Environment*, Vol. 9, No. 12, pp. 1121-1127; 1975; Pergamon Press.

Chipboard (particle board) is a common building construction material made of wood shavings held together with a urea-formaldehyde glue. Due to this composition, there is a continuous emanation of formaldehyde. Measurements in 25 rooms in 23 Danish dwellings where chipboard was used in walls, floors and ceilings showed that the average concentration was 0.62 and the range 0.08 - 2.24 mg formaldehyde m³ air, exceeding the German threshold limit for occupational exposure. In all rooms, the concentration exceeded the German limit for continuous exposure for outdoor air. The adverse health effects of low levels of formaldehyde are irritation of the upper airways and conjunctivitis. The need for air quality standards and control programs for indoor air in the home is stressed.

Andersen, Ib; Lundqvist, Gunnar R.; Proctor, Donald F.; Swift, David L. (1979). Human Response to Controlled Levels of Inert Dust. *American Review of Respiratory Disease*, Vol. 119, pp. 619-627; 1979.

The authors studied nasal mucous flow, airway resistance, and subjective response in 16 young healthy subjects during 5 hour exposures to 2, 10, and 25 mg of inert dust per m³ in an environmental chamber. The dust was fully polymerized plastic dust containing carbon black. The number of these particles in room air, expressed as a per cent of the total number of particles was 36, 41, 14, 7, and 2, respectively, for the aerodynamic size ranges less than or equal to 1.8, 1.9 to 5.3 to 8.9, 9.0 to 12.4, and greater than or equal to 12.5 μ m. No significant changes in nasal mucociliary clearance rate or nasal resistance were observed.

At all dust concentrations there was a decrease in 1-sec forced expiratory volume, but not in the forced vital capacity or the forced expiratory flow during the middle half of the forced vital capacity. The nasal penetration fraction of particles was approximately 55 percent for the smallest particles and 20 percent for the largest. Discomfort was proportional to the concentration of dust, but lagged almost 2 hours behind the changes in dust concentration. The discomfort was never excessive; the main complaints were dryness in the nose and pharynx.

Andersen, Ib (1979). Formaldehyde in the Indoor Environment—Health Implications and the Setting of Standards: Indoor Climate - Effects on Human Comfort, Performance, and Health in Residential, Commercial, and Light-Industry Buildings; Fanger, P.O., Valbjorn, O.—editors. Danish Building Research Institute, Copenhagen; 1979.

Exposure to formaldehyde vapour causes irritation especially of the eyes and upper airways. Skin irritation may also occur. The background concentration in outdoor air is about 0.05 mg/m³. Indoor concentrations up to 1-2 mg/m³ have been found in rooms with emanations from construction materials made of resins. The literature on the biological effects of formaldehyde is reviewed and the results of an exposure of 16 young healthy subjects to 0.3, 0.5, 1.0 or 2.0 mg formaldehyde/m³ air during 5 hours are described. There was no change in airway resistance. Eye irritation and dryness in the nose and throat was experienced by 3 subjects at 0.3 and by 15 at 1.0 mg/m³. A standard for continuous exposure protecting all but subjects sensitized to formaldehyde against any adverse health effect and the majority of the subjects against discomfort is suggested at or lower than 0.15 mg formaldehyde/m³ air.

Andersen, I; Seedorff, L.; Skov, A. (1982). A Strategy for Reduction of Toxic Indoor Emissions. Environmental International, Vol. 8, pp 11-16, 1982; Pergamon Press Ltd.

Many building materials emit pollutants. To reduce the exposure of the population to the major groups of toxic indoor pollutants, a strategy is put forward which concentrates on a reduction of the emission of carcinogenic substances, eye-airway irritants, and odors from building materials. It is the experience of the Danish National Inventory of Toxic Substances and Products that preventive measures in the form of either a complete removal of a substance or replacement of a toxic substance by a less toxic substance are possible. Among the banned or regulated substances in Denmark are asbestos; epoxy and polyurethane products; and another 1771 building materials including sealants, glues and adhesives, paints and lacquers, and wall/floor coverings. Based on an analysis of publicly available information on the labels of these products, the authors conclude that a notification system makes it possible to select the products with the least impact on human health and comfort, and, consequently, with the lowest need for ventilation. It is suggested that these toxicological principles should be used for the improvement of existing and future building materials.

Anderson, Darrell E. (1971). Problems Created for Ice Arenas by Engine Exhaust. *American Industrial Hygiene Association Journal*, No. 32, pp. 790-801; December 1971.

Episodes of illness in separate arenas from carbon monoxide and nitrogen dioxide exposures are described. Typical symptoms have been headache and nausea among children and headache among adults. In one episode, symptoms of illness were sore throat and tightness in the chest. Some of the unrecognized effects of low-level carbon monoxide exposure such as disturbance of coordination, judgment, psychomotor tasks, and visual acuity, and marked increases in choice discrimination errors and reaction time, which have been pointed out by recent medical research, are of added concern. Excessive air concentrations of these gases resulted from the operation of ice resurfacing machines. Carbon monoxide concentrations up to 250 ppm following one episode and nitrogen dioxide concentrations up to 40 ppm following the other were found. All 45 ice arenas in Minnesota were visited subsequently to evaluate the variables responsible. Engine exhaust and ambient air data are presented. Infrared analyzer recordings of carbon monoxide concentrations in three arenas are shown. Possible methods of controlling exhaust gas problems are discussed.

Anderson, Joan M. (1986). Ethnicity and Illness Experience: Ideological Structures and the Health Care Delivery System. *Soc. Sci. Med.*, Vol. 22, No. 11, pp 1277 - 1283; Pergamon Journals Ltd.; 1986.

This paper analyzes the experiences of Anglo-Canadian and immigrant Chinese families with a chronically ill child by using the idea that the social organization and ideology of health care services generate particular illness experiences. Immigrant families find that the ideology is not compatible with their customs for managing illness. This often leads to non-compliance and ineffective treatment. Health professionals explain non-compliance by the obvious facts of cultural differences, but the author argues that it should be understood by institutional practices that exclude families from participating in caretaking. Patients and families should be included in decisions that affect their lives. Pressures from government to economize by increasing home care services, and the increasing number of immigrants may force practitioners to negotiate culturally acceptable care with them.

Anger, W. Kent (1984). Neurobehavioral Testing of Chemicals: Impact on Recommended Standards. *Neurobehavioral Toxicology and Teratology*, Vol. 6, pp. 147-153; 1984.

Historically, the American Conference of Governmental Industrial Hygienists has served as a major source of information on recommended safe exposure limits (Threshold Limit Values or TLVs) for chemicals most frequently encountered in industry and of known toxicologic significance.

A review of the ACHIH Documentation Book, which details the basis for their judgements, has indicated that 167 of the 588 chemicals for which they assigned recommended standards have, as one of their bases, direct nervous system effects.

Angus, Douglas E.; Manga, Pran (1986). National Health Strategies: Time for a New "New Perspective": special article. Canadian Journal of Public Health; Vol. 77, pp 81 - 85; March/April, 1986.

A decade has passed since "A New Perspective on the Health of Canadians" was published by Liberal Health Minister Lalonde. But current information suggests that many of our health problems have stayed much the same. In addition to looking at overall life expectancy as an indicator of health status, one should consider the more comprehensive healthfulness of life. Looking at health from this perspective could lead to big changes in the relative priority accorded to each cause, with all the implications for health planning and policy making that would entail. There need to be firmly articulated goals and objectives, principles and rules governing the design of programs in pursuit of these goals, and a clear notion of resources that should be allocated for implementation. Also as part of a rational planning process, health needs must be assessed, priorities must be established, progress monitored and programs evaluated. Information is important, as is good communication between users and producers of health data.

The term lifestyles has been used as a synonym for personal behaviour. Social, economic and collective determinants of lifestyles tend to be ignored. This partial and restricted conceptualization of the problems is at odds with the broader message in the New Perspective. There is a danger that isolating lifestyle and personal responsibility will lead to frustration on the part of those who want a broader change, and may fail to gain the acceptance of the target groups, if not the general public.

Universal health insurance has succeeded in bringing about a high degree of equity in access to services, but did little to increase consumer responsibility and awareness for personal health care. There is no doubt that people in the bottom income groups have lower health status than those at higher income levels despite equitable access to services. This paradox may become more significant over time, since lifestyle modification has been more effective among the higher educated and income groups.

Appleyard, Donald (1976). Transportation as a Social Environment: Can We Change a Tradition?. Transportation Planning for a Better Environment, Vol. 1, section 5; Stringer, Peter and Wenzel, H., ed.; New York: Plenum Press; 1976.

A revolution is taking place in transportation planning due to public disenchantment with traditional planning methods and the closed nature of the old transport planning process. This demands a restructuring of

basic concepts and values. Transportation must be seen as a service rather than a facility and should be user- and neighbour-oriented. Systems must be seen as complementary rather than competitive. The planning must be multi-functional and multi-disciplinary and needs to be more open to public participation than the present closed professional system of planning. Environmental and social professionals, environmental planners, landscape planners, urban designers, sociologists, community organizers, political scientists, psychologists, and ecologists should be hired as full-time staff members in transportation agencies. Every discipline suffers from isolated specialization, a phenomenon that is destructive to the quality of urban life.

Appleyard, Donald (1979). The Environment as a Social Symbol: Within a Theory of Environmental Action and Perception. *APA Journal*, pp 143 - 153; April 1979.

Professionals and social scientists tend to screen out the connections between the physical environment and its social meaning. In this paper, a communications model of environmental action and perception is developed and elaborated. An environment becomes a social symbol when it is intended or perceived as a representative of someone or some social group. This can be seen when upper middle-class whites move into an area formerly inhabited by lower-class blacks and "improve" it, planting trees, and renovating buildings, to the distaste of the old residents. The attributes of environmental actions in different contexts, and the differing modes of perceiving and interpreting environments, especially during environmental conflicts and in home environments, when social meaning is dominant, are explored. The implications for public policy include a) physical planners and designers should be much more aware of the fact that the need for identity, recognition, and even some sense of power is a human need which has a necessary outlet for expression in the physical environment; b) physical planning decisions can threaten the identity and status of certain groups while enlarging the powers of others; and c) the significance of citizen participation in environmental decisions is critically important, because this is the way in which people can become identified with a new environmental action, the way in which they can possess and feel responsible for it, thus reducing their alienation. An extensive bibliography is included.

Arlien-Soborg, P.; Zilstorff, K.; Grandjean, B.; Milling Pedersen, L. (1981). Vestibular Dysfunction in Occupational Chronic Solvent Intoxication. *Clinical Otolaryngology*, Vol. 6, No. 4, pp. 285-290; 1981; Blackwell Scientific Publications.

Neurotoxic volatile organic solvents used by house and car painters may lead to professional toxic encephalopathy after several years of exposure. The symptoms are memory impairment, fatigue, personality changes, headache and dizziness. Vestibular dysfunction was found in

55 percent of 113 painters examined, mainly in the form of reduced caloric vestibular reactions. No correlation between vestibular dysfunction and the duration of exposure, cerebral atrophy or intellectual impairment could be demonstrated. Vestibular examination may be helpful in detecting early changes in exposed persons and in determining more accurate safety limits for harmful chemicals.

Aronow, Wilbert S. (1978). Effect of Passive Smoking on Angina Pectoris. New England Journal of Medicine, Vol. 299, No. 1, pp. 21-24; July 1978.

The effect of passive smoking on exercise-induced angina in a well ventilated and in an unventilated room was evaluated in 10 patients with angina. Patients exposed to 15 cigarettes smoked within two hours in either room increased their resting heart rate, systolic and diastolic blood pressure, and venous carboxyhemoglobin and decreased their heart rate and systolic blood pressure at angina. Patients exposed to passive smoking in an unventilated room had a larger increase in resting heart rate, systolic and diastolic blood pressure, and venous carboxyhemoglobin and a greater reduction in heart rate and systolic blood pressure at angina.

Asher, Janet K. (1977). Toward a Safer Design For Stairs. Job Safety & Health, Vol. 5, No. 9, pp. 27-32; 1977; U.S. Dept. of Labor, Occupational Safety and Health Administration, Washington DC.

Stairway accidents are largely caused by design shortcomings. A study of 11,000 stairway accident reports and videotapes of 30,000 people using stairs which utilized stop-motion analysis indicated that victims typically slip while descending stairs. Seven guidelines for constructing, retrofitting and use of stairs are presented. These include minimizing the use of public stairways; removal of distractions, conspicuous treads and handrails; extended handrails; elimination of deceptive floor coverings and shadows; proportion that fits the users; tight tread coverings; internally stable and slip-resistant surfaces for good traction.

Axelsson, Olav; Hane, Monica; Hogstedt, Christer (1976). A Case-referent Study on Neuro-psychiatric Disorders Among Workers Exposed to Solvents. Scand. j. work environ. & health 2, pp. 14-20; 1976.

Published reports give justification for the belief that long-term exposure to solvents might induce chronic but nonspecific neuropsychiatric conditions. This case-referent study of data from a regional Swedish pension fund register indicated a risk ratio of 1.8 in regard to nonspecific neuropsychiatric disorders among workers such as painters, varnishers and carpetlayers who are exposed to solvents as compared to workers not so exposed. Moreover a dose-response relationship seems to exist between exposure in terms of occupational years and neuropsychiatric conditions.

Axelsson, Olav; Gustavson, Jan (1978). Some Hygienic and Clinical Observations on Styrene Exposure. *Scand. j. work environ. & health* 4, suppl. 2, pp. 215-219; 1978.

Styrene exposure levels in the range of 100 to 300 ppm have been measured in small shops manufacturing glass fibre reinforced plastic products such as boats, steeping baths, etc. Exposure control through determinations of mandelic acid in the urine at the end of the workday have been suitable and convenient. Over the years during the 1970s, there seem to have been amendments in the exposure situation in Sweden if judged on this basis. More or less severe central nervous disturbances have been observed among the workers.

Axelsson, Olav; Edling, Christer; Kling, Hans (1979). Lung Cancer and Residency—A Case-referent Study on the Possible Impact of Exposure to Radon and its Daughters in Dwellings. *Scand. j. work environ & health* 5, pp. 10-15; 1979.

In view of the well-known urban-rural difference in lung cancer rates, remaining also after standardization for smoking, it is suggested that low levels of radon and its daughters in dwellings might be of etiologic importance to this disease. To test this hypothesis, a study was undertaken in a rural area; it considered residency in wooden, mixed type, and stone houses among cases of lung cancer and a control group. The results indicate an increased risk of lung cancer among residents in mixed type and stone dwellings. Additional studies are highly desirable to confirm or refute these findings, which, if valid, mean increasing lung hazards caused by a decrease in ventilation in future energy saving unless special measures are undertaken to reduce radon daughters in dwellings.

Bach, Bodil; Molhave, Lars; Pedersen, Ole F. (1985). Human Reactions During Controlled Exposures to Low Concentrations of Organic Gases and Vapours Known as Normal Indoor Air Pollutants: Performance Tests. *Proceedings, 3rd International Conference on Indoor Air Quality and Climate, held in Stockholm, Sweden, August 20-24, 1984. Volume 3: Sensory and Hyperreactivity Reactions to Sick Buildings*, pp. 397-402.

Human subjects suffering from indoor climate symptoms were exposed to various concentrations of a mixture of 22 common indoor air pollutants. Among several objective and subjective measurements, performance tests and investigations of irritation to the trigeminal nerve endings were undertaken. The digit span test, testing memory impairment, showed significantly decreased scores during exposure to organic gases and vapours, whereas the graphic continuous performance test, testing the ability to attend and concentrate, showed no effect of exposure. No significant effects on the trigeminal nerve endings were found. The authors conclude that indoor air pollution seems to impair mental performance, and that the effects of low concentrations of organic gases and vapours can be measured objectively.

Bakacs, T. (1972). Urbanization and Human Health. Akademiai Kiado, Budapest; 1972.

The growth of cities has had an impact on human environment and ecology. There are many factors of urbanization which induce damages to health, including air and water pollution, unsolved issues of sanitation, urban noise, increased risk of exposure to carcinogenic substances, and the stressors of city life. In comparison with the mass destruction caused by epidemics, these micro-hazards seem to be of minor importance, still they become very important if their chronic, cumulative effect is considered. These damaging aspects of the urban environment are discussed in detail in an international context, accompanied by statistics. Suggested general solutions are proposed. Extensive references are included.

Baldassare, Mark (1975). The Effects of Density on Social Behavior and Attitudes. American Behavioral Scientist, Vol. 18, No. 6, pp 815 - 825; July/ August, 1975.

The results of a 1965/66 study in Detroit were analyzed using two personality scales termed "powerlessness" and "need for affiliation". It appears that dense urban neighborhoods may be related to low neighboring among residents. Crowded urbanites may withdraw from local contacts with neighbors as a specific adaptation to immediate high levels of stimulation. More detailed data on social relations are needed.

Baldassare, Mark; Fischer, Claude (1975). Suburban Life: Powerlessness and Need for Affiliation. Urban Affairs Quarterly, Vol. 10, No. 3, pp 314 - 326; March, 1975.

Studies were undertaken to examine the differences in behaviour among urban and suburban residents. No substantial evidence was found to back up any hypotheses that suburbanites differ ecologically, demographically, or behaviourally from city dwellers. It was concluded that explanations for suburban life style should be sought elsewhere.

Ball, Ian (1987). Personal Interview, Ian Ball, Ottawa, Ontario. Personal interview, September 5, 1987.

The author was approximately twelve years of age at the time of the interview. He stated that adults did not often take people of his age seriously; that adults commonly treated people of his age 'like kids' and not as people in their own right; and that adults have power over him to force him to do things, like cleaning his room.

Ball, Roger; Smith, Roy (undated). Design Criteria for the Control of Health Hazards in Schools. Alberta Workers' Health, Safety and Compensation, Occupational Hygiene Branch, Edmonton Canada.

This manual sets basic design criteria for ensuring a healthy working environment and is to be used in the planning of new school facilities as well as the upgrading of old ones. It concentrates on those areas in schools where potentially hazardous conditions are most often encountered. It covers: ventilation, noise, illumination, volatile substance storage, woodworking shops, automotive shops, fine arts, music rooms, science laboratories, changing rooms, general classrooms, and various occupational training areas.

Balser, Diane (1987). Women: Their Present Situation in the World: Detailed Proposals for Policies and Actions; The Final Summary Report of an International Women's Conference, The Netherlands, October 12-17, 1984. Sisters, No. 8, pp. 59-80; 1987; Rational Island Publishers, PO Box 2081, Main Office Station, Seattle, Washington 98111, USA.

The author describes the clear challenges to the women's liberation movement as the complete elimination of sexism, and women becoming the decisive force in determining the future of mankind. The Conference concluded that wherever possible, it should be asserted that the cause of sexism is not individual people, but the institutionalization of oppressive relationships in society, and that men, while conditioned to accept sexist patterns (of behaviour), are inherently good and are also oppressed in the present society.

Sexist oppression of women takes many forms, but the underlying basis is economic exploitation. Throughout the world, most women are poor. With the economic crisis, women more and more are pushed into the ranks of the poor. Full equality between men and women includes equal pay, equal opportunity, equal valuing of the work of both sexes, equal access to all levels of employment, equal distribution of all forms of work (e.g. men and women sharing labour such as parenting and housework, which was traditionally female).

The delegates noted that the majority of women learn to live every day with the fear of physical and emotional violence, in the form of physical beatings by fathers, brothers, and lovers, as victims of rape and other forms of sexual abuse, and as victims of emotional abuse. They state that it is time to change the conditions in our society which perpetuate violence. Eliminating nuclear arms was also cited as an urgent issue facing women and all people.

Bardana, Emil J., Jr. (1980). Formaldehyde - Hypersensitivity and Irritant Reactions at Work and In the Home. *Immunology & Allergy Practice*, Vol. 11, No. 3, pp. 60-71; May/June 1980.

Formaldehyde is a widely distributed low molecular weight compound which is highly soluble in water and capable of binding and altering human protein. The bulk of this material is used in the production of resin compounds employed as industrial laminates, binders and adhesives. Major exposures occur as a result of off-gassing from a variety of wood products and foam insulation. Potential for disease has been well documented in the skin where it can induce dermatitis by irritation, delayed-type hyper-sensitivity and immediate urticarial reactions. Its capacity to irritate the eyes and upper respiratory tract are well established. Observations to date also incriminate formaldehyde in the induction of an inflammatory bronchitis as well as bronchial asthma.

Bardana, Emil J., Jr.; Montanaro, Anthony (1987). The Formaldehyde Fiasco: A Review of the Scientific Data. *Immunology and Allergy Practice*, Vol. IX, No. 1, pp 11 - 23; January, 1987.

Formaldehyde is a chemical with significant medical and industrial applications. Its potential adverse health effects triggered considerable public debate, fueled by the observation of nasal cancer in rodents with high cumulative formaldehyde exposures. In 1979, an attempt was made to analyze the literature on formaldehyde. It was felt that it had the capacity to act as both a respiratory irritant and immunogen. Since that time, there has been a literal explosion of data. This report summarized the new material as it relates to the practicing allergist.

Barker, Robert H. (1975). Additives in Fibers and Fabrics. *Environmental Health Perspectives*, Vol. 11, pp. 41-45; June 1975.

The additives and contaminants which occur in textile fibers vary widely. Synthetic fibers such as nylon and polyester contain trace amounts of contaminants such as catalysts and deactivators which remain after the synthesis of basic polymers. In addition, there are frequently a number of materials which are added to perform specific functions such as traces of metals or metal salts and antistatic agents and flame retardants. After the fibers are converted into fabric form, other substances are applied to act as lubricants, sizing agents, antistats, bleaches, wetting agents, dyes, and durable press treatments.

Barnes, J.M. (1975). Assessing Hazards From Prolonged and Repeated Exposure to Low Doses of Toxic Substances. *British Medical Bulletin*, Vol.31, No. 3, pp. 196-200; 1975.

A low dose is defined as one which by itself is less by one or two orders of magnitude than that dose of the substance that would produce an

unequivocal toxic effect. The levels of activity of bodily systems are determined by so many factors, including chemical substances, essential or non-essential to good health. Some of these may be present at different levels, because of variations in intake and disposal by different members of the population. The acquisition of more information on biological changes in response to low doses of toxic substances must be accompanied by the application of an increasingly better judgement of their significance.

Baroncelli, P.; Battisti, S.; Checcucci, A.; Comba, P.; Grandolfo, M.; Serio, A.; Vecchia, P.; (1986). A Health Examination of Railway High-Voltage Substation Workers Exposed to ELF Electromagnetic Fields. *American Journal of Industrial Medicine*, Volume 10, No. 1, 1986; pp. 45-55; Alan R. Liss, Inc.

The study surveyed health conditions of railway workers active in 258 interconnection and conversion substations all over Italy. Measurements in 220 KV substations showed maximum levels of electric field strength and magnetic flux density of 5 kiloVolts/metre and 15 micro-Tesla respectively; No differences in health measures were found between exposed and control groups, taking account organ systems assumed to be at higher risk. The researchers conclude that workers exposed to extremely low-frequency electromagnetic fields of moderate strength do not show the presence of clear effects on their state of health. They note that an electric field strength of 5 kV/m is not considered dangerous, even in the most restrictive occupational regulations.

Bates, David V.; Sizto, Ronnie (1983). Relationship Between Air Pollutant Levels and Hospital Admissions in Southern Ontario. *Canadian Journal of Public Health*, Vol. 74, pp. 117-122; March/April 1983.

Published hourly data of measurements of particulate pollution from 15 air sampling stations in southern Ontario and computerized hospital admission for the 79 acute care hospitals in the region were examined. The months of January, February, July and August in 1974, 76, 77, and 78 were studied to find possible relationships between the two sets of data. For July and August, highly significant associations were found between excess respiratory admissions and SO₂ and ozone and temperature, with 24 and 48 hour lags between the variables.

Batts Young, Bambi (1981). Chemicals That Cloud the Mind. *Environment and Behavior*, Vol. 1, No.3, pp. 1-3; August 1981; Center for Science in the Public Interest, 1755 S St., NW, Washington DC 20009.

It is not unusual for industrial workers to experience deterioration in behavior or in mental function after exposure to common industrial chemicals and pesticides. In one instance, researchers investigating the effects of "safe" levels of mercury found that a reliable prediction of certain memory problems could be based on a simple measurement of mercury exposure.

Bauman, Kay A.; Hale, F. A. (1985). Bringing the Homosexual Patient Out: Teaching the Doctor's Role. *Medical Education*, Vol. 19, pp. 459-462; 1985.

The authors note that little attention is given in medical school curricula to providing appropriate care for the 5-10% of the population that is homosexual. A comparison was made between individuals taking an elective course involving discussions with articulate homosexual people around societal biases towards homosexuality and issues of health care delivery, and a control group. The study group became more accepting towards homosexual lifestyles significantly in 12 out of 15 measures.

Beaconsfield, P.; Krebs, H.; Borlaug, N.; Rainsbury, Rebecca (1975). Man-made Chemicals and our Milieu Interieur: A Preliminary Report from the Special Commission on Internal Pollution. *Experientia*, Vol. 31, No. 7, pp. 869-872; July 15 1975.

Beall, James R.; Ulsamer, A.G. (1981). Toxicity of Volatile Organic Compounds Present Indoors. *Bulletin of the N.Y. Academy of Medicine*; Vol. 57, No. 10, pp. 978-996; December 1981.

Energy conservation measures reduce movement of air through the home and increase the atmospheric concentration of more than 40 toxic chemicals. In many instances, the effect of chronic exposure to small amounts of these chemicals is unknown. For instance, formaldehyde is a toxic chemical that illustrates the problem. Used in resins that are part of many home products, it causes adverse health effects when it offgasses into a tightly-sealed building. Most noticeably, it irritates eyes, skin and respiratory tract and causes allergic reactions. It also causes mutagenic and carcinogenic changes in test animals. Research is underway better to understand these chronic changes and to evaluate other potential effects. Physicians aware of indoor air pollutants can provide their patients with better medical care.

Becker, Henk A.; Porter, Alan L. (1986). Methods and Experiences in Impact Assessment. D. Reidel Publishing Company, Dordrecht The Netherlands; 1986.

Keywords: impact assessment; environments;

Becker, Robert O.; Marino, Andrew A. (1982). Electromagnetism and Life. State University of New York Press, Albany, New York, 1982.

The authors stress that the environment is now thoroughly polluted by man-made sources of electromagnetic radiation, with frequencies and magnitudes never before present. They conclude that the present abnormal electromagnetic environment can constitute a health risk. The authors provide background information which establishes that organisms can

receive information about their environment in the form of natural electromagnetic signals, and that this can lead to physiological and behavioural changes. They review the effects of man-made electromagnetic energy upon the human nervous, endocrine, cardiovascular, and hematological systems. They conclude that electromagnetic energy can produce varied and nontrivial biological effects, that such effects can be produced at energy levels well below the limit which initiates thermal effects in tissues, and that the electromagnetic field is a stressor. Studies are cited which confirm that both high- and low-frequency electromagnetic fields have been shown capable of impairing resistance to infection.

The authors conclude that while no biological function appears to be impervious to nonthermal electromagnetic fields, the nature, extent and physiological significance of the effects to be expected in different organisms remain, for the most part, to be determined by future studies. An extensive bibliography is included.

Befekadu, Ferede (1984). The Effect of Environmental Strain Upon Students' Comfort and Work Performance: Buildings, Ventilation and Thermal Climate; Berglund, Birgitta; Lindvall, Thomas; Sundell, Jan — editors. *Indoor Air*, Vol. 5, pp. 349-353; Swedish Council for Building Research, Stockholm Sweden, 1984.

Extensive investigations have been made in the past in order to provide a healthy and comfortable living and working indoor environment. This paper discusses the effect of thermal, visual and noise environments upon the health, comfort and work performance of students engaged in sedentary activity in the classroom. The investigation was carried out in two different buildings of different orientations and surroundings which offered varied environmental conditions. Suitable subjective criterias and corresponding scales expressing different subjective responses in terms of numerical values were used. Physical measurements on thermal, visual and noise environments were taken and simultaneous subjective assessment of 40 acclimatised subjects was made. Assessment results indicate a marked relationship between the physical measurements and the corresponding subjective assessment and also revealed that there were no marked interactions between the personal life of the subjects and the thermal, visual and noise environments.

Bell, Iris R. (1982). Stress, Immune System, and Illness. Presented at the 16th Advanced Seminar in Clinical Ecology & 17th Annual Meeting, Banff Canada, October 3-8 1982.

This paper discusses the human ecology concept of total load in relation to the scientific literature on stress. Psychosocial stressors, though difficult to assess, may cause significant biological changes. Interactions between different classes of stressors can exert markedly more severe adverse effects than can a single stressor in itself. The

nature of the stressor and the chronicity of exposures can affect the specificity and extent of the biological response to the stressor. The physiological mechanisms by which stress could affect health include not only immunological, but also hormonal and neurochemical systems. The clinical implications of this work for allergists include the need for attention to the psychosocial consequences of abrupt disruption and/or loss of customary lifestyle by avoidance measures and to the need for adequate treatment of pre-existing psychosocial stressors in newly-diagnosed patients.

Bell, Iris R. (1982). Clinical Ecology: A New Medical Approach to Environmental Illness. Common Knowledge Press, Box 316, Bolinas CA 94924; 1982.

Practitioners of clinical ecology maintain that a broad range of common physical and psychological disorders can be triggered in susceptible individuals by chronic and often low-level exposures to foods, environmental chemicals, and natural inhalants, with emphasis on the first two. Low doses of substances which singly might be benign may interact additively or synergistically on some common pathways in the body to produce illness, with the onset of illness depending on the total stress load, including all of the psychosocial, physical, chemical, antigenic or infective stressors that impinge on the individual.

Bellanti, Joseph A. (1974). Immunologic Responses to Chemical Pollutants. Pediatrics, Vol. 53, No. 5, Part 11, pp. 818-819; May 1974.

The immune system may be considered a series of adaptive mechanisms to an ever-changing and hostile environment. Any teratogenic effect early in embryonic life can have profound consequences on the immune system of the developing infant. A number of chemical pollutants are known to injure the immune response of the fully developed organism. The human fetus and newborn are especially susceptible to substances like lead, cadmium, nitrogen dioxide, and sulfur dioxide.

Benard, J.M.; McKenna, T.A.; McKinnon, D.L.; Faraht, J.P. (1985). Relating Health and Environmental Studies in a Tight Building Syndrome Investigation. Air Pollution Control Association, 1985.

A two-phased study was undertaken to investigate health and environmental complaints in a large, modern, hermetically sealed office complex. Health surveys of present and past employees were conducted to fully characterize the nature and magnitude of complaints. An environmental study was conducted to identify environmental factors but it was seriously limited by lack of time and money. Two ventilation parameters, fresh air supply and air distribution, were studied for ten floors which had been characterized as having low and high frequencies of the irritation syndrome. Ventilation was seen to be better with a uniform

office configuration on a floor. There was little indication that the type of symptom and type of environmental complaint were correlated in any meaningful way. Lessons were learned from this study and its limitations which can be applied to future studies of this type.

Berglund, B.; Berglund, U.; Lindvall, T.; Nicander-Bredberg, H. (1982). Olfactory and Chemical Characterization of Indoor Air; Towards a Psychophysical Model for Air Quality: from Indoor Air Pollution; Spengler, John—editor. *Environ. Int.*, Vol. 8, No. 1/6, pp. 327-332; 1982; Pergamon Press, Oxford, Toronto.

The relationship between the odour strength of total air samples and the odour strengths of the constituents was investigated in three field experiments in an office building and a new preschool. Perception of odour is an important aspect of indoor air quality because a majority of these contaminants are odorous. Characteristic chemical patterns of indoor air can be detected and described with methods of pattern recognition such as component analysis. Such patterns may be different for different buildings; similarly, perceived odour patterns may distinguish indoor air samples from outdoor or other sources. The overall odour strength of an indoor air sample can be predicted from the number of components most frequently reported to have a strong odour.

Berglund, B.; Johansson, L.; Lindvall, T. (1982). A Longitudinal Study of Air Contaminants in a Newly Built Preschool: from Indoor Air Pollution; Spengler, John—editor; pp. 111-115. *Environment International*, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

In about 100 newly built preschools in Stockholm, staff and children have reported medical symptoms that are associated with bad air quality, such as dry throat, irritation of eyes and lips, hoarseness, hacking cough and itching. The air quality in one such preschool was investigated in a longitudinal study. Typical air contaminants emanating from building materials were determined, their variation over time (0-18 months) was measured, and the influence of the ventilation system on contaminant concentrations was studied. The study shows that all the organic compounds decline in concentration mainly within the first 6 months of occupancy. A preschool building needs to be gassed off during the first six months after its construction with no recirculation of return air allowed. During at least 1 to 2 additional years, it is desired that the recirculation rate of return air is restricted, perhaps to 50%.

Berk, James V.; Hollowell, Craig D.; Pepper, James H.; Young, Rodger A. (1980). The Impact of Reduced Ventilation on Indoor Air Quality in Residential Buildings. University of California; Lawrence Berkeley Laboratory, Energy and Environment Division, Berkeley CA; March 1980.

The levels of air contaminants inside buildings are often higher than ambient outdoor levels. Interest in conserving energy has been motiva-

ting home-owners and builders to reduce infiltration rates in residential buildings and builders to reduce ventilation rates in institutional and commercial buildings. However, the resulting decrease of indoor/outdoor air exchange will tend to increase the concentration of many indoor air pollutants. Three indoor contaminants—nitrogen dioxide from gas stoves, formaldehyde from particleboard and urea-formaldehyde foam insulation, and radon from various building materials—are currently receiving considerable attention in the context of the potential health risks that are associated with reduced infiltration and ventilation rates. It is likely that some increased health risk will accompany an increase in indoor contaminant exposure; hence, it is desirable not to allow these concentrations to rise above human tolerance levels. There are several possible ways of circumventing increased health risks without compromising energy conservation considerations.

Bertell, Rosalie, Ph.D. (1984). Handbook for Estimating Health Effects From Exposure to Ionizing Radiation. Institute of Concern for Public Health, 67 Mowat Avenue, Suite 343 Toronto, Ontario M6K 3E3.

This handbook contains information which enables the translation of human radiation exposure levels into probable health effects, for example, a labor union faced with evaluating a list of worker radiation exposures, or a physician deciding on the risks and benefits of various X-ray procedures. Some radioactive materials emit rays which can penetrate the body even though the materials remain outside. The most common are gamma and X-ray emitters. Some beta particles are able to penetrate the outer skin layer and do some internal damage to humans. When radioactive particles are taken within the body through inhalation or ingestion, they can do more severe local biological damage to the cells immediately surrounding the bone, organ or tissue in which they lodge.

Berwick, Donald M.; Komaroff, Anthony L. (1982). Cost Effectiveness of Lead Screening. New England Journal of Medicine, pp 1392 to 1398; June 1982.

Lead screening programs may reduce childhood disabilities, but at what cost? Through a review of the literature, the authors performed a cost-effectiveness analysis in which the costs, savings, and health benefits of two lead screening strategies were compared with each other and with a strategy of no screening in a population of three-year-old children. When the prevalence of lead poisoning among the children screened is 7 percent or more, it is estimated that free screening averts morbidity and results in net savings: it is both better and cheaper than no screening. At all prevalence rates, free erythrocyte protoporphyrin screening is most cost effective type.

Besch, Emerson L.; Dart, David M.; Goldman, Ralph F.; Horton, Robert J.M.; Logsdon, Robert F.; McNall, Preston E. Jr.; McQuiston, Faye C.; Turk, Amos; Woods, James E. (1981). Position Paper on Legionellosis: Part I and Part II. American Society of Heating, Refrigerating and Air Conditioning Engineers July 1981 and February, 1981.

This is the report of an Ad Hoc Committee. It discusses the micro-organism which causes the disease which is found in wet soil or water such as is found in lakes, domestic water taps, sewage treatment facilities and recreational waters. It is amplified by such things as cooling towers, humidifiers, hot water systems and is thought to be transported by airborne means. Treatment of water is discussed as an amplifier control. Filtering of the make-up air for a building is practical and can provide removal efficiency approaching 100%. Within buildings, special attention should be given to warm and/or standing water such as in humidifiers, shower heads, air conditioning ductwork. Research needs are discussed such as isolating the bacteria in their natural environment, methods of practical measurements of airborne concentrations, control strategies for amplifiers, threshold concentrations for infection, effectiveness of filtration, and susceptibility of various populations.

Billings, Charles E.; Vanderslice, Sandra F. (1982). Methods for Control of Indoor Air Quality: from Indoor Air Pollution; Spengler, John—editor; pp. 497-504. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Keywords: buildings; ventilation; air filters; indoor air pollution; risk assessment;

Bjorklid, Pia (1985). Children's Outdoor Environment From the Perspective of Environmental and Developmental Psychology. Children Within Environments, Chapter 6, pp 91 - 106; Garling, Tommy and Valsiner, Jaan editors; Plenum Press; 1985.

Some of the findings are reviewed of a study (Bjorklid, 1982) which in 1969 was initiated by the Swedish Public Committee on Children's Outdoor Environment. The purpose of the study was to provide a frequency description of the environment-behaviour interactions on two housing estates of children ages 4 to 12. Boys spent much more time outdoors than girls. Some implications are drawn concerning the importance of the outdoor environment for children's social and psychological development. Examples of factors in the physical environment that restrict outdoor stay are high-rise buildings, traffic, long distances to attractive play spaces, and bad weather. An indirect social factor is a lack of suitable activities for adults. Recommendations include setting aside all open spaces on housing estates for play, providing playgrounds with play-leaders, allowing children opportunities for concrete action upon the environment, helping them to develop a sense of competence, and encouraging them actively to bring about changes in the environment.

Blume, Kathleen A. (1976). Air Pollution in the Schools. from *Clinical Ecology*, Lawrence D. Dickey ed; chapter 39; pp. 369-376; Charles C. Thomas; 1976.

There are many sources of indoor air pollution in our schools. Many children are reacting to a toxic exposure when they become irritable, excited, depressed, unreasonable or antisocial. Many children who demonstrate an inability to learn, who are slow learners; who have reduced reading comprehension and speed, reduced memory, or who show mental confusion, could be satisfactory students if there were no toxic fumes present in their school environment. The major causes of air contamination in schools are found to be aerosol sprays, janitorial supplies, scholastic supplies, building materials and furnishings, heating, ventilating and cooking equipment, cosmetics and toiletries, smoking and school buses.

Blumenthal, James A; Burg, Matthew M.; Roark, Steven F. (1986). Behavioral Approaches to Primary and Secondary Prevention of Coronary Heart Disease: from *Handbook of Prevention*; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 287 - 306.

Research has demonstrated that risk factors for the development of coronary heart disease can be modified using behavioral interventions. Hypertension, hyperlipidemia, obesity, cigarette smoking, and Type A behavior all have been shown to be potentially modifiable in this way. However, with the exception of smoking, there is inconclusive evidence that behavioral interventions will prevent it in later life or that such intervention will reduce morbidity or mortality once it has become evident. The most prudent approach is to identify children at risk while relying on a variety of educational and behavior change strategies to promote wellness behaviors. Education of adults is also important in a variety of settings. In addition to clinical trials for primary and secondary prevention, future research should include models to study mechanisms by which behavioral factors link to heart disease. Public health policy needs to be redirected toward preventing illness rather than treating it.

Blumenthal, Monica D.; Davie, James W. (1980). Dizziness and Falling in Elderly Psychiatric Outpatients. *American Journal of Psychiatry*, Vol. 137, No. 2, pp. 203-206; February 1980; American Psychiatric Assoc.

Keywords: elderly; ambulatory care; blood pressure; effects of drugs; coronary disease; dizziness; equilibrium; female; psychiatric hospitals; orthostatic hypotension; mental disorders; middle age; psychotropic drugs;

Boegman, R.J.; Bates, L.A. (1984). Neurotoxicity of Aluminum. *Can J Physiol Pharmacol*, No. 62, pp. 1010-1014; August 1984; National Research Council of Canada.

The toxic effects of aluminum, the earth's most abundant metal, on the nervous system are reviewed. Soluble Al salts can be absorbed from the stomach; nutritional deficiencies of calcium and magnesium can enhance Al absorption. Al is deposited in the gray matter of the brain and affects many neuronal processes: it stimulates protein synthesis, inhibits axonal transport, and increases the breakdown, while decreasing the re-uptake, of neurotransmitters. Behavioural consequences may include memory loss, impaired motor coordination, decreased learning ability, psychotic reactions, and seizures. Common sources of Al include antacids, deodorants, paper towels, drinking water (Al sulfate is used in water purification), airborne dust, and food. High levels of Al in soil, and acid rain, which can leach Al from the soil, increase the level of Al and tend to decrease the amount of selenium in locally grown plants.

Boer, J. de (1986). Community Response to Soil Pollution; A Model of Parallel Processes: Part D: Community Processes and Social Participation. Methods and Experiences in Impact Assessment; Part D1; pp 185 - 200; Reidel Publishing Company, Dordrecht The Netherlands; 1986.

A field study investigated the perception of environmental hazards in residential areas polluted by chemical wastes. Soil pollution is an important problem in The Netherlands. In four cases of soil pollution experiences of the inhabitants, the information process, social and medical assistance and participation by the inhabitants are studied. The data were collected in comprehensive interviews with involved professionals. Increasing understanding of individual reactions to cases of soil pollution could have important consequences for policy. The Authorities could improve their handling of cases such as this by improving the information provided to residents, improving citizen participation, and acknowledging their responsibility for future living conditions in the affected neighbourhoods.

Bokina, A.I.; Eksler, N.D.; Semenenko, A.D.; Merkur'yeva, R.V. (1976). Investigation of the Mechanism of Action of Atmospheric Pollutants on the Central Nervous System and Comparative Evaluation of Methods of Study. *Environmental Health Perspectives*, Vol. 13, pp. 37-42; National Institutes of Health; 1976.

Some aspects of the mechanism of action of atmospheric pollutants (acetone, benzene, ammonia, formaldehyde and ozone) on the central nervous system were studied by using methods of functional electroencephalography. Effects of the compounds were determined for the various structures of the cerebral cortex of experimental animals. The most sensitive structures were those which were first to associate in the reaction to

toxic agents. EEG indices were observed which were indicative of an adverse effect. During long-term action of toxic materials at low concentrations, changes were observed in the parameters of the primary and secondary responses of the visual evoked potential which were indicative of a disturbance of the cortical inhibition processes. This can be considered one effect of atmospheric pollutants at low concentrations.

Bonham, Gordon Scott; Wilson, Ronald W. (1981). Children's Health in Families with Cigarette Smokers. *American Journal of Public Health*, Vol. 71, No. 3, pp. 290-293; March 1981.

Recent studies have indicated higher rates of certain respiratory conditions among children who live in households with adults who smoke cigarettes. This paper analyzes data from the 1970 National Health Interview Survey. Children in families with no smokers had an average of 1.1 fewer restricted activity days and 0.8 fewer bed disability days per year than did children in families with two smokers. Children in families with one smoker were in between. Acute respiratory illness accounted for the difference in disability days among children in families with different smoking characteristics. Family smoking was also measured by the combined number of cigarettes smoked by adults; children in families which smoked 45 or more cigarettes a day had 1.9 more restricted activity days and 0.9 more bed disability days due to acute respiratory conditions than did children in families who did not smoke cigarettes. The age of the child, the number of adults in the family, the education of the family head, and the family income were all controlled and did not eliminate the relationship between children's health and family smoking.

Borland, Barry L.; Rudolph, Joseph P. (1975). Relative Effects of Low Socio-Economic Status, Parental Smoking and Poor Scholastic Performance on Smoking Among High-School Students. *Soc. Sci & Med*, Vol. 9, pp 27 - 30; Pergamon Press; 1975.

Previous studies have established that parental smoking, socio-economic status and scholastic performance are all related to smoking in high school students. A study of 1814 students at a high school in Western Pennsylvania confirmed the three factors previously identified as factors influencing the smoking of teenagers. Findings of this study also showed that of the three factors, scholastic performance was clearly the strongest, with parental smoking next in strength and socio-economic status a weak third. These results are discussed in relation to similar findings on antisocial behavior in teenagers.

Borrows, Peter; Turner, Richard (1984). How Safe is Science in Schools?. *New Scientist*, No. 13, pp. 12-15; September 1984.

Safety regulations for school science laboratories are going unobserved for lack of funds. But school is the ideal place to instill a lifelong awareness of potential hazards.

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

Boulding, Elise (1979). Children's Rights and the Wheel of Life. Transaction Books; 1979.

Although they make significant contributions to the world's work and well-being, often despite severe legal and economic handicaps and social victimization, people under age twenty-one and over age sixty are excluded sectors of the world population. Striking at the roots of ageism strikes at the root pathology of human relationship: the drive to dominate and to mold. The centres of power will always let the young and the old be the first to go hungry, especially if they are poor, and most of the world's poor are either very young or very old. This book examines the active, participatory roles of children in their respective societies and then looks at children as objects of legal protection and explores the extent to which this protection helps or hinders their welfare. An extensive bibliography is included.

Bourne, Patricia Gerald; Medrich, Elliot A.; Steadwell, Louis; Barr, Donald (1971). Day Care Nightmare — A Child-Centered View of Child Care: Working Paper No. 145. Institute of Urban & Regional Development, University of California; February, 1971.

Child care services need to be child-centered rather than focused on the needs of adults in the labour force. When a child's eligibility for a program is tied to his or her mother's participation in a training program or a particular job, frequent damage to the child's sense of stability and security is inevitable. When franchisers and industry, in search of cheap programs, seek to avoid expensive public agency standards, it is the children and their chances for emotional and educational development who are the losers. A set of criteria needs to be formulated based on the needs of children for extra-parental care. The existing constellation of services needs to be looked at as a system and evaluated as to whether that system is able to respond to the current and evolving needs of children.

Bowles, A.M.; Shirliffe, C.J. (1981). Development of a Canadian Standard for Urea Formaldehyde Thermal Wall Insulation: reprinted from Thermal Insulation Performance, American Society for Testing and Materials, Special Technical Publication 718, 1980, pp. 361-394. National Research Council of Canada, Division of Building Research, Ottawa Canada K1A 0R6; 1981.

Keywords: urea formaldehyde; foam; insulation; plastic; thermal insulation; material standards; formaldehyde; fungal growth; shrinkage; freeze-thaw; hydrolysis; field investigations; pH; derating; thermal resistance; NHW?

Bozzelli, Joseph W.; Russell, Joel F. (1981). Airborne Asbestos Levels in Several School Buildings Before and After Bulk Asbestos Removal. Intern. J. Environmental Studies, Vol. 20, pp. 27-30; 1982; Gordon and Breach Science Publishers, Inc.

Airborne particulate samples were collected in several public schools before and after friable asbestos containing (5-20 percent) insulation material had been removed from the ceiling areas. Transmission electron microscopy with selected area electron diffraction (SAED) was used for fiber identification and counting. Asbestos fiber concentrations in the air were in the range 5-40 ng/M³ before removal of the friable asbestos ceiling insulation. The airborne asbestos concentrations approximately one week after removal showed reductions of 56 to 90 percent.

Bradley, Eileen; Erman, Tahire; Fanos, Irene; Gottlieb, Nina; Jackson, Gerald, Mikula, Amy; Rivlin, Leanne; Rosenblum, Ilith; Sorensen, Lena; Walker, Peter; Wolfe, Maxine (1986). Workshop; Diversity as an Environmental Issue: Introduction. Environmental Design Research Association, April 1986.

The working group is composed of 11 participants — women and men, from different cultural backgrounds, including Irish, Jewish, Black American, Danish, Turkish, Israeli, Polish, Greek, Protestant, Catholic, and Moslem. Some were immigrants, others were US-born. Economics, age, family structure, sexual orientation, family structure, education, and work experience were all diverse. Their commonality was the value they placed on diversity, on exploring differences and similarities, and understanding ways in which as individuals and design researchers they could integrate that value into their work.

In order to help create environments which will sustain and support the positive or change the negative realities in our lives, a point of departure is needed that acknowledges and sustains difference and diversity, both in people and in environments, while building community based on acknowledging difference. In order to do this, it is necessary to understand the ways in which we have been taught to ignore diversity and to develop ways of unlearning this limited way of thinking and acting. There is an umbrella of oppression: within our society different groups have different amounts of power and a person could in one respect be part of a dominating group, for example, white or male, and at the same time be part of an oppressed group, for example, a white woman or a Black male.

Bradley, Eileen; Erman, Tahire; Fanos, Irene; Gottlieb, Nina; Jackson, Gerald, Mikula, Amy; Rivlin, Leanne; Rosenblum, Ilith; Sorensen, Lena; Walker, Peter; Wolfe, Maxine (1986). From Stereotyping to Revealing Diversity; The Role of Language in Environmental Research. from Diversity as an Environment Issue Workshop; Environmental Design Research Association, April 1986.

Words are frequently used to perpetuate the status quo in discrimination against women, ethnic/racial groups, people with disabilities, and so on. Making words more accurately describe the specific population under discussion will weaken overt and insidious biases by causing both writer and reader to think about people and environments in a less pre-conceived, stereotypical way. The first step in the direction of changing the status quo could have the researcher asking a person, "How would you prefer to be described?"

Bradley, Eileen; Wolfe, Maxine; (1987). Where Do the 64-Year-Old Jewish Latina Lesbians Live?: Diversity of People as an Environmental Issue. Proceedings of the EDRA 18 Conference, Ottawa, Canada, May 29-June 2, 1987, pp. 175-181, edited by Joan Harvey and Don Henning; Environmental Design Research Association, Washington, D.C.; 1987.

This paper analyzes titles and contents of published EDRA research material, based on a framework which acknowledges and supports the diversity of people. The findings show that we are ignoring diversity by homogenization, invisibility, treating people as deviant, specifying non-dominant groups and not specifying dominant groups, only focusing on a non-dominant group in comparison to a dominant group, inferring membership in a category on spurious grounds, and using descriptors as "throwaways", neither defining them nor relating them to the issues at hand, and supporting a reader's use of normative images. The authors discuss the relationship between these findings and issues of oppression, and show how ignoring the diversity of people affects environmental descriptors and explanation of phenomena.

Bradley, Jim (1987). Remarks to the Canadian Society for Clinical Ecology and Environmental Medicine. unpublished speech to the Canadian Society for Clinical Ecology and Environmental Medicine; April 3, 1987.

Environmentally sensitive people are the first to feel the effects of low-level environmental contaminants. These people should not be forgotten in pollution control decisions. Source reduction, not dilution is the solution. Alternatives to agricultural chemicals must be developed, chemical input into drinking water sources from polluting industries must be reduced, and sulphur dioxide emissions must be reduced, among other actions.

Breysse, Peter A. (1979). Formaldehyde Exposure in Mobile Homes and Conventional Homes. Presented at the 43rd Annual Education Conference of the National Environmental Health Association, June 23-28, 1979.

Keywords: formaldehyde; mobile homes; health hazards; indoor air pollution;

Breysse, Peter A. (1981). The Health Cost of 'Tight' Homes. Journal of the American Medical Assoc., Vol. 245, No. 3, pp. 267-268, Jan. 16 1981; 535 No. Dearborn St., Chicago IL 60610.

Keywords: airtight homes; health cost

Bridbord, Kenneth; Brubaker, Paul E.; Gay, Bruce, Jr.; French, Jean G. (1975). Exposure to Halogenated Hydrocarbons in the Indoor Environment. Environmental Health Perspectives, Vol. 11, pp. 215-220; June 1975.

Keywords: vinyl chloride; carcinogen; aerosol propellants; freons; ISO-butane; propane; Trichloroethane; methylene chloride; carbon tetrachloride; cardiac arrhythmia; carbon monoxide; metabolism;

Broadbent, D.E.; Broadbent, M.H.P.; Male, J.C.; Jones, M.R.L. (1985). Health of Workers Exposed to Electric Fields. *British Journal of Industrial Medicine*, Volume 42, 1985; pp. 75-84;.

The results of health questionnaire interviews with 390 electrical power transmission and distribution workers are reported, along with long term estimates of their exposure to 50 Hz electric fields, and short term measurements of actual exposure, for 287 of them. After allowing for the effects of job and location, the authors found no significant correlations of health with the subjects' measured or estimated exposure to electromagnetic fields. The general level of health was higher than found in manual workers in other industries, but there were significant differences in the health measures between different categories of job, different parts of the country, and in association with factors such as overtime, working alone, or frequently changing shift.

Brundrett, G.W. (1979). Maintenance of Spray Humidifiers. US Govt.; February 1979.

Keywords: spray humidifiers; occupational diseases; indoor air pollution; microorganisms; air conditioning equipment; culture medium; dust; NHW

Budd, Roger A.; Czerski, Przemyslaw; (1985). Modulation of Mammalian Immunity by Electromagnetic Radiation. *Journal of Microwave Power*, Volume 20, No. 1, 1985; pp. 217-231;.

This paper examines reports that electromagnetic radiation alters the function of mammalian immune systems. The authors conclude that there is no convincing evidence that electromagnetic radiation effects on the human immune system are a health hazard. Available data indicate that electromagnetic radiation exposure does not affect the ability of cells of the immune system to respond to a subsequent challenge; however, the time-course and magnitude of the response may be affected by exposure following stimulation. Research to date provided evidence that at least at some frequencies and/or amplitude and pulse modulations, the site of primary interaction of electromagnetic radiation is at the cell membrane. The authors conclude that electromagnetic radiation alters the responses of the immune system, but that much of the literature on the subject is confusing, and contradictory. They found it difficult to assess the literature in terms of its implications to human health.

Budnitz, R.J.; Berk, J.V.; Hollowell, C.D.; Nazaroff, W.W.; Nero, A.V.; Rosenfeld, A.H. (1979). Human Disease from Radon Exposures: The Impact of Energy Conservation in Residential Buildings. *Energy and Buildings*, Vol. 2, No. 3, pp. 209-215; August 1979; Elsevier Sequoia S.A., Lausanne Switzerland.

Keywords: houses; energy conservation; radon; environmental impact; air pollution; indoor air pollution; public health;

Cain, W.S.; Leaderer, B.P. (1982). Ventilation Requirements in Occupied Spaces During Smoking and Nonsmoking Occupancy: from Indoor Air Pollution; Spengler, John—editor. *Environ. Int.*, Vol. 8, No. 1/6, pp. 505-514; 1982; Pergamon Press, Oxford, Toronto.

Ventilation requirements in occupied spaces have traditionally derived largely from odor control. The requirements have rested on the notion that an environment that seems subjectively acceptable to a visitor will in fact be healthful and comfortable for both occupant and visitor. They have also derived from criterion concentrations of notable contaminants such as carbon dioxide. This investigation looked again at both sensory and physical criteria of acceptability, paying particular attention to the difference between smoking and nonsmoking occupancy in a well-controlled environmental chamber. More than 200 visitors made judgements of odor intensity and acceptability under various conditions of occupancy. The results implied that under nonsmoking conditions and moderate humidity, only about 7.5 cfm of fresh air per occupant sufficed to satisfy visitors, but that under smoking conditions at least 5 times as much fresh air is necessary.

Calabrese, Dorothy V. (1986). Fungi: Molds and Yeasts A World of New Discoveries for the Sensitive Patient. *Serendipity Environmental Medicine Newsletter*, Vol. 1, No. 4, pp 1 - 6; November, 1986.

Very little information is available in the medical literature about the health effects of molds and yeasts, compared to the wealth of knowledge that has been gained by clinicians who have been specifically treating mold sensitive individuals. This is a major problem in modern medicine: physicians look for what they have been trained to see, rather than practise the true art of medicine which is being creative and using basic knowledge as a springboard to further discoveries about patients' problems.

Sensitivities to molds can cause a variety of symptoms such as headaches, ear and hearing problems, respiratory problems like bronchitis and asthma, gastrointestinal difficulties, weight gain and loss, food cravings, as well as dermatologic, muscular, urologic, and neuropsychiatric symptoms.

Calabrese, Dorothy V. (1986). Concept of Total Load: Man and His Environment. *Serendipity Environmental Medicine Newsletter*, Vol. 1, No. 2, p 1; September, 1986.

A person's total load of exposures to toxic and allergenic substances is comprised of external load factors as well as internal individual factors. There are many factors simultaneously at work in a system that varies considerably with each individual.

Calabrese, E.J. (1978). Pollutants and High-Risk Groups: The Biological Basis of Increased Human Susceptibility to Environmental and Occupational Pollutants. John Wiley & Sons, New York, 1978.

To be at high risk with respect to a pollutant, an individual would experience the adverse health effects of the pollutant significantly before the general population because of some genetic, developmental, nutritional, physiological, behavioural, psychological, or disease state factors present which predispose the individual to the harmful effects. The author stresses that at present, the identification and especially the quantification of the numbers of individuals at high risk in the population in question is still in its rudimentary stages. The question addressed is: what percentage of the population is actually being protected from the toxic or carcinogenic activities of a pollutant, by clean air standards.

Calabrese, E.J. (1978). Methodological Approaches to Deriving Environmental and Occupational Health Standards. John Wiley & Sons, New York, 1978.

This work discusses the methodological bases for deriving pollutant exposure standards, with particular reference to the fact that there is a high degree of variability in the response of humans to different levels of air pollutants.

Calabrese, Edward J. (1979). The Influence of Ambient Ozone on the Incidence of Bone Fractures Especially Among the Elderly. Medical Hypotheses 5, pp. 201-207, 1979.

Keywords: ozone; ultraviolet radiation; vitamin D; osteomalacia; hypocalcemia; rickets;

Calabrese, E.J. (1980). Nutrition and Environmental Health: The Influence of Nutritional Status on Pollutant Toxicity and Carcinogenicity; Volume 1 - The Vitamins, and Volume 2 - Minerals and Macronutrients. John Wiley & Sons, 1980.

This two-volume work discusses the complex interaction of nutrition and pollutants on human health, itemizing research results which reveal the positive and negative effects of vitamins, minerals and other nutrients on a person's sensitivity to the adverse effects of pollutants.

Camp, Janice E.; and Morgan, Michael S. (1985). Upper Respiratory Irritation from Carbonless Copy Paper. Proceedings, 3rd International Conference on Indoor Air Quality and Climate, held in Stockholm, Sweden, August 20-24, 1984. Volume 3: Sensory and Hyperreactivity Reactions to Sick Buildings, pp. 393-396.

Clerical workers were given controlled exposures to fumes from carbonless copy paper forms approximating ambient air levels associated with normal daily usage. Nasal congestion was assessed by measuring the nasal impedance to air flow. The results provided quantitative evidence that exposure to airborne emissions of carbonless copy forms can cause acute nasal congestion.

Campbell, Dugal (1986). Research Priorities in Mental Health: Commentary. Can. J. Psychiatry, Vol. 31, November 1986; pp 746 - 749.

This paper proposes a scheme for estimating the relative merits of different branches of research within the whole field of mental health. The scheme is based upon 3 factors: the need for research in a particular topic; the possibilities of doing good science; the estimated time to a clinical payoff. It is argued that reasonable assessments of these 3 factors can be combined to give a relative weight to any particular research area. The argument is also put that one way to obtain more funds for mental health research overall is to improve the ordering of the different areas within the field of mental health.

Campbell, J.S.; Day, James; Clary, John J.; Kinloch, David; and Golberg, Leon; (1981). Final Report of the Department of National Health and Welfare Expert Advisory Committee on Urea-Formaldehyde Foam Insulation. Department of National Health and Welfare, Ottawa, Canada, April 1981.

The Committee declared that compared to other building materials, UFFI is an unstable material which deteriorates, and the rate of deterioration is dependent on the conditions to which it is exposed. Degradation results in the release of formaldehyde gas which can be carried to the living space by air leakage. Where the cavity containing the foam has not dried properly after installation, fungus growth may occur leading to possible structural and health problems.

The Committee noted the role of formaldehyde as a potential carcinogen, and irritant, and a potential allergen affecting skin and ocular and nasal mucous membranes as well as the lungs. They note that some individuals may become highly responsive to low doses leading to debilitating dermatitis, rhinitis, conjunctivitis and asthma. Superimposed on these health considerations is the likelihood of respiratory sensitization from exposure to fungal spores. The Committee states that it is not prepared to recommend any level of formaldehyde exposure that it safe. UFFI was banned under the Hazardous Products Act, Section 8b.

Campbell, Joan (1983). Ambient Stressors. Environment and Behavior, Vol 15, No. 3, pp. 355-380; May 1983.

The author concludes from previous research on environmental stressors that there is ample reason to presume that environmental stressors

interact with numerous psychological variables to produce adverse effects on health and well-being. Evidence suggests that psychological factors such as perceived control, information about the stressor, coping resources, and certain personality traits all may mediate the influence of physical environmental stressors on human well-being. The author argues that it is "heuristically useful to view global, chronic phenomena such as air pollution, community noise, and crowding as a distinct class of stressors, namely ambient stressors". Such stressors represent noxious stimulation and place demands upon us to cope. Ambient stressors over the long run are expected to affect several dimensions of human well-being, including motivation, emotions, attention, somatic health, and behaviour.

Campbell, K.L.; George, E.L.; Washington, L.S. Jr. (1980). Enhanced Susceptibility to Infection in Mice After Exposure to Dilute Exhaust from Light Duty Diesel Engines. from Health Effects of Diesel Engine Emissions, Vol. 2; pp. 772-785; Pepelko, W.E., Danner, R.M., Clarke, N.A., editors; US Environmental Protection Agency, November 1980.

A series of experiments was conducted in which groups of mice were first exposed for various durations to diluted exhaust from light duty diesel engines and then briefly to an infectious aerosol generated by nebulizing cultures of a bacterial pathogen (streptococcus). Typically, post-infection mortality was significantly greater in groups exposed to exhaust than in their corresponding control groups exposed to purified air only. Data of recent diesel and of past diesel- and catalyst-treated gasoline engine exhaust experiments suggest a somewhat greater excess mortality from enhanced susceptibility to bacterial infection in mice exposed to diesel exhaust than in those exposed to catalytic gasoline exhaust. Limited data on acute tests of NO₂ and acrolein vapor alone suggest that the infectivity-enhancing effect of diesel exhaust could be accounted for in large part by these components. Exposures to diesel exhaust, NO₂, or acrolein did not enhance the mortality to a viral pathogen.

Canada Mortgage and Housing Corporation (1983). Housing and the Elderly. Canada Mortgage and Housing Corporation; Ottawa ON; 1983.

This is an advisory document dealing with desirable standards of housing designed specifically for elderly people who are sufficiently healthy and mobile to live independently in self-contained dwelling units. Its purpose is to assist those intending to organize, finance, design, or build housing for the elderly. Designs compensate for health and mobility limitations; ensure affordability and ease of access to public facilities; allow for socialization, communication, and interaction; and provide a sense of home for residents.

Canadian Advisory Council on the Status of Women (1987). Integration and Participation: Women's Work in the Home and in the Labour Force. Canadian Advisory Council on the Status of Women; 1987.

Much of this book reviews the legal, regulatory, and practical considerations that govern women's economic status and the quality of their working lives, whether they work at home, in the paid labour force, or both. Issues examined include marriage and divorce, conditions of work, unemployment, health and safety, unionization, and pensions.

Canadian Advisory Council on the Status of Women (1987). No Vacancies! Women and Unemployment. Integration and Participation; chapter 3, pp 37 - 51; Canadian Advisory Council on the Status of Women; 1987.

Unemployment causes the woman in the household to tap emotional resources to keep the home together. Feelings of guilt, anger and despair, turned inward or outward, are a powerful part of the chronic unemployment picture. When personal and financial resources are stretched to the breaking point, women as a group are affected particularly severely, as they continue to try to provide mothering, support and household comforts in an atmosphere of tension and strain.

Canadian Association for Children and Adults with Learning Disabilities (1984). A Brief on Emission Control Standards. Canadian Association for Children and Adults With Learning Disabilities, Kildare House, 323 Chapel, suite 101, Ottawa Canada K1N 7Z2; September 1984.

This submission focuses on the effects of auto emissions on child health, specifically concerning their potential for adverse effects on brain development and function. Minimal brain dysfunction affects an estimated 10 to 15% of the population today. Lead emissions from cars are one of the major causes. Carbon monoxide is a known teratogen and another component of automobile exhaust that should be better regulated. Acid precipitation and leaching of heavy metals such as mercury, cadmium and aluminum which are known neurotoxins is another major problem which should be better monitored.

Canadian Centre for Occupational Health and Safety (1985). The Trade Names Data Base at CCOHS. Canadian Centre for Occupational Health & Safety, 250 Main St. E., Hamilton ON Canada L8N 1H6; February 1985.

This document describes the Trade Names Data Base which is a centralized source of current information on products used in Canadian workplaces. It is a computerized file of material safety data sheets supplied by manufacturers and distributors.

Canadian Home and School & Parent-Teacher Federation (1986). Preventing Child Abuse: Everybody's Responsibility. Canadian Home and School & Parent-Teacher Federation; 1986.

Child abuse is any act by an adult, or the omission of any act, which results in harm to a child. It can include physical, sexual, and emotional abuse or neglect. Physical abuse involves any kind of injury or extreme punishment of a child and also includes the failure to provide a child with the food, clothing, shelter, and health care needed for the child's optimum development. Sexual abuse is the exploitation of a child by an adult for sexual gratification and includes incest, sexual molestation, sexual assault, and the exploitation of the child for the purposes of pornography or prostitution. Emotional abuse or neglect is the constant ill-treatment of a child through the withholding of affection or through repeated humiliations. The organization has prepared a resource kit on child abuse and neglect aimed at promoting greater public awareness and understanding of the problem; encouraging individuals and community groups to become involved in positive action on the issue; and helping to prevent child abuse by showing support for families in the community.

Canadian Mental Health Association (1984). Work and Well-Being: The Changing Realities of Employment: Cross-Canada Perspectives and an Emerging Agenda on Mental Health and the Workplace. Canadian Mental Health Association, Toronto ON; September, 1984.

People who are unable to work undergo not only economic hardship, but also experience physical, psychological, and spiritual distress. In contrast, simply having a job is no guarantee of personal well-being. There are many factors in the workplace which may enhance or diminish our sense of well-being. This report is a result of action research in 5 communities across the country. The findings included, among many things, that most people have experienced negative workplace stress. Many of the work-associated pressures were found to be age-specific. Negative workplace stress affected women's personal physical and psychological health, due to the kind of work which they performed and their additional burden of family and household responsibilities. Skilled and unskilled manual labourers suffered the highest incidence of stress-related illnesses of all occupations. The issue for former psychiatric patients centres on access and opportunity.

During the next decade, an emerging priority in preventive mental health care will be to assess, anticipate, and respond to the social impacts of employment patterns. Government must establish a policy framework and program supports which would encourage positive mental health practices in both the public and private sectors. Many cooperative measures can be taken by employers and unions to respond sensitively and adaptively to the mental health needs of employees. Individual employees share a responsibility for greater sensitivity, consideration and tolerance of differentness. Strictly self-interested approaches must be abandoned in favour of concerted, collaborative action among all the parties involved. An extensive bibliography is included.

Canadian Mental Health Association (1987). Women and Mental Health in Canada: Strategies for Change. Canadian Mental Health Association, National Office, Toronto ON; April 1987.

Although the majority of those using mental health services are women, relatively little attention has been paid to their special or unique needs. The report concludes that significant improvements in women's health and well-being can be achieved only if there are general gains in the status of women in education, employment, and representation in decision-making roles. Actions directed to improving women's mental health need to be integrated and coordinated with efforts to improve women's social status and women should be active participants in the process of developing mental health programs at all levels and stages. It is recommended that funding for women's mental health research should be made a priority and that a conference be organized to identify research priorities. Health and Welfare Canada should develop appropriate materials for the promotion of information about women and mental health for use in training health professionals. Other recommendations involve promoting the awareness of sexual harassment, increased funding for rape crisis centres and shelters for battered women, the development of more comprehensive standards for occupational health and safety, assisting more women to seek training in the field of mental health planning and policy development, support of initiatives to improve women's social and economic status, and child care. A comprehensive list of references is included.

Cannon, Margaret (1987). Give Them Air: Breathe New Life into Your Workers' Productivity. Canadian Business, pp. 58-62, 95-99; April 1987.

This author cites recent investigations of 'sick building syndrome', which suggest that employees who work in sealed office buildings make more mistakes, have more office accidents, take longer to do tasks and are sick more often.

Brodsky, C.M. (1983). Allergic to Everything: A Medical Subculture. Psychosomatics, Vol 24, 1983, pp. 731-742.

Carp, Frances M. (1967). The Impact of Environment on Old People. Gerontologist; Vol. 7, No. 1, pp 106 - 108, 135; 1967.

Analysis of data on elderly applicants for new housing showed dramatic improvement deriving from a new life setting: more favorable attitudes, better mental health and social life, diminished interpersonal frictions, greater emotional independence, and fewer health complaints. The median age of the applicants was over 72. Of the 352 individuals, 204 were

successful applicants and 148 were not. Unsuccessful applicants showed little change in behavioral attitudes. Results suggest that new, improved settings foster new friendships and social activity, that old people are not necessarily rigid in behavior.

Carson, Bonnie L.; McCann, Joy L.; Ellis, Harry V. III; Herndon, Betty L.; Baker, Larry H. (1981). Methanol Health Effects. Environmental Protection Agency; December 1981.

Health effects literature primarily related to inhalation exposure to methanol was collected, evaluated, tabulated, and summarized. Approx. 160 documents were collected from computerized and manual literature searches covering the period 1901-1980. Pharmacologists and an MD epidemiologist rated the documents according to their applicability to the study and their methodology. The approximately 25 documents considered useful for deriving a range of concern for human exposure to methanol from automotive emissions were tabulated. The pages of tables detail the results of acute, repeated dose, and chronic testing of mice, rats, rabbits, dogs, monkeys and humans as well as human occupational studies. A brief summary of oral and skin absorption toxicity is included. Most of the documents evaluated are described in an annotated bibliography.

Cassel, John (1977). The Relation of the Urban Environment to Health: Towards a Conceptual Frame and a Research Strategy: A Report of the Inter-University Board of Collaborators. The Effect of the Man-Made Environment on Health and Behavior, Chapter 5, pp 129 - 142; Centre for Disease Control, Public Health Service, U.S. Dept. of Health Education and Welfare, Atlanta GA; 1977; Hinkle, Lawrence E., Loring, William C., ed.

Intervention research is proposed by which a partnership would be formed between those agencies of society charged with the task of improving the quality of the urban environment and research scientists largely drawn from academic institutions. Such a partnership would have the responsibility for introducing changes in physical aspects of housing and the residential neighborhood, and would use this entree as a means for deliberate alteration in some social factors as experimental variables. Thus, improvements in housing could be introduced with greater and lesser degrees of community participation, and with varying degrees of community control and decision making, with and without attempts to modify use, crowding, etc.

Cassidy, Michael W.A. (1970). Social Indicators: Accidents and the Home Environment: working paper. unknown; October 1970.

This paper draws together two apparently disconnected topics: first, the search for ways of evaluating the quality of physical environment, and second, the use of accident statistics as social indicators. Each of

these is covered separately, but the main focus will be on whether data on the incidence of home accidents can be used to indicate the quality of the home environment. A good theory about environmental constraints has not yet been developed. Most attempts to develop such a theory have themselves been constrained by avoidance of the most obvious limitations on interaction. Little attention has been paid to the legal, political, and economic constraints on any activities undertaken. The conflict of attitudes and expectations between people in different contextual relations with the environment — the employee compared with the manager, owner compared with renter, user compared with passer-by — clearly needs more explicit recognition and theoretical treatment.

Catalano (198?). Health, Behavior and the Community: An Ecological perspective. Pergamon Press, 198?.

In the past, public health was concerned with health problems traceable to germs. But recently, illness has been more related to man-made pollutants, safety hazards, and individual resistance. An ecological paradigm is suggested for public health. It would assume, among other things, that an individual becomes ill when he or she cannot resist or avoid biological organisms and their by products, pollutants, or safety hazards; that resistance is lowered by stress; that the type of health problems attributable to pollutants will depend on the community's economic base and natural setting; and that health problems attributable to safety hazards are predictable from a community's economic base. It would also recognize that, in relation to genetic disease, in a population equally stressed and equally exposed to the same toxin, some persons will become ill and others not because of constitutional factors inherited from their parents; as stress levels rise in a given community, those with genetic predispositions will become ill before those with average or above average tolerance.

If public health is to contend with the illnesses precipitated by stress, pollutants, and safety hazards, it must become more familiar with the economic and social processes which shape the human community. The relationship among demographic and economic characteristics of a community and the health of its population will have to be carefully measured. Such measurement will facilitate both the remedial and preventive functions. Prediction is prerequisite to effective preparation.

Chalupiak, Sharon (1987). Telecommuting Means Never Being Late For Work. *Toronto Computes*, Vol. 3, No. 6; pp 4 - 6; June, 1987.

Running a computer-based business from a home office is a work alternative that allows flexibility and control over the working environment. Women and some men are able to integrate family and work lives when they work at home. Higher efficiency, economy, reduction of stress, and the elimination of commuting time are other benefits. Telecommuting has possibilities for creating employment for the disabled.

Chenier, Nancy Miller (1982). Reproductive Hazards at Work: Men, Women and the Fertility Gamble. Canadian Advisory Council on the Status of Women, Box 1541, Stn. B., Ottawa Canada K1P 5R5; December 1982.

Keywords: indoor air pollution; occupational health and safety; reproductive hazards;

Chester, Edward H.; Martinez-Catinchi, Fernando L.; Schwartz, Howard J.; Fleming, Gerald M.; McDonald, Eugene W. (1979). Patterns of Airway Reactivity to Asthma Produced by Exposure to Toluene Di-isocyanate: supplement. CHEST, Vol. 75, No. 2, pp. 229-231; February 1979.

Keywords: airway reactivity; asthma; toluene diisocyanate;

Chipman, Clark (1981). What Does it Mean When a Patient Falls? Part I: Pinpointing the Cause: Geriatric Emergencies section. Geriatrics, Vol. 36, No. 9, pp. 83-85; September 1981.

Keywords: accidents; aged; elderly; arrhythmia; dizziness; drugs; hemorrhage; hypotension; shoes; vertebrobasilar insufficiency; vision disorders;

Chisolm, J. Julian (1974). Heavy Metal Exposures: Toxicity From Metal-Metal Interactions, and Behavioral Effects. Pediatrics, Vol. 53, No. 5, Part 11, pp. 841-842; May 1974.

The extent to which pediatric exposure to a heavy metal contributes to the body burden of that metal must be considered. There is an important interaction between metals, including the essential ones, creating greater health hazards than would exposure to the metals individually. Thus the susceptibility of the young child to heavy metals may be related to diet, or the intake of protein, calcium or iron. Added to this is his or her rapid growth, which may be considered a stress factor. Recently there have been fewer children with acute disorders due to lead absorption; rather, it seems that children with infantile autism and other disabilities after normal development up to age 3 are occurring in greater numbers. In these cases, elevated blood lead levels were present and it is quite possible that their chronic increased lead absorption contributed to the behavioral disorders. They suggest that the role of lead should be considered in a wide variety of central nervous system disorders.

Chown, G.A.; Bowen, R.P.; Shirliffe, C.J. (1981). Urea Formaldehyde Foam Insulation. National Research Council Canada, Division of Building Research, Ottawa Canada K1A 0R6; April 1981.

Urea formaldehyde foam insulation has been identified as causing medical problems for inhabitants and structural deterioration of buildings. Its

use was banned by Health and Welfare Canada and the Department of Consumer and Corporate Affairs in 1980. The likelihood of health problems occurring depends on the duration of exposure to, and the concentration of, formaldehyde. This paper was prepared to help occupants of homes insulated with urea formaldehyde foam insulation to become familiar with basic information on the characteristics of the material, the factors that affect the severity of a situation, problems that may be encountered, remedial measures that can be taken, consequences of inappropriate actions, and sources of further information and assistance.

Christensen, Kathleen E. (1986). Impacts of Computer-Mediated Home-Based Work on Women and Their Families. accepted for publication by Office Technology and People, June 1986.

This is the report of a study of professional and clerical women who work at home. It concludes that women who work at home as a way of balancing child care and paid employment typically live in traditional two-parent households, where the father is the major breadwinner. These women work part-time, primarily for 'bonus' money and the psychological benefits of doing something other than being full-time home-maker and mother. Secondly, they do not work and care for their children simultaneously. They most often work when their partners can care for the children, or when their children are at school or asleep. When a professional woman has dependable, steady work, she is apt to employ paid child care. Third, corporations that hire home-based workers as independent contractors run the risk of creating second-class corporate citizens.

Christensen, Lary; Krietsch, Kelly; White, Beth; Stagner, Brian (1985). Impact of a Dietary Change on Emotional Distress. *Journal of Abnormal Psychology*, Vol. 94, No. 4, pp 565 - 579; 1985.

A single-subject design was used to investigate the impact of a dietary change on the emotional state of four individuals selected by means of the Behavioral Index of Metabolic Imbalance and a subsequent interview. The dietary change for three subjects consisted of a high protein-low carbohydrate diet void of sucrose and caffeine, whereas only caffeine and sucrose were eliminated for the fourth subject. Results revealed that subjects reported many symptoms and/or presented a distressed profile during baseline assessment. However, following a 2-week dietary change symptoms declined, and tests revealed a more stable and less distressed individual. Overall, the results suggest that a dietary change can remediate the emotional distress exhibited by some individuals.

Church, Kathryn (1986). From Consumer to Citizen: Building a Framework for Support. Canadian Mental Health Association; May, 1986.

This reports on the proceedings of a conference on mental health advocacy designed to encourage self-help development for people with mental disabilities. The conference structure was meant to give former

and present users of mental health services a structure within which to participate in a process of joint planning for a system which would better respond to their needs.

Ciaranello, Roland; Lipton, Morris; Barchas, Jack; Barchas, Patricia R.; Bonica, John; Ferrario, Carlos; Levine, Seymour; Stein, Marvin (1983). Panel Report on Biological Substrates of Stress: Analysis and Implications of Research/A Study by the Institute of Medicine, National Academy of Sciences. Stress and Human Health; chapter 8, pp 189 - 254; Elliott, Glen R.; Eisdorfer, Carl ed; Springer Publishing Company, New York; 1983.

The biological mechanisms relating to stress are examined. Psychological and social stressors have been found to affect the immune response and disease susceptibility. The processes are complex and need further clarification. There is a great need for genetic studies to be integrated with work on environmentally caused variation in stress reactions. Different individuals can respond quite differently to the same environmental stressor, and patterns of immediate and delayed responses to stressors may differ markedly. An extensive bibliography is included.

Clark, Don (1987). The New Loving Someone Gay: Revised and Updated. Celestial Arts, PO Box 7327, Berkeley, CA USA 94707; 1987.

The author specifically addresses the task of helping someone who is non-gay overcome feelings of prejudice towards homosexual people. He notes that prejudice breeds on ignorance: in order to keep your prejudice intact you must be sheltered from too much information about the people who are the targets of your prejudice. The invisibility of gay people, in response to overt discrimination, perpetuates a situation in which non-gay people have little or no accurate information about lesbians and gay men, to counter stereotyped images of gays.

Clark, Elizabeth Johns; Rieker, Patricia Perri (1986). Gender Differences in Relationships and Stress of Medical and Law Students. Journal of Medical Education; Vol. 61, pp 32- 40; January 1986.

A small-scale comparative study of medical and law students was undertaken at a large, southern state university to examine the sources and consequences of stress during professional training. Specifically, the impact of stress on personal relationships was explored. Gender differences were found in the degree and source of stress perceived by the students. Women reported significantly more stress than men. Unlike the men, women found sexism and difficulties with partners to be particular sources of stress. Although both men and women reported that the stress of the professional training had resulted in strained personal relationships, proportionately more women than men stated that their personal relationships had ended.

Coalition for Gay Rights in Ontario (1986). Discrimination Against Lesbians and Gay Men; The Ontario Human Rights Commission: A Brief to the Members of the Ontario Legislature. The Coalition for Gay Rights in Ontario; October 1986.

The range of extent of discrimination experienced by Ontario's lesbians and gay men is described, in the hope of persuading members of the legislature to include sexual orientation in the Ontario Human Rights Code. The brief explains some of the attitudes behind anti-gay discrimination and the process whereby they might be changed. Unjustifiable prejudice against lesbians and gay men is expressed in harassment, hate propaganda and even murder. Public policy in Ontario reflects and reinforces homophobic prejudice, particularly in such areas as education, family law, police/community relations and the administration of health (a matter of increasing concern in light of the AIDS crisis). Governmental complicity in anti-gay discrimination contributes to discrimination in the private sector.

Cohen, Martin A. (1985). Air Pollution Exposures To Campers Inside of Tents; A Study of the Use Of Camping Stoves and Lanterns. Air Pollution Control Association, 1985.

The use of combustion appliances for the purpose of cooking or lighting inside camping tents has the potential of producing high indoor concentrations of combustion gases. These products may produce a substantial exposure to the individuals inside of the tent. This study investigates the build up of two of these products of combustion, NO₂ and CO while using a gas lantern inside of a family tent and a small gas backpacking stove inside of a backpacking tent.

Colley, J.R.T.; Brasser, L.J. (1980). Chronic Respiratory Diseases in Children in Relation to Air Pollution. World Health Organization, Regional Office for Europe, Copenhagen Denmark; 1980.

The World Health Organization, Regional Office for Europe has organized an international collaborative study on the relationship between air pollution and respiratory diseases in children, beginning in 1971. Studies were carried out in 8 countries: Czechoslovakia, Denmark, Greece, the Netherlands, Poland, Romania, Spain and Yugoslavia. In each country areas with relatively high and low levels of air pollution were selected and investigation of groups of school children around the age of 10 was carried out. Peak expiratory flow rate (PEFR) was measured in all children and a questionnaire on respiratory symptoms was completed with the assistance of the parents. Air pollution data were obtained from existing monitoring systems in the areas concerned. The report yielded the following results: a close association exists between air pollution and various respiratory indices in children, and smoke appears to have a greater effect on health than SO₂. It is likely that initial effects on children's respiratory indices should be sought in areas with an annual mean concentration of SO₂ and smoke of 50-200 ug/m³.

Colligan, M.J. (1981). Psychological Effects of Indoor Air Pollution. Bull. N.Y. Acad. Med., Vol. 57, No. 10, pp. 1014-1026; 1981.

Keywords: carbon monoxide; memory; pathology; anxiety; indoor air pollution;

Collishaw, Neil E.; Kirkbride, John; Wigle, Donald T. (1984). Tobacco Smoke in the Workplace: An Occupational Health Hazard. Canadian Medical Association Journal, No. 131; November 1984.

Tobacco smoke, which contains over 50 known carcinogens and many other toxic agents, is a health hazard for nonsmokers who are regularly exposed to it while at work. Involuntary exposure to tobacco annoys and irritates many healthy non-smokers. Serious acute health effects are probably limited to the one fifth of the population with pre-existing health conditions that are aggravated by exposure to tobacco smoke. The consequences of long-term exposure include decreased lung function and lung cancer. Existing air quality standards for workplaces do not directly specify an acceptable level for tobacco smoke. The evidence on the composition of tobacco smoke and on the health hazards of involuntary exposure suggests that there may not be a safe level for such exposure.

Colome, S.D.; Spengler, J.D.; McCarthy, S. (1982). Comparisons of Elements and Inorganic Compounds Inside and Outside of Residences: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 197-212; 1982; Pergamon Press, Oxford, Toronto.

The results of more than 1 year of air monitoring inside and outside of five homes in each of two communities are presented for SO₂, NO₂, mass respirable particles, SO₄, Al, Br, Cl, Mn, Na, and V. Outdoor measurements across the home site in each city are consistent with proximity to outdoor sources. Looking across indoor residential sites in each city, the home appears to alter outdoor concentrations in several ways. Indoor levels of SO₂, SO₄, Mn, and V are lower than those measured outdoors. These constituents are thought generally to result from outdoor sources. The other constituents studied are at times found in excess within homes. In some cases, the source of excess concentration of a particular constituent could be identified; often, however, the source of excess indoor concentration could not be identified.

Combs, E. Radene; Hanzal-Kashi, Amy (1985). Conversion of a Community School into Living Quarters for the Elderly. Environmental Change/Social Change; EDRA 16; pp 354; Environmental Design Research Association; 1985.

Research has shown that it is important for people to have control, security, continuity, identity and stimulation in their lives. Environmental design that enables the elderly to more adequately meet these needs can be found in a variety of housing alternatives. This paper

investigates and analyzes one such alternative — the conversion of a community school building into living units for the elderly. The community is described (land use patterns, demographic characteristics, connections with the larger metropolitan area). The decision-making process used in determining the design of the individual units and of the common space is described. The results of the initial design process and the modifications that have taken place on the basis of feedback are presented. Also presented is an analysis of current residents' evaluation of their living arrangement in terms of control, security, continuity, identity and stimulation.

Committee on Indoor Pollutants, National Academy of Science (1982). Indoor Pollutants. US Govt.; NTIS PB82-180563; March 1982.

This report is intended to characterize the quality of the indoor environment—primarily with respect to airborne pollutants, although others are discussed—and to determine the potential adverse health effects of indoor pollutants. The charge was to review, compile, and appraise the available knowledge. The Committee has also identified the research needed for abatement of indoor pollution. "Indoor" refers to the environments in homes, schools, and public buildings.

Comptroller General of the U.S. (1980). Indoor Air Pollution: An Emerging Health Problem: Report to the Congress by the Comptroller General of the United States. U.S. General Accounting Office, Document Handling and Information Services Facility, Box 6015, Gaithersburg MD 20760; Sept. 24 1980.

Keywords: toxicology; air pollution; asbestos; nitrogen dioxide; radon; carbon monoxide; formaldehyde; indoor air pollution; particulates;

Consumer Federation of America (1987). Canada Sets Exposure Limit Guidelines for Indoor Pollutants. Indoor Air News, Vol III, No. 4, p. 3; Fall 1987; Exposure Guideline Report available from Communications Directorate, Health and Welfare Canada, 5th floor, Brooke Claxton Building, Ottawa, Ontario K1A 0K9.

The articles reports on the release of the April 1987 report entitled "Exposure Guidelines for Residential Indoor Air Quality", compiled by a Federal/Provincial Advisory Committee on Environmental and Occupational Health. The report outlines maximum acceptable concentrations for specified substances, in the form of 'Acceptable Long-Term Exposure Ranges' (ALTER) and 'Acceptable Short-Term Exposure Ranges' (ASTER) for nine substances or groups of substances (aldehydes, carbon dioxide, carbon monoxide, nitrogen dioxide, ozone, particulate matter, sulphur dioxide, and water vapour). The committee found that the data base was inadequate, or human exposure limits inappropriate, for eight other contaminants. The report notes that an important consideration in deriving the acceptable exposure ranges was the possibility of interactive effects. Where possible, due account was taken of the potential for synergistic and additive effects.

Contenta, Sandro (1987). Minorities Face Racial Barriers, College Study Says. Toronto Star, page A7; December 11, 1987.

Community colleges discriminate against ethnic minority groups by severely restricting access to the training and education they need. This study for George Brown College in Toronto indicated that there were problems with admission requirements, lack of bridging programs, failure to acknowledge skills obtained in other countries, too few courses combining learning English with studying a trade, little attempt to reach out to minority groups, lack of child-care facilities, and insensitivity of college staff and course content to cultural diversity. Immigrant groups interviewed for the study made it clear that they face similar discrimination in all community colleges.

Cooper, Cary L.; Marshall, Judi (1976). Occupational Sources of Stress; a Review of the Literature Relating to Coronary Heart Disease and Mental Ill Health. J. occup. Psychol., No. 49, pp 11 - 28; 1976.

A great deal of research has been done linking the working conditions of a particular job and its relationship to physical and mental health. Poor mental health is directly related to unpleasant work conditions, the necessity to work fast and to expend a lot of physical effort, and to excessive and inconvenient hours. Poor physical health is linked to repetitive and dehumanizing environments such as paced assembly lines. Research in this area is desperately needed. Qualitative (too difficult) and quantitative (too much) work overload also negatively effects health, being strongly linked to cigarette smoking, coronary heart attacks, low self esteem, skin resistance, and high cholesterol levels.

Another major source of occupational stress is associated with a person's role at work. Role ambiguity, which exists when an individual has inadequate information about his/her work role, creates low job satisfaction, high tension, greater futility, and low self confidence and is related to increased blood pressure and pulse rate. Role conflict exists when an individual is torn by conflicting job demands or doing things he/she really does not want to do or does not think are part of the job specification. This causes physiological strain and job dissatisfaction.

Other major sources of stress at work involve the nature of relationships with one's boss, subordinates and colleagues, and the organizational structure and climate. The latter includes little or no participation in the decision-making process, lack of effective consultation, restrictions on behaviour, and office politics. Little work has been done into some of the more important extra-organizational factors, particularly the relationship between family and work. Stresses may arise over the allocation of time and commitment to the areas of occupational/productive and domestic/consumer activities.

investigates and analyzes one such alternative — the conversion of a community school building into living units for the elderly. The community is described (land use patterns, demographic characteristics, connections with the larger metropolitan area). The decision-making process used in determining the design of the individual units and of the common space is described. The results of the initial design process and the modifications that have taken place on the basis of feedback are presented. Also presented is an analysis of current residents' evaluation of their living arrangement in terms of control, security, continuity, identity and stimulation.

Committee on Indoor Pollutants, National Academy of Science (1982). Indoor Pollutants. US Govt.; NTIS PB82-180563; March 1982.

This report is intended to characterize the quality of the indoor environment—primarily with respect to airborne pollutants, although others are discussed—and to determine the potential adverse health effects of indoor pollutants. The charge was to review, compile, and appraise the available knowledge. The Committee has also identified the research needed for abatement of indoor pollution. "Indoor" refers to the environments in homes, schools, and public buildings.

Comptroller General of the U.S. (1980). Indoor Air Pollution: An Emerging Health Problem: Report to the Congress by the Comptroller General of the United States. U.S. General Accounting Office, Document Handling and Information Services Facility, Box 6015, Gaithersburg MD 20760; Sept. 24 1980.

Keywords: toxicology; air pollution; asbestos; nitrogen dioxide; radon; carbon monoxide; formaldehyde; indoor air pollution; particulates;

Consumer Federation of America (1987). Canada Sets Exposure Limit Guidelines for Indoor Pollutants. Indoor Air News, Vol III, No. 4, p. 3; Fall 1987; Exposure Guideline Report available from Communications Directorate, Health and Welfare Canada, 5th floor, Brooke Claxton Building, Ottawa, Ontario K1A 0K9.

The articles reports on the release of the April 1987 report entitled "Exposure Guidelines for Residential Indoor Air Quality", compiled by a Federal/Provincial Advisory Committee on Environmental and Occupational Health. The report outlines maximum acceptable concentrations for specified substances, in the form of 'Acceptable Long-Term Exposure Ranges' (ALTER) and 'Acceptable Short-Term Exposure Ranges' (ASTER) for nine substances or groups of substances (aldehydes, carbon dioxide, carbon monoxide, nitrogen dioxide, ozone, particulate matter, sulphur dioxide, and water vapour). The committee found that the data base was inadequate, or human exposure limits inappropriate, for eight other contaminants. The report notes that an important consideration in deriving the acceptable exposure ranges was the possibility of interactive effects. Where possible, due account was taken of the potential for synergistic and additive effects.

Cooper, W. Clark (1973). Indicators of Susceptibility to Industrial Chemicals. *Journal of Occupational Medicine*, Vol. 15, No. 4, pp. 355-359; April 1973.

Individuals differ in their susceptibility to toxic agents and other environmental stresses. All individuals do not react similarly, nor do lower animals, when exposed to toxic agents. Some individuals show effects at concentrations which do not affect the majority, while at the other end of the dose-response curve there are individuals who are unusually resistant. Terms like idiosyncrasy or hypersusceptibility have been used to describe, but not explain, extreme deviations in the direction of lack of resistance.

Couchman, Robert (1986). Provider and Parent: Fatherhood Renewal as a Dynamic in the World of Work: A New Work Agenda for Canada. Canadian Mental Health Association, Toronto Canada; 1986.

The steadily increasing involvement of fathers with their children may well be the single most important positive development emerging from the familial turbulence of the past 20 years. However, the struggle to correct the imbalance in work loads between husbands and wives continues. In two recent studies, for example, it was discovered that women working full time experience a major imbalance in family and household work loads. The fatherhood revival is having a powerful influence upon the world of work. A significant number of male workers are beginning to exercise influence so as to restore the balance in their family responsibilities and parenting relationships. This results in work time flexibility, part-time work, and job sharing.

Crawford, T. (1987). There is Something About Welfare Itself That Creates Victims. *Toronto Star*, November 30, 1987.

The article reviews a study reported in the October 1987 issue of the *Canadian Journal of Psychiatry*, in which Dr. Dan Offord of McMaster University conducted a survey of child victims of welfare. The study is reported to conclude that children from families living on welfare are three times more likely than their classmates to end up flunking a grade or sent to remedial classes for slow learners. Ontario's Minister of Community and Social Services, John Sweeney, was quoted as saying in response to a question about the study, "There is something about welfare itself, the atmosphere and the environment, that creates (these problems)". The Minister is quoted as suggesting that the solution to the welfare trap must lie in programs which lead to self-sufficiency, "so they can escape the feeling that 'Someone else is running my life'".

Statistics Canada had announced a week earlier that 1 million Canadian children live in families with incomes below the poverty line. Penny Moss, former chair of the Toronto Board of Education, was also quoted

as saying at a conference on poverty and education the week before, that schools are failing the poor, citing the example that only 46 per cent of the children of the poor end up in advanced high school programs needed for university entrance, compared with 88 per cent of middle-class children.

Crawshaw, G.H. (1978). The Role of Wool Carpets in Controlling Indoor Air Pollution. Textile Institute and Industry, Vol. 16, No. 1, pp. 12-15; January 1978.

Keywords: carpet manufacture; dust abatement; air pollution; sulfur dioxide;

Cullen, Joseph; Greenwald, Peter (1986). Prevention of Cancer: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 307 - 341.

Current scientific knowledge about cancer has led us to a new avenue toward disease prevention and health promotion. The way to health and a life without cancer requires changes in traditional lifestyles. Lifestyle and environmental factors cause or promote at least 70% of cancer cases. Most of these are controllable at the personal level. These include tobacco, diet, occupation, alcohol, viruses, excess sunshine, medicine and medical procedures, food additives, and environmental pollution. Clarifying this knowledge for the public and medical community and motivating them to make changes in their lifestyles and professional practices are the next advances required to make cancer prevention a probability.

Culot, Michel V.J.; Olson, Hilding G.; Schiager, Keith J. (1978). Field Applications of a Radon Barrier to Reduce Indoor Airborne Radon Progeny. Health Physics, Vol. 34, pp. 501-503; May 1978; Pergamon Press Ltd.

Keywords: radon; indoor air pollution; abatement;

Current Intelligence System staff/Div. of Criteria Documentation and Standard Development: NIOSH (1981). Formaldehyde: Evidence of Carcinogenicity. US Dept. of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health; April 15 1981.

Keywords: industrial medicine; formaldehyde; carcinogens; aldehydes;

D'Arcy, Carl; Siddique, C.M. (1985). Unemployment and Health: An Analysis of "Canada Health Survey" Data. International Journal of Health Services, Volume 15, Number 4, 1985; pp 609 - 635; Baywood Publishing Co., Inc.

This paper provides a cross-sectional analysis of the physical and emotional well-being of employed and unemployed workers. The data used consists of a sub-sample drawn from the Canada Health Survey's national probability sample. The analysis indicates substantial health differences between employed and unemployed individuals. The unemployed showed significantly higher levels of distress, greater short-term and long-term disability, reported a large number of health problems, had been patients more often, and used proportionately more health services. Consistent with these measures, derived from self-reported data, physician-diagnosed measures also indicate a greater vulnerability of unemployed individuals to serious physical ailments such as heart trouble, pain in heart and chest, high blood pressure, spells of faint-dizziness, bone-joint problems and hypertension. While these health differences persisted across socio-economic and demographic conditions, females and older unemployed individuals reported more health problems and physician visits whereas people under 40 reported more psychological distress. The blue-collar unemployed were found to be considerably more vulnerable to physical illness whereas the unemployed with professional background reported more psychological distress. The low-income unemployed who were also the principal family earners were the most psychologically distressed. The need for social policies that effectively reduce unemployment and its detrimental impact is clear.

Dadd, Debra Lynn; Levin, Alan S. (1982). A Consumer Guide for the Chemically Sensitive. Nontoxic Lifestyles Inc., 450 Sutter, suite 1138, San Francisco CA 94108; 1982.

Many daily processes can cause health problems for the chemically sensitive. Commercial inks and papers, certain types of reproduction processes, as well as various types of school and art supplies can give off immunotoxic fumes and cause reactions in susceptible individuals.

Dadd, Debra Lynn (1984). NonToxic and Natural. Jeremy P. Tarcher Inc.; 1984.

Most consumers assume that if products are on the market they must be safe, but thousands of products in daily use contain chemical substances that we are now discovering may cause health problems from minor discomfort to major illness. Those who seek a healthier lifestyle can significantly reduce their exposure to these toxins by paying attention to what they buy. This book lists over 1000 products in over 300 categories which are non-toxic and naturally-based, without chemical additives. It reflects the efforts of a researcher who specializes in finding consumer products free of toxic chemicals.

Daglish, Susan (1987). How Many Suffer From Food Additive Reactions?. Allergy Quarterly, Volume 23, p 5; Summer, 1987.

Questionnaires indicate that 15.5% of the population of Great Britain believe that they are adversely affected by food additives. Dr. Maurice Lessof, a professor at Guy's Hospital in England and a world-renowned researcher into food sensitivities, is conducting research to examine the actual incidence of reactions to food additives. He has concluded that the best way to handle a suspected reaction to a food additive is to eliminate it from the diet for a few months.

Daglish, Susan (1987). How To Reduce or Eliminate Indoor Air Pollution: Part IV of Environmental Concerns. Allergy Quarterly, Volume 23, No. 4, pp. 17-19; 1987.

This article by the Allergy Information Association (25 Poynter Dr., Weston, Ont M9R1K8) reviews ways of reducing indoor air pollution for the allergic person, including dust control, mould avoidance, and advice on animals, tobacco smoke, heating, pesticide use, and general chemical odours in the home. It emphasizes that many indoor substances, both natural and man-made, can adversely affect the health of susceptible individuals.

Dahl, Alan R.; Hadley, William M. (1983). Formaldehyde Production Promoted by Rat Nasal Cytochrome P-450-Dependent Monooxygenases with Nasal Decongestants, Essences, Solvents, Air Pollutants, Nicotine, and Cocaine as Substrates. Toxicology and Applied Pharmacology, No. 67, pp. 200-205; 1983.

The results of this study indicate that a variety of materials which often come in contact with the nasal mucosa can be metabolized to formaldehyde by nasal enzymes. The released formaldehyde may influence the irritancy of inhaled compounds and has been suggested to play a role in the tumorigenicity of some compounds. Thirty-two potential substrates for cytochrome P-450-dependent monooxygenases were screened with rat nasal and, for comparison, liver microsomes. Tested substrates included 6 nasal decongestants, cocaine, nicotine, 9 essences, 3 potential air pollutants, and 12 solvents.

Five substrates, namely, the solvents HMPA and dimethylaniline, cocaine, and the essences dimethyl anthranilate and p-methoxyacetophone, were metabolized to produce formaldehyde at rates exceeding 1000 pmol/mg microsomal protein/min. by nasal microsomes. Eight substrates, including four nasal decongestants, nicotine, and an extract of diesel exhaust particles, were metabolized to produce formaldehyde at rates of 200 to 1000 pmol/mg microsomal protein/min. Five other substrates were metabolized to formaldehyde at detectable rates.

Dainty, E.D.; Mitchell, E.W.; Schnakenberg, G.H., Jr. (1986). Diesel Emissions Reduction. Occupational Health in Ontario, Vol. 7, No. 4, pp 170 - 191; Fall, 1986; Ontario Ministry of Labour, Occupational Health and Safety Division, Toronto ON.

The Collaborative Diesel Research Advisory Panel was established to resolve a number of issues including: a criterion to evaluate the comprehensive toxicity of the major components of diesel exhaust, research and development to produce add-on exhaust hardware, the study of other techniques to reduce emissions from diesel engines, measurement of the impact of such devices on the underground environment and the strategy to improve mine environments, reduce ventilation costs, increase productivity and improve safety underground, depending on the circumstances of each case.

Damstra, Terri (1978). Environmental Chemicals and Nervous System Dysfunction. The Yale Journal of Biology and Medicine, No. 51, pp. 457-468; 1978.

Subtle, subclinical effects may result from chronic low-level exposures to many chemicals. The nervous system may be particularly vulnerable to many of these exogenous chemicals. Selective damage to particular areas of the nervous system has been noted with numerous toxins, and certain groups, such as the young and the elderly, may be especially vulnerable. Selected examples of associations between nervous system diseases and exposures to occupational and environmental chemicals such as inorganic and organo-lead compounds, elemental, inorganic and organo-mercury compounds, kepone, organo-phosphates, n-hexane and methyl-n-butyl ketone, have been reviewed.

Daniell, Harry (1976). Try Not to Exhaust Yourself. Runner's World, Vol. 11, No. 9, pp. 54-55; September 1976.

Carbon monoxide from automobile exhaust, paint and varnish removers, certain prescription medications such as Phenobarbitol and Dilantin, and cigarette smoke can negatively effect the distance runner in a number of ways. Commonly encountered carbon monoxide concentrations interfere with endurance performance for many hours after exposure to the gas has ended. Breathing air containing 50 ppm of carbon monoxide at rest causes the concentration in the bloodstream to rise to 5% in five hours. But during vigorous exercise, this level of air pollution can produce a level of 5% in less than one hour. This limits oxygen transport to muscle tissue by 5%. It is not metabolized but must be dissipated by being exhaled.

Darlington, L.G.; Ramsey, N.W.; Mansfield, J.R. (1986). Placebo-Controlled, Blind Study of Dietary Manipulation Therapy in Rheumatoid Arthritis. The Lancet; pp 236 - 238; February, 1986.

In a blind, placebo-controlled study of dietary manipulation therapy in outpatients with rheumatoid arthritis there was significant objective improvement during periods of dietary therapy compared with periods of placebo treatment, particularly among "good responders". Possible explanations for improvement include reduced food intolerance, reduced gastrointestinal permeability, and benefit from weight loss and from altered intake of substrates for prostaglandin production. A proportion of the improvement was due to a placebo response, but this was not sufficient to explain the whole improvement.

Davey, Paula G.; Shearer, Ruth W. (1986). Hypersensitivity Causes by Environmental Chemicals, in Particular Pesticides. Clinical Ecology, Vol. IV, NO. 1, Spring 1986.

There is an inadequate data base concerning many petrochemical compounds and pesticides. Chemically sensitized patients often share common symptoms which include the inability to concentrate, irritability, anxiety, recurrent flue-like symptoms, shortness of breath, gaseous distension, nausea, aching, weakness, and fatigue. The health test data is of very poor quality as studies in animals are not designed to look for this chemical sensitivity problem.

Cytotoxic tests of 100 patients' blood against two pesticides revealed cell damage of 50% of patients at the 1:10 dilution of Ficam and all patients tested had cell damage at the 1:80 dilution of Dursban. Case histories are cited.

Davidson, Marilyn J.; Cooper, Cary L.; Chamberlain, Deborah (1980). Type-A Coronary-Prone Behavior and Stress in Senior Female Managers and Administrators. Journal of Occupational Medicine, Vol. 22, No. 12, pp. 801-805; December 1980.

Davies, Freda Lynn (1988). Skipping School in Earnest: (advanced pre-publication draft). Afore-the-Wind Publishers, Hwy 595, NR608, South Gillies, Ont P0T 2V0.

The author examines the beliefs underlying educational institutions, and discusses the alternative of home-schooling in Canada. She demonstrates that conventional educational thinking is incompatible with the development of a future-oriented vision of promise. She notes that children are one of a number of groups who have suffered injustices as a result of ingrained attitudes which regard such groups as somehow inferior. Like racial and cultural minorities, like women, like those missing one or several of the "normal" human traits, children have been victims of a stereotyped characterization based on unexamined premises. Some of these other groups have begun to obtain release from their castes, and to gain recognition as individual humans. Children, as yet, are nowhere near achieving such liberation. Many older people still feel that it is acceptable, perhaps even necessary, for children to be treated in ways unacceptable for the treatment of adults: to be humiliated, insulted, judged constantly, interrogated and compelled to obey.

Davis, Karen (1985). Biochemical Individuality, Nutrition, and Learning Disabilities: Supported by a grant from the ACLD Foundation. University of Tennessee, Centre for the Health Sciences, Memphis, TN 38163.

Learning disabilities, a heterogenous group of entities of multiple causes are, in the most basic sense, an expression of underlying central nervous system dysfunction. This shows up as specific deficits in perceptual, integrative, or expressive processes which severely impair learning efficiency. A myriad of causes has been proposed such as genetic constitution, prematurity, complications of pregnancy, labour or delivery, metabolic disorders, malnutrition, infectious disease sequelae, trauma and chronic illness. This paper concentrates on nutrition as a factor which plays a relevant role in the development of, and perhaps in the prevention of learning disabilities.

Day, David M.; Page, Steward (1986). Portrayal of Mental Illness in Canadian Newspapers. Can. J. Psychiatry, Vol. 31, December 1986; pp 813 - 817.

Results are reported from a content analysis of 103 newspaper reports taken from eight major Canadian newspapers and selected at random from the Canadian Newspaper Index. The portrayal of mental illness and mentally ill persons in these reports was compared with that in samples of articles taken from two comparison mental health publications not receiving popular circulation. As compared with these latter publications the content analysis indicated that the newspapers portrayed mental illness and the mentally ill in a manner which could be described as essentially pejorative, thus seeming to support frequent observation and complaints from the mental health establishment about inadequate or unfair coverage of mental illness in the popular print media. Implications for the attitudes and beliefs of the general public about mental illness are offered, although the correlation between media information and the attitudes of the public remains speculative. It is suggested that further content analyses of print media be undertaken.

Day, James (1986). UFFI-Fungal Interaction. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Molds can cause a variety of ailments in humans, both allergic responses and infections. Work in the area to pinpoint the specific causative agents is hampered by measurement problems, but in some instances a positive correlation has been established. There is an increasing effort to correlate symptoms with the indoor as well as the outdoor environment. Further research should include complete mold testing of subjects from different populations with seasonal and non-seasonal asthmatic symptoms from UFFI and non UFFI containing households; making up of new antigens from available preparations for in vitro use; identification of all molds in households with subjects complaining of multiple symptoms for

which there is no medical explanation; determination of biological effects of toxins from molds obtained from UFFI containing households; and controlled exposure over time in a specially designed chamber to appropriate mold antigens, and the measurement of end organ responses. An extensive bibliography is included.

Dean, Alfred; Lin, Nan (1977). The Stress-Buffering Role of Social support: Problems and Prospects for Systematic Investigation. *The Journal of Nervous and Mental Disease*; Vol 165, No. 6, pp 403 - 417.

Over the past 20 years, a sizable body of literature has developed which serves to establish that stressful life events are associated with the onset, incidence, and prevalence of a wide range of physical and psychiatric disorders. Several prominent researchers have emphasized the importance of studying the role of social support systems as possible buffers or mediators of stress. This paper contributes to the advancement of such studies by clearly identifying key empirical, theoretical and methodological problems and suggesting some approaches to their resolution. Detailed proposals are made for approaching problems of measurement and research design. An extensive bibliography is included.

Dean, Jack H.; Ward, Edward C.; Murray, Michael J.; Lauer, Lloyd D.; House, Robert V. (1983). Immunotoxicity of Polycyclic Aromatic Hydrocarbons. NIH Immunotoxicology Workshop; pp 259 - 273; October, 1983.

Polycyclic aromatic hydrocarbons have been shown to be immunosuppressive specifically with reference to the antibody response to the T-dependent antigen SRBC. This suppression has been shown to occur within 2 days after exposure to carcinogenic PAH. Several indicators of general immunotoxicity were assessed in mice exposed to selected doses of DMBA. While pathological examination of the immune system following exposure to chemicals may give general indications of immunotoxicity, assessment of functional responses of lymphoid cells are usually more sensitive indicators of immune alteration. In the studies summarized here, mice exposed to the carcinogenic PAH CMBA, at exposure levels which produced no mortality or other obvious clinical symptoms of general toxicity, were found to have profoundly suppressed IgM antibody response to the thymus-dependent antigen SRBC. This occurrence is explained and discussed in detail.

Dershewitz, Robert A.; Williamson, John W. (1977). Prevention of Childhood Household Injuries: A Controlled Clinical Trial. *Am J Public Health*, Vol. 67, No. 2, pp 1148 - 1153; 1977.

Injuries claim the lives of more children each year than the next six leading pediatric disorders combined, and produce injuries that require medical attention for one in three children. In the preschool age group, 91 percent of these accidents and over one-half the resultant fatalities

occur in the home. This paper reports the results of a controlled clinical trial conducted to evaluate the implementation of a health education program intended to reduce the risk of childhood household injuries. The study population was randomly assigned into two demographically comparable groups. Only the experimental group mothers received an educational intervention consisting of a tutorial, home safety-proofing assignments, and follow-up. The homes of the two groups were later assessed for hazards during an unannounced visit by an interviewer who did not know to which group each home belonged. A home safety score mean for the two groups was almost identical. The program stimulated heightened interest and stated intent to improve, but did not result in actual reduction of household hazards. Active health education, as used and evaluated in this study, appears to have limited effectiveness when applied to home safety.

Diamond, Philip (1970). Health Hazard Potential from Pre-Polymerized Foam Packaging Materials: Final Report. US Govt.; January 1970.

Keywords: plastics; packing materials; isocyanic acid; industrial medicine; polymerization; indoor air pollution; confined environments;

Dickey, Lawrence D. (1976). Clinical Ecology. Charles C. Thomas, Springfield IL; 1976.

This author is concerned with adverse reactions to environmental insults modified by individual susceptibility in terms of specific adaptation. The outcome of the interreaction of excitants versus host is modified by the host's individual susceptibility which is governed by genetic background, present state of physical and mental well-being, and the individual's ability to adapt. Chemical contaminants, pesticides, herbicides, additives, vehicles, and excipients that are to be found in the air, water, food, drugs and our habitat are dealt with in this publication, with specific reference to diagnostic and therapeutic procedures, avoidance procedures, and detailed explanations of the susceptibility mechanism.

Didriksen, Nancy; Goven, Arthur; Butler, Joel R. (1986). Psychological Stress: Effects on Phagocytic Immune Functioning. Clinical Ecology, Vol. IV, No. 1, Spring 1986.

The purpose of this study was to determine whether psychological stress, specifically examination stress, would decrease immune system functioning. Subjects were 25 first-year master's and doctoral students. They were psychologically and immunologically assessed during two high- and two low-stress periods. Immunological assessments includes a white blood cell differential count and nitroblue tetrazolium test (NBT) to measure neutrophil functioning. Psychological instruments administered at each assessment period included Clinical Analysis Questionnaire, State-Trait

Anxiety Inventory, and a Brief Stress Questionnaire. Stepwise discriminant function analysis of data revealed five variables which contributed significantly to change under stress with the NBT test being the most significant. These variables yielded an average canonical correlation of .79 (p less than .002) providing evidence of support for the hypothesis that increased psychological stress will alter immune functioning and heighten psychological responses.

Didriksen, Nancy A. (1987). Psychological Stress: Effect on Humoral Immune Functioning As Measured by Immunoglobulin Levels: Abstracts. Symposium on Man and His Environment in Health and Disease; 1987.

23 volunteers were subjected to a series of psychological tests to measure stress, personality factors, emotional states and anxiety levels. Results showed increased immunoglobulin levels during periods of stress; anxiety related to external events; increase in anxiety under stress; and anxiety inversely correlated with emotional stability and coping skills while positively related to tension, increased number of somatic complaints, and obsessive-compulsive trends. It was concluded that the concept of body/mind interaction was the most realistic approach to understanding the total response patterns.

Dillon, Carolyn (1985). Families, Transitions, and Health: Another Look. Social work in Health Care; Vol. 10, No. 4, pp 35 - 44; Summer 1985.

The fate of humane, family-centered health care may well rest on the ability of caregivers to identify and empathically respond to complex family-like forces arising when large numbers of desperate people converge to interact around issues of health and illness, life and death, competence and vulnerability. The health social worker is in a position to observe how the illness family and the caregiving professionals meet in moments of crisis and stress, sometimes responding to and assessing each other in regressive ways. The social worker can also play a major facilitating role in negotiating between these two groups, building consensus for positive collaboration.

Dimmick, R.L.; Wolochow, H. (1980). Effects of Energy Conservation Measures on Air Hygiene in Public Buildings: Final Report. University of California; Lawrence Berkeley Laboratory, Energy and Environment Division, Berkeley CA; February 1980.

Keywords: hospitals; schools; aerosols; bacteria; indoor air pollution; public buildings; ventilation systems;

Dockery, Douglas W.; Spengler, John D. (1981). Indoor-Outdoor Relationships of Respirable Sulfates and Particles. Atmospheric Environment, Vol. 15, No. 3, pp. 335-344; 1981; Pergamon Press Ltd.

Keywords: cigarette smoking; gas stove; air conditioning; sulphates;

Dodge, Russell (1982). The Effects of Indoor Pollution on Arizona Children. Archives of Environmental Health, Vol. 37, No. 3, p. 151; May/June 1982.

The respiratory health of a large group of Arizona school children who have been exposed to indoor pollutants—tobacco smoke and home cooking fumes—is reported. A significant relationship was found between parental smoking and symptoms of cough, wheeze, and sputum production. Also, children in homes where gas cooking fuel was used had higher rates of cough than children in homes where electricity was used. No differences in pulmonary function or yearly lung growth rates occurred among subjects grouped by exposure to tobacco smoke or cooking fuel. Thus, parental smoking and home cooking fuel affected cross-sectional respiratory symptom rates in a large group of Arizona school children. Study of pulmonary function, however, revealed no lung function or lung growth effects during 4 years of study.

Dohrenwend, Bruce; Pearlin, Leonard; Clayton, Paula; Hamburg, Betty; Riley, Matilda; Rose, Robert M. (1983). Report on Stress and Life Events: Analysis and Implications of Research/A Study by the Institute of Medicine, National Academy of Sciences. Stress and Human Health; chapter 4, pp 55 - 80; Elliott, Glen R.; Eisdorfer, Carl ed; Springer Publishing Company, New York; 1983.

Substantial changes in future studies of life events and health are needed. Linkages between life events and health are influenced by many mediators: social characteristics, position along the life course, personality traits and coping repertoires, and the supportiveness of social networks. To provide a clearer view of the many elements that affect the stress process, studies should be as inclusive as possible. Thus a study of associations between life events and disease consequences could be improved if it also considered the effects of social supports and potentially relevant personality characteristics. A comprehensive bibliography is included.

Douglas, Carolyn J.; Kalman, Concetta M.; Kalman, Thomas P. (1985). Homophobia Among Physicians and Nurses: An Empirical Study. Hospital and Community Psychiatry, Vol. 36, No. 12, pp. 1309-1311; December 1985.

The authors note that the current epidemic of acquired immune deficiency syndrome (AIDS) has had a great impact on public attitudes toward homosexuals, leading to many instances of overt discrimination, social ostracism, and even the deprivation of various rights such as housing, employment, transportation, and funeral services. The study attempted to determine the degree of homophobia among physicians and nurses working in a large urban teaching hospital where many male homosexuals with AIDS have been treated. The investigators defined homophobia as the constellation of affective responses, including fear, disgust, anger, discomfort, and aversion, that many individuals may experience in contact with homosexuals.

Mean scores for both physicians and nurses fell in the low-grade homophobic range. Greater homophobia was found among female respondents, compared to males. Having a close relative or friend who is gay significantly reduced personal anxiety about homosexuals. The results indicated that a disturbingly high percentage (31%) of the health professionals studied acknowledge more negative, even overtly hostile feelings toward homosexuals than they had before the emergence of the AIDS epidemic. The authors speculate that health professionals may be more aware than the general population that most homosexuals do not fit stereotypes of homosexuality. While increased contact with homosexuals may serve to dispel false stereotypes and lessen homophobia in some individuals, they suggest that it may heighten anxiety and elicit greater hostility in others, if those stereotypes served important defensive functions.

Drader, Darla (1982). The Design of Geriatric Assessment Units: Psychosocial Considerations: Final Report. Health and Welfare Canada; September, 1982.

This report reviews literature concerning the design-related needs of the elderly in acute care settings. Design for the elderly should be based on an understanding of their special needs because they are less able to adapt to environments and have more difficulty coping with stress. Environments should enhance their adaptive capacity by providing a range of behavioural alternatives in a flexibly designed setting. Since place and possessions have a special significance for the elderly, design should also allow for a sense of continuity with the past as well as for privacy. Physical changes that accompany aging, such as hearing losses, visual changes, reductions in motor skill, changes in memory and learning, and diseases, should also be considered in design. An extensive reference list is included.

Dravnieks, A.; Whitfield, J. (1971). Gas Chromatographic Study of Air Quality in Schools. ASHRAE Trans, Vol. 77, Pt. 1, pp 113-123; 1971.

By use of high-capacity organic vapor collection devices and appropriate sample transfer analysis techniques, combined with psychophysical (sensory) odor evaluation of the gas chromatographically resolved components at the effluent port of a gas chromatograph, it is possible to gather considerable insight into the inventory of the odorous components of an air sample. These procedures were applied to a study of air in several Chicago metropolitan area schools. The outside air was also examined, to permit assessments of the contribution of outside air pollution to the air pollution of the school air by organic vapors and odorants. The gas chromatographic parameters were then inspected, using non-parametric statistics for correlations to the air quality which was first independently ranked using engineering data and on-the-spot observations.

Dudney, C.S.; Walsh, P.J. (1981). Report of Ad Hoc Task Force on Indoor Air Pollution. US Govt.; NTIS ORNL/TM-7679; April 1981.

This Ad Hoc Task Force on Indoor Pollution has reviewed information on indoor air pollution and the potential impact on human health of energy conservation measures. The Task Force concluded that: the indoor environment is likely to include exposure to radon daughter nuclides, formaldehyde, carbon monoxide, nitrogen dioxide, respirable particulates and asbestos, as well as other undefined pollutants; indoor exposure may constitute 80-95% of the total exposure for some pollutants; some energy conservation measures reduce the rate of exchange between indoor and outdoor air; reduced air exchange can lead to increased indoor levels of some pollutants; increased indoor pollutant levels lead to increased exposure and increased risk of health impact.

Duhl, Leonard J. (1969). Health — 2000 A.D. American Journal of Public Health, Vol. 59, No. 10, pp 1809 - 1815; October, 1969.

A broadly-based health system is envisioned with increased choices for the consumer who will increasingly control the marketplace of health as well as the regulatory apparatus. New infrastructures are needed so that the health care system can become a person-oriented one which deals with human needs.

Duhl, Leonard J. (1976). The Promotion and Maintenance of Health: Myth and Reality. Health Promotion Through Designed Environments, pp 27 - 64; Health and Welfare Canada, Health Programs Branch; October, 1976.

In order to effect positive change in the area of physical and social environment and its relationship to health, there must be a new way of conceptualizing the development of new images, and possibly the changing of an institution's constituency. In order to promote the most effective policy-making possible, there should be a meaningful dialogue between those affected by decisions and those making them. A proposal is made for the requirement of Health Impact Statements for every proposed project within and outside the health field to see how it would affect the health of people.

Duhl, Leonard J. (1984). The Promotion and Maintenance of Health: Myth and Reality: Working Paper No. 436. Institute of Urban and Regional Development, University of California, Berkeley CA; November 1984.

The environment is a tool for dealing with the concerns of human beings because human development — people's lives and relationships — are the cornerstones of life. Through policies and actions, we should attempt to maximize the potential of the individual for aliveness, for development, and for quality of life. The processes of healing involve not only the

individual, but the community, tribe, and family as part of the larger social and physical environment. Though we have a responsibility for optimizing the development of individuals toward attaining their health, we have an equal one to look at the healthy development of the environment.

Duhl, Leonard J. (1984). The Healthy City: Working Paper No. 435. Institute of Urban and Regional Development, University of California, Berkeley CA; November 1984.

A healthy city must be a competent one for all its members. Its responses to its developmental needs, its organizations, and its people must be appropriate and effective and it must have the ability to cope with breakdowns of the system and its members. Solutions must come from the grass roots level, and must deal with the underlying issues, not the symptoms.

Dunlop, Marilyn (1987). Soil, Plant Fungi Infecting Humans U.S. Doctors Say: What's New in Medicine. Toronto Star, January 17, 1987.

Types of fungus that once only affected foods such as tomatoes and peaches are turning up as the cause of some diseases, according to Dr. John Rippon, professor of medicine at the University of Chicago. A report in Medical World News is described which detailed the finding of tomato rot fungus in 24 patients and cases of patients infected with about 100 different soil and plant fungi. According to Rippon, the new fungal infections are caused by the increase of people whose immune systems have been suppressed by drugs.

Durlak, Elizabeth (1987). Personal Interview, Elizabeth Durlak, Toronto, Ontario. Telephone contacts, November 19th and November 30th, 1987.

The author speaks as a person who happens to be asthmatic and highly sensitive to inhaled gaseous and particulate pollution. Her illness has been life-threatening, and her inability to work has left her without sufficient independence to arrange all the environmental conditions necessary for survival without heavy dependence on anti-asthmatic medication. She cautions that there are environmental crises in every aspect of our lives, and that the home is no exception to this: much of available housing is dangerous for environmentally susceptible individuals.

She also suggests that one reason that the needs of the handicapped and the elderly have received inadequate attention is the fact that their stories are commonly deleted from public attention, and many able-bodied people are not at all familiar with what life is like for handicapped or elderly individuals, let alone conscious of them as complex human beings of value in their own right. She notes that most handicapped people are

in fact expert problem-solvers, and that they often spend a great deal of their time and effort addressing their own environmental needs.

She also stresses that it is a fallacy to think that a disability or impairment is necessarily stable, or that a handicapped person cannot get more hurt. People who are not able-bodied must diligently avoid environmental hazards that could worsen their disability (e.g. wheelchair accidents on ramps or curbs, slips and falls for the elderly, chemical exposures for asthmatics or other chemically sensitive individuals, etc.) They are also faced with the burden of educating able-bodied people as to what conditions may be hazardous for them, that present fewer risks to the able-bodied, and they commonly encounter misunderstanding on the part of able-bodied people as to the extent of danger presented by seemingly minor hazards.

Edelstein, Barry A.; Michelson, Larry, editors (1986). Handbook of Prevention. Plenum Press, New York, 1986.

A collection of essays about preventive behavior. Includes comprehensive bibliographies in all subject areas.

Edwards, D.D. (1987). ELF Under Suspicion in New Report. Science News, Vol. 132, July 18, 1987; p. 39.

This article reviews a newly released report on a five-year research program funded by the New York State Power Lines Project. The program included 16 studies of possible health effects from 60 Hz electric and magnetic fields.

The research included an epidemiological study performed in the Denver Area by Dr. David A. Savitz of the University of North Carolina (Chapel Hill). Savitz and his co-workers concluded that the cancer risk among children in higher-exposure homes is 1.7 times greater than that among children in lower-exposure homes, and that the chances of developing leukemia in particular are 2.1 times greater.

David O. Carpenter of the State University of New York at Albany reported that there are some learning and neurological effects from extremely low frequency (ELF) electromagnetic field exposure. Pregnant rats exposed to ELF fields for 30 days developed temporary learning problems. Their offspring, exposed both in the womb and for nine days after birth, developed permanent learning deficits.

The study did not prove a definite cause-and-effect relationship between the extremely low-frequency (ELF) fields studied and the biological and behavioural effects observed.

Ehrlich, Richard (1966). Effect of Nitrogen Dioxide on Resistance to Respiratory Infection. *Bacteriological Reviews*, Vol. 30, No. 3, Sept. 1966, p. 604-614.

The effects of exposure to nitrogen dioxide on man and on animals are confined almost exclusively to the respiratory tract. With increasing dosage, the progressive effects of this gas are: odor perception, nasal irritation, difficulty in breathing, acute respiratory irritation, edema and death. Experimental and epidemiological data pertaining to nitrogen dioxide effects in man are sparse, especially in the low concentration level found in community air pollution.

The work reported in this paper suggests a more sensitive indicator of biological effects of nitrogen dioxide, namely, a synergistic effect or secondary effect, demonstrated by reduction in resistance to infection. A single 2-hr. exposure of Swiss albino Webster strain mice or of inbred mice to 3.5 ppm of nitrogen dioxide before or after respiratory challenge with aerosol of *K. pneumoniae* significantly increased mortality. Continuous exposures to 0.5 ppm for 3 months or longer as well as intermittent daily exposures over a 30-day period produced the same effect in mice. The author cautions that extrapolation to humans is speculative only.

Ehrlich, Richard; Findlay, J.C.; Fenters, J.D.; Gardner, D.E. (1977). Health Effects of Short-term Inhalation of Nitrogen Dioxide and Ozone Mixtures. *Environmental Research*; No. 14, pp. 223-231; 1977.

The effects of single and multiple daily 3-hour exposures to nitrogen dioxide (1.5 to 5.0 ppm) and ozone (.05 to .5 ppm) mixtures on the resistance to streptococcal pneumonia were investigated. It is suggested that a synergistic effect might be present upon repeated inhalation of pollutant mixtures, that made them more effective in reducing resistance to respiratory infection.

Eisenbud, Merrill (1978). *Environment, Technology, and Health: Human Ecology in Historical Perspective*. New York University Press; 1978.

Various sources of environmental pollution which create toxic substances are dealt with in the framework of environmental toxicology: the manner in which they reach man after passing through intricate ecological pathways, and their biological effects on human health. Substances covered include pharmaceuticals, food additives, pesticides, dyes, detergents, soaps, atmospheric and liquid wastes from industrial processes, ionizing radiation, microwaves, combustion byproducts from automobiles, and agricultural chemicals. Nuclear power is written about in detail. Many people are influenced by the psychological association between nuclear power and nuclear weapons, while others are concerned about the gaseous

and liquid radioactive wastes discharged during normal operation, the potential dangers from the radioactive clouds that could escape in the event of a nuclear accident, the difficulty of disposing of radioactive waste, the toxicity of plutonium, or the possibility that plutonium might be diverted clandestinely into nuclear weapon production.

Social and behavioral factors have not been sufficiently emphasized in the contemporary environmental movement. Neither have the environmental problems of the impoverished attracted sufficient attention.

Elinson, Lynn (1984). Analysis of Health Status in the Neighbouring Areas of the Junction Triangle. City of Toronto Dept. of Public Health; September 1984.

This study resulted from the results of a study by Dr. Walter O. Spitzer, entitled "A Study of the Health Status of Residents of the Junction Triangle, Toronto", in which health status was examined in the city's highly industrialized Junction Triangle, a comparison census tract and in the census tracts adjacent to the Junction Triangle. For certain health status indicators it appeared that these indicators occurred more often or at higher levels in the adjacent areas compared to either the Triangle or comparison areas, while the Spitzer report only made statistical comparisons between the two latter tracts. The Spitzer data concerning this adjacent area was therefore re-examined and compared statistically with the comparison census tract data and the conclusion that there is not a major health problem in the area was cautiously drawn.

Elliott, Glen R.; Eisdorfer, Carl (1983). Stress and Human Health: Analysis and Implications of Research/A Study by the Institute of Medicine, National Academy of Sciences. Springer Publishing Company, New York; 1983.

There is marked individuality in response to stressful experiences. Both genetic and environmental factors pertinent to such individuality of response are beginning to be clarified and require more study in the future. There is a great need for genetic studies to be integrated with studies of environmentally caused variation in responses to stress. The extent of community disintegration, a lack of social supports, and the number of stressful events in the life of an individual have been found to affect illness, productivity, and life expectancy. Individuals who experience any of a wide range of stressful events or situations are at increased risk of developing a physical or mental disorder. How individuals attempt to cope with stress has been a neglected area of great potential importance. A deeper understanding of coping behaviour can be useful in devising reasonable therapeutic and preventive interventions. The promise is clearest with respect to mental health; but such interventions also have direct relevance to general health, because health-damaging coping efforts such as smoking, alcohol use, and risky driving weigh heavily in the burden of illness.

Elwood, J.M.; Williamson, C.; Stapleton, P.J. (1986). Malignant Melanoma in Relation to Moles, Pigmentation, and Exposure to Fluorescent and Other Lighting Sources. *Br. J. Cancer*, No. 53; pp 65 - 74; 1986.

Interviews were performed on 83 patients with malignant melanoma and on age and sex matched controls from the same hospitals. Significantly increased risks of melanoma were found in subjects with 3 or more raised moles on the upper arms, in association with heavy freckling of the face and arms, and with a tendency to sunburn easily and tan poorly, these factors having independent effects. While no significant and consistent association with exposure to fluorescent light was seen, the observed risks were higher in subjects with greater exposure, and higher in association with exposure to undiffused than to diffused light. Cases had a significantly greater number of hours' exposure to undiffused light than did controls. The associations with fluorescent light exposure were stronger when based on interview data than on a subsequent postal questionnaire. 21 cases and 11 controls reported exposure to unusual occupational lighting sources which may have had an ultraviolet component; these included various intense lighting sources and lamps used in printing and dyeline copying.

Engen, T. (1986). Odors and Sensory Irritants: Effects on Health and Comfort. *Indoor Air*, Vol. 6, 1986; Swedish Council for Building Research, Stockholm Sweden; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

The study of sensations of odour and irritation has been a part of the study of indoor air from the beginning. These sensations are the effects experienced by people due to stimulation of the olfactory and trigeminal nerves — perhaps the first warning signal a person receives that the atmosphere is contaminated. While things such as body odour and tobacco smoke are still topics of research, there is now perhaps even greater concern with the odour of new building materials and related sensory effects associated with modern tight buildings. The human being is an analyzer of indoor air who makes a crucial contribution to the scientific and technological analysis of indoor air. It must be recognized that while scientific and objective measurement of the environment seem generally favoured over potentially subjective information received from the people living in the environment, it is the effect on them as they experience it which provides much of the norms for indoor air research.

English, Kathy (1987). No One Should Have To Live Like This. *Toronto Star*, June 7, 1987.

A single mother with an employment income of \$800 a month pays almost half of her income to live, with her two teenaged sons, in a tiny, poorly maintained, second-floor, one-bedroom apartment in Toronto. She recently broke some ribs falling down a steep staircase that has no protective

rails and one son's asthma is aggravated by poor ventilation. The basic services like heat and plumbing are inadequate. While municipal officials are aware of this and many similar rental properties, they fear pressuring landlords too hard in case scarce accommodation will be taken off the market. The tenant has been waiting more than a year for subsidized housing, but is relatively low on the priority points system that ranks the 6,000 families now on the waiting list and feels that she would be in a better situation if still on welfare assistance.

Epstein, S.S.; Grundy, R.D. (1974). Radiation Exposures from Electronic Products. from Consumer Health and Product Standards—Chemicals, Electronic Products, Radiation; MIT Press, 1974; pp. 218-221.

Ettema, J.H.; Zielhuis, R.L. (1975). Effects of Alcohol, Carbon Monoxide and Trichloroethylene Exposure on Mental Capacity. *Int. Arch. Occup. Environ. Hlth.*, 35, pp 117-132.

Groups of adult male volunteers were exposed to alcohol, carbon monoxide or trichloroethylene. performance in mental loading tests and effects on physiological parameters were studied. Using alcohol as reference substance, producing marked effects at blood alcohol levels greater than 0.3g/l, the effects from carbon monoxide and of trichloroethylene in dosages as encountered in social and occupational life proved to be much less severe. The experiments suggest the possibility of hazardous impairment of performance in older subjects, particularly in cases of combined exposure to various external loads.

Evans, John R. et al (1987). Toward a Shared Direction for Health in Ontario. Ontario Health Review Panel, June 1987.

This is the report of a panel appointed in November 1986 by the Premier of Ontario to review the health status of the province's residents and its health care system. The panel's membership included representatives of various areas of Ontario, health care providers, consumer organizations, and researchers. The review was undertaken within a framework of a broad concept of health which goes far beyond the absence of illness and disease. Seven values were identified as benchmarks for assessing health and health care: equity, quality, comprehensiveness, informed choice, accountability, cost-effectiveness, and commitment to the future.

Although Ontario has a high standard of health and a good health care system, there are pressures for change, created by rising public expectations for health care, constraints on public resources and frustration of health providers caught in the middle. There is a need to place greater emphasis on primary care, to integrate and coordinate services, to achieve a community focus for health, and to increase the emphasis on health promotion and disease prevention. Well-founded recommendations made by credible groups over a period of fifteen years have rarely been

translated into action. Three general issues have been selected which might provide an overall context in which specific problems might be addressed: strengthening the role of the individual, linking the elements of health care delivery, and increasing the emphasis on ambulatory and community-based care. It is recommended that a Premier's Council on Health Strategy be established with membership from cabinet ministers, health care providers and residents.

Farber, Michael (1987). Police Racism Has Deep Roots. The Gazette, Montreal PQ, p A - 3; August 10, 1987.

Despite efforts by the Montreal Police Director to improve race relations there is an uneasy truce between the police and minority groups. Six civil suits have been filed against Montreal Urban Community police this year for alleged racism, brutality and false arrest. A police force merely represents the society it is sworn to protect and does not have a corner on racism. But the columnist suggests that the problem is not so much racism as ignorance of what might constitute a slight, hardly shocking, he says, considering the historically closed nature of French-speaking Quebec.

Faust, Halley S.; Brilliant, Lawrence B. (1981). Is the Diagnosis of "Mass Hysteria" an Excuse for Incomplete Investigation of Low-Level Environmental Contamination?. Journal of Occupational Medicine, Vol.23, No. 1, pp. 22-26; January 1981.

Fein, Greta G.; Schwartz, Pamela M.; Jacobson, Sandra W.; Jacobson, Joseph L. (1983). Environmental Toxins and Behavioral Development: A New Role for Psychological Research. American Psychologist, Vol. 38, No. 11, pp. 1188-1197; November 1983; American Psychological Assoc., Inc.

Childhood exposure to chemicals routinely encountered in the environment has become an issue of scientific and public concern. Recent research has revealed the inadequacy of traditional notions in which chemically induced illness was likened to overt biological disease. The new multiple-effects model emphasizes subtle behavioral alteration as an early sign of toxicity and as evidence that a particular chemical agent may produce long-term impairment in susceptible individuals. For instance, a case is cited where a group of children who had been hospitalized for overt lead poisoning were evaluated using psychometric tests. Of the 19 children for whom IQ scores were available, 17 scored better than 80. This rather mild retardation was typically accompanied by emotional lability, difficulty in concentrating, and poor school performance. Of the six children whose IQ scores ranged from 95 to 106, only one did well in school. The permeability of the placenta to a variety of chemical agents and the special sensitivity of the fetus to some of these agents draws attention to prenatal exposure and the need for prospective longitudinal studies of affective, social, and cognitive development in exposed individuals. Chemicals discussed in this paper are lead, methylmercury and PCBs.

Ferguson, Roy (undated). The Hospital as a Therapeutic Environment. Alberta Children's Hospital.

The physical environment of a hospital should be considered an important part of the patient's treatment. Hospitalization is often a frightening and stressful experience to children. The unfamiliar environment can induce stress; the child may feel insignificant, anonymous, overpowered, isolated, and intimidated by the scale and procedures of the hospital. The provision of familiar environmental cues through attempts to make certain areas of hospitals more home-like can help. And to the young child in particular, play may be instrumental in enabling hospitalization to be a positive experience. For this reason, the recreational areas and play facilities are important to a children's hospital and should be well staffed. For adolescents, greater opportunity to exercise choice and control in the physical environment will most likely result in less time spent resisting authority. Design also needs to take into consideration the need for family participation in the child's hospital stay.

Ferris, B.G. Jr.; Speizer, F.E.; Bishop, Y.M.M.; Spengler, J.D. (1979). Effects of Indoor Environment on Pulmonary Function of Children 6-9 Years Old. American Review of Respiratory Disease, No. 119 (4 pt 2), p. 214; 1979.

A study was conducted into the type of fuel used for cooking in the home. Indoor levels of mass respirable particulate (MRP), sulfur dioxide (SO₂) and nitrogen dioxide (NO₂) were monitored in homes using different cooking fuels. Results of pulmonary function in the various cities in 6 to 9 years olds were analyzed. It was found that the pulmonary function is lower in those children who live in homes where gas is used for cooking as compared with electricity. Households with no air conditioning tended to have children with lower levels of pulmonary function than those households with partial or central air conditioning. These findings suggest that models taking into account effects of air pollution should include the characteristics of the indoor environment.

Ferris, B., Jr. (1986). Epidemiological Studies of Health Disorders Related to Housing. Indoor Air, Vol. 6, 1986; Swedish Council for Building Research, Stockholm Sweden; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

Epidemiology has come of age and should prove to be a useful tool in future evaluations of the effects of indoor pollutants on health. It is essential to institute quality control procedures for both air monitoring data and health data. There is a need for well-designed and carefully executed epidemiologic studies; for direct measurements of exposure of populations; to develop better estimates of dose to develop dose-response relationships; to control for confounding factors such as active or passive cigarette smoking, other pollutants, temperature, season, etc.; and for longitudinal studies to answer more precisely the medical significance of small changes.

Feuerstein, Michael; Sult, Susan; Houle, Manon (1985). Environmental Stressors and Chronic Low Back Pain: Life Events, Family and Work Environment. *Pain*, No. 22, pp 295 - 307; Elsevier Science Publishers; 1985.

Stressors in the family and job environments have been proposed to play a role in the modulation of pain, yet direct empirical support for such a role is limited. This study investigated the relationship between general stress, family and work environments, psychological distress, and pain experience in 33 ambulatory chronic low back pain subjects and 35 healthy controls matched for sex, age, socioeconomic status, weight, and height. Results indicated that environmental stressors, including family conflict, family control, and general stress, were greater in the group with chronic low back pain. Increased family conflict was associated with increased distress and increased pain, while increased family independence was correlated with less distress and increased pain. Less peer cohesion, less physical comfort, and less job clarity were correlated with increased pain, but not distress. Work pressure was associated with decreased depression and less pain. These findings suggested the presence of both stress and operant mechanisms in the modulation of pain in the family, while operant and distraction mechanisms appear to characterize the relationship among work environment factors and pain.

Ficner, C.A.; Riley, M.; Lubun, M. (1985). Improving Indoor Air Quality Through Energy Conservation Programs. Air Pollution Control Association, 1985.

This paper outlines the approach being taken by the Department of Energy, Mines and Resources to ensure that conservation measures do not adversely affect air quality. The view is presented that it is important to focus on developing means which will ensure that accepted good practices can easily be put into effect in the field. A case is made that conventional housing units do not consistently comply with accepted ventilation practice. In larger buildings, ventilation systems frequently do not perform in the way in which they were intended. Thus, the air quality question affects all buildings, whether or not they are energy efficient. It is suggested that we must emphasize the development of techniques for allowing the industry to construct, modify and operate buildings in a way which complies with existing standards.

Finberg, Laurence (1974). Interaction of the Chemical Environment with the Infant and Young Child. *Pediatrics*, Vol. 53, No. 5, Part II, pp. 831-836; May 1974.

Chemical products and by-products of our society have special implications for infants and children, because maturation and growth processes are often qualitatively or quantitatively different from mature systems and because the infant and child have different environments from adults. In turn the effects may be immediate or delayed, and they may have pro-

found effects on the organism at different points in life. Finally, it would appear advisable to encourage both close epidemiologic surveillance and careful research on the effects of known pollutants. In particular, the tools of modern cell biology and molecular biology may have useful application to these problems.

Finkel, A.S.; and Duel, W.C. editors (1976). Clinical Implications of Air Pollution Research. Publishing Sciences Group, Inc., Acton, MA, USA; McAinsh & Company, Ltd., 1835 Yonge Street, Toronto Canada M4S 1L6..

This proceedings document describes the state-of-the-art at the time of the conference on the relationship between air pollution and cardiovascular and respiratory illnesses. Highlights include a number of chapters on carbon monoxide and cardiac performance (Ch. 9, Edward Haak, Ch. 10, D.A. DeBias), effects on immune defenses and asthma (Ch. 12, G.M. Green), and clinical implications of air pollution research for hypersensitivity diseases (Ch. 15, R.E. Smith). Chapter 16 (R.I. Henkin) discusses the effects of vapour phase pollutants on nervous system and sensory functions.

Finkel, Joe M.; James, Ruby H.; Miller, Herbert C. (1979). Residual Monomers in Acrylic and Modacrylic Fibers and Fabrics: Final Report. US Govt.; July 1979.

Keywords: acrylic resins; textile industry; acrylonitrile; vinylidene chloride resins; monomers; indoor air pollution;

Finn, Peter; McNeil, Taylor (1987). The Response of the Criminal Justice System to Bias Crime: An Exploratory Review. National Institute of Justice, U.S.; October 7, 1987.

This U.S. federally commissioned report, was prepared by Abt Associates for the U.S. National Institute of Justice (a division of the Department of Justice). The study deals with bias crimes against minority groups, and the response of the criminal justice system. The authors conclude that homosexuals are probably the most frequent victims of hate-motivated violence, and are targeted for assault, verbal intimidation, and vandalism more often than blacks, Hispanics, Southeast Asians, and Jews.

Finn, Ronald (1986). Environmental Medicine and the New Allergy. Clinical Ecology, Vol. IV, No. 1, Spring 1986.

Although there are many thousands of diseases, they must all be due to one of three causes — genetic, degenerative, or environmental. Environmental disease is of major clinical importance because genetic degenerative diseases have limited treatment possibilities, whereas removal of an environmental factor which is causing a disease will halt the progress of a disease or even prevent it from occurring.

A wider view of allergy is now gaining acceptance. Reactions are not limited to IgE and can be mediated by other immunoglobulins, and clinical sensitivities can also be caused by non-immunological mechanisms. The clinical spectrum of conditions in which environmental factors are of clinical importance is being expanded. Clinical observations and controlled studies are appearing which confirm that environmental control is of value in the management of many diseases.

Fischer, Claude S.; Baldassare, Mark; Ofshe, Richard J. (1975). Crowding Studies and Urban Life: A Critical Review. *Journal of the American Institute of Planners*, No. 41, pp 406 - 418; November, 1975.

The crowding literature is reviewed with particular concern for its implications for urban life. It was found that much of the research is methodologically or conceptually limited, and that density, though perceived as unpleasant, does not appear to have definite and consistent detrimental effects. Several theoretical frameworks for guiding future research on the effects of urban density are presented. A comprehensive bibliography is included.

Fischer, Claude S. (1982). What Do We Mean by 'Friend'? An Inductive Study. *Social Network*, No. 3, pp 287 - 306; 1982.

The study of friendship is a popular and important one to the study of how social support networks influence health and stress coping mechanisms. However, the concept of 'friend' is ambiguous. This study was undertaken in order to determine what aspects of real relations are correlated with applications of the label 'friend'. In a survey of 1050 adults, the authors obtained the names and descriptions of 19,417 associates. Of these, 59% were labelled 'friends'. Several regression analyses suggest that this label is likely to be applied: to an overwhelming majority of non-relatives in a largely unsystematic way; to associates lacking other, specialized role-relations; to people of the same age; to people known a long time; and to people with whom respondents had primarily sociable, rather than intimate or material, involvements.

Fischer, Claude (1982). *To Dwell Among Friends: Personal Networks in Town and City*. University of Chicago Press; 1982.

Modern urban life has destroyed community and caused it to disintegrate into a mass of atomistic and alienated individuals. This is a study of personal networks in city and small town environments. All else held constant, respondents with below-average household incomes who lived in semirural communities tended to have more local nonkin associates, bound up in denser networks, than did low-income respondents residing in other places. The semirural people tended also to report a slightly more positive sense of well-being than their counterparts elsewhere. This is a thread that ought to be followed in future research. A comprehensive bibliography is included.

Fishbeck, William A.; Townsend, Jean C.; Swank, Marlene G. (1978). Effects of Chronic Occupational Exposure to Measured Concentrations of Benzene. *Journal of Occupational Medicine*, Vol. 20, No. 8, pp. 539-542; 1978.

Fisher, Alexander A. (1983). Paper Dermatitis: Current Contact News. *CUTIS*, Vol. 31, pp. 392-405; April 1983.

Various types of chemicals are found in different papers. Formaldehyde, ammonia, hydrogen sulfide, sulfuric acid, carbon monoxide, sodium hydroxide and chlorine, among others are used. The handling of most newspapers, magazines, hardcover and paperback books, art paper, paper tissues, plates and cups and glossy paper readily produces dermatitis in formaldehyde-sensitive individuals. Carbonless paper can produce mild dermatitis and severe upper respiratory symptoms as well as eye irritation. Chemical agents in various types of photocopier papers also causes dermatitis in some people.

Fleishman, Jane (1984). There's Something in the Air...: from Double Exposure: Women's Health Hazards on the Job and at Home, edited by Wendy Chavkin, Monthly Review Press, New York NY. Working Mother, pp. 114-118; October 1984.

Occupational health experts, health professionals, working women's organizations, unions and clerical workers themselves have begun to pay close attention to the health hazards of the office workplace. Such things as low-level exposure to various office chemicals, including those used in office machinery, to artificial lighting, to inadequate ventilation, to high noise levels and to low-level radiation, are all being looked at as health hazards creating actual sickness and stress.

Foote, Robert S. (1972). Mercury Vapor Concentrations Inside Buildings: Reports. *Science*, Vol. 177, pp. 513-514; August 1972.

Franck, Karen A. (1985). Change: A Central But Unheralded Theme in Environmental Design Research. *Environmental Change/Social Change*, pp 4 - 9; Klein, S., Wener, R., Lehman, S, editors; Environmental Design Research Association; June 1985.

If we are interested in encouraging environmental improvements which we judge to be important, we need to hone our analytic skills and to apply them to describe existing environments. One recommendation is to analyze a building's hidden program, that is, the system of mutually reinforcing social, political, and economic forces that lead to a recurring spatial relationship and physical form; and to restructure that program to produce an alternative set of spatial relationships and an alternative physical form that are more supportive of the forces that one feels should be reinforced.

Franklin, Claire A; Burnett, Richard T.; Paolini, Richard J.P.; Raizenne, Mark E. (1985). Health Risks from Acid Rain: A Canadian Perspective. Environmental Health Perspectives, Vol. 63, pp 155 - 168; 1985.

Acid rain is causing serious environmental damage in eastern Canada. The revenues from forest products, tourism and sport fishing are estimated to account for about 8% of the gross national product. The impact on human health is not as clearcut. A multi-department program on the Long-Range Transport of Airborne Pollutants (LRTAP) was approved by the federal government in June 1980. The objectives of the program are to reduce wet sulfate depositions to less than 20 kg/ha per year in order to protect moderately sensitive areas. This will require a 50% reduction in Canadian SO₂ emissions east of the Saskatchewan/Manitoba border and concomitant reductions in the eastern USA. The objectives of the health sector of the program are to assess the risk to health posed by airborne pollutants which are subjected to long-range transport and to monitor the influence of abatement programs. Two major epidemiology studies were undertaken in 1983. Preliminary analysis of the data do not indicate major health effects, but definitive conclusions must await final analysis. Studies on the indirect effects of acid depositions on water quality have shown that acidified lake water left standing in the plumbing system can adversely affect water quality and that federally set guidelines for copper and lead are exceeded. Flushing of the system before using the water rectifies the situation. Additional studies are planned to further delineate the magnitude of the health effects of acidified lake water.

Friedman, Mitchell; Dougherty, Richard; Nelson, Steven R.; White, Robert P.; Sackner, Marvin A.; Wanner, Adam (1977). Acute Effects of an Aerosol Hair Spray on Tracheal Mucociliary Transport. American Review of Respiratory Disease, Vol. 116, pp. 281-286; 1977.

Garling, Tommy; Valsiner, Jaan (1985). Children Within Environments. Plenum Press, New York; 1985.

The most frequent harm to children is accidental injuries. This book presents a view on the prevention of childhood accidents emerging from a discussion of children and how they relate to their environments.

Gaudert, P.C. (1985). Briefing Paper on the Status of DBR Research on Contaminant Emissions from Wood-Based Composition Boards. unpublished; enquire P.C. Gaudert, DBR, NRC K1A 0R6.

Wood based composite boards bonded with urea formaldehyde (UF) resin - especially particleboard and interior type plywood grades - have become one of the most widely used base materials for interior construction and furnishings. Such board can introduce about 50 to 150 g. of resin for

each cubic metre of air in the living space. A yearly release of as little as one to two percent of the formaldehyde contained in the resin into the air of moderately "tight", i.e. energy efficient, homes can cause the ambient concentrations to exceed the recommended 0.1 ppm limit for formaldehyde. A surplus of formaldehyde from the manufacturing process and a slowly decreasing generation of formaldehyde from hydrolysis result in an exponential emission decay with a half-life of about three to five years, i.e. the emission will decrease by half in this time.

Simulation of emissions from samples of representative, unfinished board, using dynamic chambers, have indicated that an unrestricted flow of emissions can cause ambient indoor concentrations to approach 1 ppm. In practice this concentration is usually not reached because of the diffusion resistance introduced by various surface finishes, stagnant air layers or the restriction of flow due to enclosures such as cupboards and closets.

Gelfand, Donna M.; Ficula Teresa; Zarbatany, Lynne (1986). Prevention of Childhood Behavior Disorders: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 133 - 152.

Prevention of behavior problems in children depends largely on our ability to predict future maladjustments. Accurate prediction is often hampered by our lack of knowledge regarding the process by which developmental and environmental factors combine to influence the child's psychological functioning. Risk factors for children's behavioral and psychological disturbance involve a dysfunction in some aspect of the child's environment that is critical to the child's mental health such as the social community, the family, the schools. Many suggestions are made for primary prevention programs which involve restructuring the social community to remove the offending features such as poverty and lack of good housing, violence in the media, child abuse, parents who engage in antisocial behavior and alcohol abuse. Other suggestions are to provide support services for children and to help them develop coping skills for dealing with a less than optimal environment.

Geller, E. Scott (1986). Prevention of Environmental Problems: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 361 - 383.

Many solutions to environmental problems require changes in human behavior, and behavioral science can offer suggestions for attacking the behavioral aspects of environmental protection. A multi-level program is proposed for environmental control and an organizational framework is presented for an energy conserving community.

Gerdas, Kendall (1987). Letter from Dr. Kendall Gerdes, Environmental Medicine Associates. Environmental Medicine Associates, 1617 Vine St., Denver CO 80206-1117; July 10, 1987.

The author, a specialist in environmental medicine, notes that we have in the past defined health as an absence of disease. He suggests that it is considerably more than that, embodying features of sustained optimal functioning or doing every day what is possible on a person's best day. He observes that the concept of health does not include the highest functions of human beings, as typified by the Ghandis or the Mother Teresas of the world. He is convinced that we cannot create an environment which grows that type of individual, but by making the environment negative, we can prevent that type of individual from developing. One suggestion is to avoid xenobiotic chemicals. He also suggests gathering a small group of healthy Canadians and studying them in terms of both biochemistry and in terms of social structure and environment.

Gershon, S.; Shaw, F.H. (1961). Psychiatric Sequelae of Chronic Exposure to Organophosphorus Insecticides. *Lancet*, 1(7191), pp 1371 - 1373; 1961.

A study of 14 men and 2 women exposed to organophosphorus insecticides for periods ranging from 1 1/2 to 10 years showed schizophrenic and depressive reactions with severe memory impairment and concentration difficulty. A field survey gave some support to the possibility that psychiatric disorders might be less frequent in towns than in fruit-growing regions of Australia where the insecticides were used. Follow-up showed that the effects of poisoning lasted for at least 6 months, but nearly all patients reverted to normal within a year.

Gibson, Bonnie (1985). Spotlight on Cadmium. *Canadian Environmental Law Association Newsletter*, Vol. 10, No. 1, pp. 10 -13; February 1985.

Cadmium is a naturally-occurring heavy metal that is increasingly being released into the environment as a result of human activity. It is a highly toxic substance with no known biological function and cannot be destroyed by any environmental process. It is produced as a by-product of zinc manufacturing. The major source of exposure is inhalation in the industrial setting and as a result of accidental releases from chemical and electro-plating plants. Another source of exposure is through food and tobacco, both as a result of soil uptake of naturally-occurring cadmium as well as fertilizer residue. Concentrations of cadmium are considerably higher in second-hand tobacco smoke than to the smoker. Canadians, on an average, exceed the World Health Organization's suggested maximum permissible dietary intake level of .07 milligrams per person per day. Health effects can be lung damage and kidney damage or failure. A lethal dose is possible without warning due to lack of discomfort at the time of exposure. It accumulates in tissues and its chemical half-life can be as high as 30 years.

Giddings, Michele J. (1986). Occupational Health Problems Due to Air Exposure to Fungal Contamination. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Occupational hazards from the exposure to organic dusts, specifically fungi, in the workplace do exist. These diseases may be so mild as to be asymptomatic or be extremely debilitating. There are other diseases which may also be considered as resulting from occupational exposure. They include buildings whose humidifier and/or ventilation systems are contaminated with fungi allowing for the development of symptoms among the occupants. Recognizing the possible hazards associated with one's chosen occupation, whether it be exposure to organic dusts, chemicals or dangerous machinery is of greatest importance for prevention of injury or infection. Persistent symptoms, no matter how small, should be regarded as important and measures taken to locate their source. Care must be taken when identifying the cause of a worker's complaint, as these diseases may be misdiagnosed which in turn may lead to irreversible lung damage. An extensive bibliographic reference list is included.

Gilka, L. (1978). Childhood Experiences As Causes of Criminal Behaviour: The Report to Senator Fred A. McGrand, Chairman, The Senate Subcommittee on Childhood Experiences as Causes of Criminal Behaviour. The Society for the Understanding of Nutrition, Ottawa ON K2B 5W6; April 1978.

An increasing number of scientists and physicians are now concluding that in many children learning disabilities may be the result of nutritional deficiencies and food and other allergies. There is now also sufficient evidence of the connection between juvenile delinquency and learning disabilities.

Girman, J.R.; Apte M.G.; Traynor, G.W.; Allen, J.,R.; Hollowell, C.D. (1982). Pollutant Emission Rates from Indoor Combustion Appliances and Sidestream Cigarette Smoke. University of California, Lawrence Berkeley Laboratory, Energy & Environment Division, Berkeley CA; May 1982.

Keywords: gas appliances; particulate emissions; ventilation rate; gas fired cooking stoves; unvented kerosene space heaters; unvented natural gas space heaters; nitrogen dioxide; NHW

Godish, Thad (1984). Indoor Air Pollution—Offices and Other Public Access Buildings. Natural Resources Notes, No. 8; Fall 1984.

Within the last decade there have been hundreds of reported outbreaks of illness among occupants of new or recently remodelled offices, schools, and other public access buildings. These outbreaks have

characteristically affected a large number of individuals, in some cases 30-40% of the building's population. Illness symptoms are usually of a non-specific nature often including headaches, dizziness, unusual fatigue, eye, nose and throat irritation, chest tightness and shortness of breath. The building materials and lack of ventilation are often the cause, but furnishings, office equipment, maintenance materials, accidental spills of chemicals, cross contamination from outside, tobacco smoke and microbial contaminants can also be at fault. Investigation procedures and mitigation measures are discussed.

Goldberg, Susan (1985). Facing the Nuclear Age: Parents and Children Together. Annick Press Ltd., Toronto Canada; 1985.

Researchers are demonstrating that children are aware of the arms race much earlier than adults suspect. This awareness and the perception that few adults are doing anything to prevent a nuclear war are described as resulting in a profound distrust of adults and adult values. This resource book examines the extent of the problem and is designed to help initiate discussion around the issue with children. It suggests practical responses to the questions. A complete resource list of films, videos, books, organizations, and articles on the psychological aspects of the arms race is included.

Good, Bennie W.; Vileins, G.; Harvey, W.R.; Clabo, D.A., Jr.; Lewis, A.L. (1982). Effect of Cigarette Smoking on Residential NO₂ Levels: from Indoor Air Pollution; Spengler, John—editor; pp. 167-175. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Two studies evaluating the levels and sources of nitrogen dioxide in approximately 90 employee homes in the Richmond area were performed using samplers in various rooms and outdoors. Additional data were collected concerning appliance usage, heating/cooling plant, ventilation and cigarette smoking. The largest contributor to NO₂ concentration was found to be gas-fired kitchen appliances, completely overpowering that due to the contribution from cigarettes. Nitrogen dioxide is corrosive, reactive and highly oxidizing, and may be toxic at high concentrations.

Gordis, Leon (1986). Geographic and Environmental Factors in Pediatric Cancer. Cancer, No. 58, pp 546 - 549; 1986.

It is important to determine the relative contributions of genetic and environmental factors to the etiology of childhood cancer in order to discover the pathogenic mechanisms involved and to develop effective means of primary prevention. Geographic differences in cancer incidence as well as changes in incidence over calendar time have long been used to generate clues to possible etiologic agents. The important role of genetic factors in childhood cancer is clear. The importance of the

contributions of environmental factors in general and of specific factors in particular has proven more difficult to determine. A variety of environmental factors have been implicated to varying degrees in the etiology of different childhood cancers. These factors include physical agents such as radiation, chemical agents such as nitrosamines, and organic solvents, and infectious agents such as the Epstein-Barr virus. The observations that certain compounds may act as teratogens when a pre-natal exposure occurs early in pregnancy and as carcinogens when the exposure occurs late in pregnancy, suggests that there may be a continuum of teratogenesis and carcinogenesis. This finding has major implications for the possible biologic mechanisms that could be involved in childhood cancers and for the design of future research of their etiology and prevention. The etiology of childhood cancer should be viewed as an interaction of environmental factors to which the child or his or her parent were exposed together with varying degrees of genetically determined susceptibility of the child to the carcinogenic effects of these factors.

Gordon, Michael (1982). Falls in the Elderly: More Common, More Dangerous. *Geriatrics*, Vol. 37, No. 4, pp. 117-120; April 1982.

Keywords: accidents; aged; aging; arrhythmia; dizziness; women;

Gosnell, Denise (1987). Minutes of the Pollution and Education Committee, Toronto Board of Education, November 17, 1987. Toronto Board of Education, 155 College St., Toronto M5T 1P6..

The Toronto Board of Education's Pollution and Education Committee (Trustee level) considered total building performance investigations as a means of addressing indoor environment concerns in Toronto schools and other buildings. The procedure recognizes that a variety of elements contribute to occupants' satisfaction with a building environment, and also recognizes the special needs for those who experience greater sensitivity to the local environment. The procedure encourages all occupants to participate in the planning, design, problem-solving and operating practices of their building. Part of the rationale for the procedure is to give individuals or groups the tools to solve building environment problems.

The Committee also authorized the Director of Education, in cooperation with the Working Group of Ontario School Boards on Pollution and Education, to investigate with recognized testing laboratories the development of tests and standards for pollutants, which will assist the Toronto Board in evaluating materials which are being considered for purchase and use in the school system. The Committee voiced a parallel intent to investigate applications for rezoning of land within approximately 1000 feet of school sites, with respect to pollution from outdoor sources.

Gottlieb, Nell H.; Baker, Judith A. (1986). The Relative Influence of Health Beliefs, Parental and Peer Behaviors and Exercise Program Participation on Smoking, Alcohol Use and Physical Activity. Soc Sci Med, Vol. 22, No. 9; pp 915 - 927; 1986.

This research specifies a model for lifestyle health behavior which includes socialization, social environmental and cognitive influences on smoking, alcohol use and exercise. Survey respondents were participants in university conditioning classes, academic health classes and a community fitness program. Of the socialization influences, drinking by both parents was directly related to drinking by females and drinking by the mother to that of males. Physical activity level of the father was associated with the activity of both genders. Parental smoking was not related to that of the respondents. Income was positively correlated with alcohol consumption in both genders and with smoking among women. In the immediate environment, drinking and smoking by male and female friends were directly associated with that of both male and female respondents with congruent gender relationships being strongest. Exercise by male friends was positively associated with activity level for both genders. For cognitive factors, belief in the efficacy of lifestyle change was inversely related to alcohol consumption for both genders and to smoking for women. Susceptibility to heart disease was associated with a low exercise frequency and smoking. With one exception, the peer modeling variables had the strongest relationships of any of the model elements. These findings suggest that prevention programs incorporate strategies to maximize peer support for healthful behavior and to counteract the effects of unhealthful behavior modeled by peers. Interventions to increase beliefs in the efficacy of lifestyle change to reduce risk are appropriate to encourage behavior change. When the change is underway, discussion of lowered susceptibility as a function of program compliance should reinforce the new behavior.

Gravesen, S.; Larsen, L.; Gyntelberg, F.; Skov, P. (1986). Demonstration of Microorganisms and Dust in Schools and Offices: An Observational Study of Non-Industrial Buildings. Allergy, No. 41, pp 520 - 525; 1986.

The sick building syndrome is reported with increasing intensity in non-industrial places of work, such as schools, kindergartens, and offices, all of which have a heavy load of traffic (people). The construction of the buildings, with flat roofs, often leads to water damage with subsequent microbial growth. Further, reduced cleaning budgets in connection with wide use of needle-felt carpets, as well as ventilation systems not regularly maintained, will lead to pollution by dust and microorganisms. A systematic registration of dust and microbial parameters has been carried out since 1980 in buildings with indoor climate complaints, in order to elucidate the possible influence of these factors.

Green, G.H. (1979). Field Studies of the Effect of Air Humidity on Respiratory Diseases: from Indoor Climate: Effects on Human Comfort, Performance, and Health in Residential, Commercial, and Light-industry Buildings; Fanger, P.O. and Valbjorn, O.—editors; pp. 207-223. Danish Building Research Institute, Copenhagen Denmark; 1979.

Studies have been undertaken of the effect of increasing indoor relative humidity on respiratory diseases. Kindergarten children, schoolchildren, soldiers and office workers were studied. The groups were split into rooms that were identical except for differing levels of humidity. There were significantly lower absenteeism and/or occurrences of respiratory illnesses in the spaces with the higher humidities. Increased winter relative humidity of 8 to 10% in the 22 to 50% range reduced the absenteeism from 10% in adults to 50% in kindergarten children.

Green, K.B.; Pasternack, B.S.; Shore, R.E. (1982). Effects of Aircraft Noise on Reading Ability of School Age Children. Archives of Environmental Health-CISTI; Vol. 37, No. 1, pp. 24-31; 1982.

The percent of students reading below grade level from 1972 to 1976 was regressed on racial, socioeconomic, educational, and noise level variables for all elementary schools in Brooklyn and Queens, New York. Schools were assigned noise exposure scores based on Noise Exposure Forecast contours for New York City airports. The correlations between these noise scores and a variety of noise level metrics ranged from 0.74 to 0.97. The regression coefficients adjusted for confounding factors, indicated that an additional 3.6% of the students in the noisiest schools read at least 1 year below grade level with 95% confidence limits from 1.5 to 5.8%. The dose-response relationship indicated that the percent reading below grade level increased as noise level increased.

Green, Kendall B.; Pasternack, Bernard S.; Shore, Roy E. (1982). Effects of Aircraft Noise on Hearing Ability of School-Age Children. Archives of Environmental Health, Vol. 37, No. 5, pp. 284-289; September/ October 1982.

201 cases with permanent bilateral high-frequency hearing loss and 208 controls with normal hearing were identified through the New York City hearing screening program. Aircraft noise exposure was estimated for the residences of cases and controls from Noise Exposure Forecast contour maps of the New York City airports. The noise exposure estimates had a correlation on 0.89 with noise level measurement made in the area. Birth certificates and questionnaires were used to provide additional information on other sources of noise exposure and potential confounders. The results showed a positive, but not statistically significant association between aircraft noise exposure and the risk of high-frequency hearing loss.

Greer, Nora Richter (1986). The Search For Shelter. The American Institute of Architects, Washington DC; 1986.

Many Americans are homeless. They have a right to dignified shelter which is investigated in this book. Case studies are presented.

Greer, Nora Richter (1986). Paths to Homelessness. The Search For Shelter; pp. 17 - 22; The American Institute of Architects, Washington DC; 1986.

The immediate causes of homelessness are diverse: loss of a job, physical or mental disability, loss of benefits, divorce, etc. But the plight of the homeless represents the most severe housing shortage since the Depression. The problem centres on the dwindling supply of housing for those with the lowest incomes.

Greer, Nora Richter (1986). Dignified Shelter. The Search For Shelter; pp. 31 - 35; The American Institute of Architects, Washington DC; 1986.

When designing shelters and programs for the homeless, it is perhaps most important to create a sense of dignity. Design considerations are presented which would create shelters and more permanent housing, as well as service delivery vehicles which would be dignified and take into account the physical as well as the psychological needs of the homeless.

Greer, Nora Richter (1986). The Right To Shelter. The Search For Shelter; pp. 37 - 41; The American Institute of Architects, Washington DC; 1986.

One of the biggest obstacles facing developers of shelter or low-income housing is community resistance. The 'not in my backyard' phenomenon is occurring in countless communities. Strategies for helping communities feel comfortable with this type of housing are presented. These include communicating with the community at an early stage about the proposed facility and how it works, promising priority placement in the shelter to people from the surrounding community, and designing it to be compatible with surrounding buildings. Reevaluation must also occur of municipal building code definitions of suitable, affordable housing, as well as that of emergency and transitional shelter.

Gregory, Judith (1981). Office Air Quality, 'Tight Buildings', and Job Stress - The Impact on Women Office Workers' Occupational Health. Working Women Education Fund, 1224 Huron Rd., Cleveland OH 44115; May 25 1981.

Keywords: indoor air pollution; occupational hazards; stress; workplace; ventilation; offices;

Griffith, Ezra E.H.; Griffith, Elwin J. (1986). Racism, Psychological Injury, and Compensatory Damages. Hospital and Community Psychiatry, Vol. 37, No. 1; pp 71 - 75; January, 1986.

Psychiatrists and other mental health professionals have paid only modest attention to the idea that discriminatory conduct causes emotional suffering for those who are the object of it. However, courts have held that if such racist conduct is willful and outrageous and the ensuing suffering is severe, the plaintiff has a reasonable claim to compensatory damages. The authors trace these developments in the legal area in both tort actions and complaints under civil rights statutes. They also point out how psychiatrists could be more influential in sharpening considerations about the idea that racism causes psychological injury.

Grimsrud, D.T.; Sherman, M.H. (1982). Energy Efficient Domestic Ventilation Systems for Achieving Acceptable Indoor Air Quality: A Comparison of Alternate Ventilation Strategies. The Air Infiltration Centre, Old Bracknell Lane W., Bracknell, Berkshire RG12 4AH England; 1982.

Keywords: ventilation; measurement of pollution; indoor air quality;

Guidotti, Tee L.; Goldsmith, John R. (1983). Air Pollution and Family Health. American Family Physician, Volume 27, number 4, pp 165-172, April 1983.

Air pollution is classified according to chemical characteristics of the pollutants and sources: reducing air pollution, photochemical air pollution, point-source emission and indoor air pollution. The primary pollutants are particulates, oxides of sulfur and nitrogen, ozone, hydrocarbons, carbon monoxide and lead. A knowledge of the effects of these on healthy people and impaired patients can provide the physician with specific measures to apply in individual cases. These effects are described under the headings: respiratory system effects, cardiovascular effects, mucosal irritation, carcinogenesis, neurologic effects.

Gupta, K.C.; Ulsamer, A.G.; Preuss, P.W. (1982). Formaldehyde in Indoor Air: Sources and Toxicity: from Indoor Air Pollution; Spengler, John—editor; pp. 349-358. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Formaldehyde, a highly reactive gas with a pungent odour, is released from a variety of sources including urea-formaldehyde foam insulation, particle board, and plywood, as well as various combustion processes. Concentrations of formaldehyde associated with the presence of these products are higher indoors than outdoors. Under controlled conditions, formaldehyde causes eye and nasal irritation at air concentrations of 0.24 mg/m³ and above. Exposure, residential or occupational, has been

associated with eye, nose, and throat irritation, coughing, wheezing, skin rashes, nausea, and other symptoms. Formaldehyde is also a sensitizer; individuals who are allergic to it, or who suffer from respiratory diseases, are likely to suffer its effects at even lower concentrations. Based on its known metabolism, reactivity with DNA and other macromolecules, as well as its mutagenic effects, formaldehyde is thought to be genotoxic. Recent studies have also indicated that it is a carcinogen in rats and probably in mice.

Haan, Mary; Kaplan, George A.; Camacho, Terry (1987). Poverty and Health: Prospective Evidence from the Alameda County Study. American Journal of Epidemiology, Vol. 125, No. 6, pp 989 - 998; 1987.

To examine the reasons for the association between socioeconomic status and poor health, the authors examined the nine-year mortality experience of a random sample of residents aged 35 and over in Oakland, California. Residents of a federally designated poverty area experienced higher age-, race-, and sex-adjusted mortality over the follow-up period compared with residents of nonpoverty areas. This increased risk of death persisted when there was multivariate adjustment for baseline health status, race, income, employment status, access to medical care, health insurance coverage, smoking, alcohol consumption, physical activity, body mass index, sleep patterns, social isolation, marital status, depression, and personal uncertainty. These results support the hypothesis that properties of the sociophysical environment may be important contributors to the association between low socioeconomic status and excess mortality, and that this contribution is independent of individual behaviours.

Haider, M.; Groll-Knapp, E.; Holler, H.; Neuberger, M.; Stidl, H. (1976). Effects of Moderate CO Dose on the Central Nervous System—Electrophysiological and Behaviour Data and Clinical Relevance. from Clinical Implications of Air Pollution Research, Finkel and Dael, editors; Chapter 17, pp. 217-232; Publishing Sciences Group, Inc.; 1976.

Many aspects of human performance are apparently affected even by low CO levels, while others show an impairment only considerably later. The capacity for observation in situations involving comparatively weak stimulation could be a very sensitive indicator, while complex psychomotor dexterity is only slightly affected. Sleeplessness, irritability, restlessness, headaches, disturbances of well-being are all symptoms that have many causes. One of these may be CO, but each case has to be tested individually and further research is needed to clarify the extent to which such a statement can be verified.

Hall, Chris (1987). Aging Well: Focus on Growing Old and Staying Gay. Xtra, No. 89, p. 5; Nov, 27, 1987.

A November 7-8, 1987 Conference on Gays and Lesbians Aging reviewed the problems and options faced by older gay people in Canada. While gay people have many of the same needs in aging as their heterosexual counterparts, gay people also have many needs that are unique to them. In housing, for example, there is no existing structure that allows for lifelong lovers to share facilities in an institutional setting. The conference stressed the importance of maintaining good health to increase the quality of life in the later years. Workshops were held on maintaining independence in the home, creating change within the system, and developing a supportive social network.

Hall, Muriel (1987). Frustrations of Hypersensitivity. unpublished; July, 1987.

The most demoralizing and frustrating experience for the person with chemical hypersensitivity is that, while being seriously ill, weak, exhausted, confused, in pain, and isolated, one must also fight spouse, children, extended family, neighbors, friends, community, all levels of government, doctors, hospitals, social services, OHIP, OMA, psychiatrists, employers, insurance companies, pension and/or welfare agencies — in fact, the whole world — in order to survive.

Hall, Muriel (1987). Healthy Living. unpublished; September, 1987.

Some of the frustrations coping with hypersensitivities in a world full of toxins, in a society geared to oppression of misfits (with psychosomatic illness), are perfumes built into stationery and periodicals; visitors who lack the knowledge or sensitivity to eliminate completely the use of cosmetics, body and clothing care products which contain perfumes; relatives who treat the illness as psychosomatic and who ignore the environmental causes of the illness, refusing to change them; and the incredible feeling of hopelessness at being misunderstood, put-down, ignored, talked over, assumed to be retarded or hard of hearing. Education of patients, families, communities, church groups, the medical profession, educators, social and welfare agencies, law enforcement officers, penal institutions, and other providers of housing and services is necessary.

Halton, David M. (1983). Occupational Exposures from Spirit-Duplicator Operations. Canadian Centre for Occupational Health & Safety, Hamilton ON L8N 1H6; October 1983.

The current technology used in spirit duplicating is discussed. A number of studies have indicated the potential for methanol overexposure from these machines. These are described here, along with the negative health effects. The various exposure-control strategies are reviewed.

Halton, David M. (1983). Photocopiers—Do They Pose a Health Hazard?. Canadian Centre for Occupational Health & Safety, Hamilton ON; April 1983.

The current technologies used in photocopying machines are briefly described. Concerns that have been raised about health and safety at each step of the copying process are discussed. Particular emphasis has been placed on the chemical components of the technology. Areas of specific concern are : the refilling and disposal techniques for toner in dry machines; levels of isodecane produced by LTT machines; levels of ozone produced; heat and noise levels; and escape of ultraviolet light.

Hambraeus, A. (1986). Microorganisms Related to Buildings. Indoor Air, Vol. 6, 1986; Swedish Council for Building Research, Stockholm Sweden; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

Very little knowledge is available concerning the effect of the building-related microorganisms on people or buildings. More information on the clinical significance is needed.

For instance, in a simulated office space, it was shown that the HVAC system operational parameters affect airborne fungal levels in occupied spaces. The reduction of airborne fungi correlated well to the number of airchanges when the filter system was intact. When the main air handling unit was operated without its filter bank, fungi were dispersed into the air. The fan coil units were found to be important sources of microbial aerosols. Indoor air always contains viable mould spores but in buildings that do not generate their own sources of mould, the numbers are usually low and the species found are similar to those found in outdoor air. The classical sources of in-house generated mould are found in high water vapour risk areas such as bathroom and kitchen. Problems in this area can be increased due to reduced ventilation which has been introduced for energy saving. There are other problems that might be caused by microorganisms in buildings. For instance, the presence of occupational asthma and humidifier fever was investigated in two air conditioned office buildings which had humidifiers designed to produce around 50% relative humidity. And in newly built homes where self-leveling cement has been used, problems with foul smell and miscoloured floors have occurred. Occupants and office workers have complained of symptoms such as headache, tiredness, and allergic reactions. Microorganisms have been suspected to cause degradation of the protein in the cement.

Hancock, Trevor (1980). The Soft Health Path: An Alternative Future for Health in the 80's. March, 1980.

A new approach to health care is emerging which can be described as the Soft Health Path. It is a synthesis of environmental health, community-based public health, the primary health care concept, the holistic health movement and the self-care movement. It is compatible with and leads towards a sane, humane and ecological future.

Hancock, Trevor (1987). Healthy Public Policy in the Nordic Countries: Preliminary Report: CPHA Study Tour Report. Canadian Journal of Public Health, Vol. 78, pp 9 - 10; February, 1987.

Fourteen Canadians from 5 provinces spent 3 weeks studying health policy and promotion in Denmark, Norway, Sweden, and Finland as a result of the October 1984 Beyond Health Care conference on healthy public policy. The policies in these countries are briefly described. Preliminary conclusions are drawn: 1) Central government leadership is essential in formulating a health policy that moves beyond health care to address health for all and health promotion policy but efforts must be directed to the Provincial ministers of health and their staffs; 2) Local action must be initiated to develop public awareness of Health for All and to develop intersectoral policies to promote health; 3) A greater awareness must be developed of the social nature of health, identifying inequalities in health and making them apparent to policy makers and the public.

Hane, Monica; Axelsson, Olav; Blume, Jan; Hogstedt, Christer; Sundell, Lennart; Ydreborg, Berit (1977). Psychological Function Changes Among House Painters. Scand. j. work environ. & health, No. 3, pp. 91-99; 1977.

Keywords: brain damage; hemoglobin concentration; psychological function tests; solvents; work environment;

Hanvey, Louise (1987). Communiqué. Canadian Institute of Child Health, Ottawa ON; 1987.

Even playgrounds with equipment specifically designed for children can be hazardous places. A 1983 study found that Canadian hospitals 934 injuries resulting from playground apparatus, many from falls. The Canadian Institute of Child Health has released "When Child's Play Is Adult Business", publication designed for parents and concerned citizens which spells out the difference between a safe and dangerous playground.

Harding, Douglas H. (1982). Health Effects of Formaldehyde. Occupational Health in Ontario, Vol. 3, No. 2, pp. 64-80; April 1982.

The possibly erroneous idea that formaldehyde is a carcinogen is explained in this review. The argument is presented that this substance is an epigenetic carcinogen based on its irritancy and high reactivity. It is the irritancy and not the formaldehyde chemical that induces the chain of events leading to malignancy. The threshold for irritation of formaldehyde is 0.1 ppm, and if such a level were used to control exposure to the chemical, it would be expected that the danger from formaldehyde-induced cancer would be greatly reduced. Possible sources of formaldehyde are listed: cigarette smoke, automotive exhaust, photochemical smog, incinerators, urea formaldehyde in fabrics and building materials. There are individuals who have become hypersensitized to formaldehyde and must avoid it at all costs.

Harrison, John (1983). Artsmart. Living Safety, No. 4, pp. 26-29; Winter 1983.

Art and craft materials can be hazardous to health. Particularly vulnerable are the elderly, children, and people with health problems. Particularly hazardous are ceramics, printmaking, painting and sculpting. Hazards and precautions are described.

Hart, Dr. Roger (date?). Sex Differences in the Use of Outdoor Space. Perspectives on Non-Sexist Education, Chapter 8, pp 101 -109.

The physical environment is essentially boys' domain. A study was conducted of young children who were asked to identify the area in which they were allowed to play outside your house without special parental permission. There were two age groups: five- to eight-year-olds and nine-to-twelve-year-olds. In both age groups the boys' range was larger than the girls' but most markedly so in the older age group. It was concluded that parentally-defined range restrictions restrict the movement of girls in the outdoor environment. It was also found that boys modify the landscape more frequently and more effectively than girls who instead use their imaginations to create outdoor spaces. Girls and boys receive multiple reminders from adults and from peers as to what are and are not suitable activities for them. It seems that girls are practicing and being prepared for roles in the home and boys for roles outside. The very different opportunities given to girls versus boys to freely manipulate the environment affects their spatial abilities and other types of problem solving. It is important that all children have the freedom, ability, and confidence to define and carry out their goals.

Hathaway, Warren E. (1983). Lights, Windows, Color: Elements of the School Environment. Alberta Education, 11160 Jasper Ave, Edmonton Alberta T5K 2L0; January 1983.

Recent research indicates that there are a great many non-visual effects of light on people. Of these effects, the suntan effect and the control of rickets are two that are well understood. Other important but less well known effects include synchronization of a number of physiological rhythms, and prevention or control of infantile jaundice. Still other research indicates that variations in the quality of light can influence student achievement and behaviour. Closely related to light is colour and there are studies which indicate that colour too can produce measurable effects in the classroom. The reported research is sufficiently conclusive to allow the suggestion that light and colour have predictable effects on students and that learning environments can be designed to foster these predictable effects.

Health and Welfare Canada (1975). Health and Built Environment. Health Facilities Design Directorate, Health and Welfare Canada; 1975.

This conference examined the evidence of ill health and other effects on human well-being resulting from interaction with the built environment. Society often militates against individuals adopting life-styles leading to good health. There is a need for a new type of cost-accounting which looks at long-term health effects of good environments. Environmental design should be for promotion of health, and although some risk-taking is healthy, people are not always in a position to evaluate environmental risk in that respect. Design for health may become a question of consciousness raising, increasing perception and broadening fields of concern. People should be given a feeling of competence in dealing with their designed environment. Also discussed are computer-related problems, self-help systems and alternatives to health care, research, education, professional training, facility design, risk, and standards.

Health and Welfare Canada (1976). Health Promotion Through Designed Environment: Conference Report. Health and Welfare Canada, Health Programs Branch; October, 1976.

The conference was convened to examine the evidence of ill health and other effects on people's well being resulting from interaction with the built environment; deal with what may be called the institutionalization and individualization of health; to discuss the promotion and maintenance of health; and to examine the application of design to the planning for health.

Health and Welfare Canada (1980). Workshop on Window Design: Considerations For Health and Well-Being. Health Services and Promotion Branch, Health Facilities Design Division, Department of National Health and Welfare, Ottawa Canada; March 1980.

This volume contains the proceedings, transcribed in discussion format, of a three-day meeting of window and building design experts from the United Kingdom, the United States, and Canada. It addresses the history and the state of the art of window design, the building code, biological/emotional/aesthetic considerations, and the disadvantages of windowless environments. It was concluded that evaluation of windowless environments, and especially deep-core buildings, is a research priority.

Health and Welfare Canada (1982). Canadian Governmental Report on Aging. Department of National Health and Welfare, Ottawa ON; June 1982.

The health of older persons affects, and is affected by, a number of factors including income security, employment opportunities, life with-

in the family and community, leisure activities, living arrangements, and social service needs. A holistic approach to health in old age is needed. The health care system should promote the highest possible functional independence and health among the aging, provide appropriate care in an integrated and coordinated way, ensure equitable and reasonable access to appropriate services, and provide for the needs of those among the aging who require special care. Suggestions include health promotion programs from an early age, increased safety awareness to prevent falls and other accidents, development and use of vaccines to prevent respiratory illness, physical and mental fitness promotion, improved psychiatric services, more intensive and extensive education about normal age-related processes, proper packaging and labelling of drugs and the monitoring of adverse drug reactions, and on-going evaluation and control of medical devices. There is a need for improved coordination in the planning, development, and implementation stages of policies and programs, and in the delivery of services. Integration of health and social services delivery needs to be improved country-wide. The uneven distribution of health services between provinces, as well as the larger urban centres and smaller urban and rural communities within provinces, also affects the health of the aged as do limits to access to services due to language, distance, cultural background, and so on.

Heath, Lormar (1981). *Gay Fathers: Some of their Stories, Experience, and Advice*. Gay Fathers of Toronto, PO Box 187, Stn F, Toronto M4Y 2L5; 1981.

The authors describe difficulties experienced by fathers who are gay, citing specific experiences involving marriage breakup, access to children, discrimination, and the process of coming out. For whatever reasons, many men who are homosexuals do marry or establish continuing relationships with women, and do father children. The life of the gay father is often fraught with the fears and anxiety that come from leading a double life. They experience the conflict between what society expects of them, and what they come to realize is a deep and unalterable part of themselves. They feel a need to express their identity but do not want to jeopardize their relationship with their children or damage them in any way. The authors attempt to contradict common stereotypes, and to reduce the isolation of gay fathers. Additional bibliography available from publisher.

Helsing, K.J.; Comstock, G.W.; Meyer, M.B.; Tockman, M.S. (1982). *Respiratory Effects of Household Exposures to Tobacco Smoke and Gas Cooking on Non-Smokers: from Indoor Air Pollution: Proceedings of the International Symposium on Indoor Air Pollution, Health and Energy Conservation*; Spengler, John— editor; pp. 365-370. *Environment International*, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

The records of 708 nonsmoking white adult residents of Washington Co, MD, who had participated in two studies of respiratory symptoms were

analyzed to evaluate the effects of exposure at home to two potential sources of indoor air pollution: cigarette smoking by other household members, and use of gas as a cooking fuel. After adjustment for the effects of age, sex, socioeconomic level, occupational exposure to dust, and years of residence in household, the presence of one or more smokers in the household was only suggestively associated with a higher frequency of chronic phlegm and impaired ventilatory function. The use of gas for cooking was associated with a significantly increased frequency of chronic cough and a significantly greater percentage with impaired ventilatory function.

Henderson, A.S.; Grayson, D.A.; Scott, R.; Wilson, J.; Richwood, D.; Kay, D.W.K. (1986). Social Support, Dementia and Depression Among the Elderly Living in the Hobart Community. *Psychological Medicine*, No. 16; pp 379 - 390; 1986.

In a community sample of the elderly in Hobart, Tasmania, cases of dementia and depression were ascertained by the Canberra Geriatric Mental State and the Mini Mental State Examination. Social relationships and support were examined by means of the Interview Schedule for Social Interaction. The elderly had fewer social relationships than younger adults, but were more content with what they did have. Elderly women had more affectional ties than elderly men. The presence of offspring in the same town increased the number of close ties and of social relationships, but was more important for men than for women. Persons with cognitive impairment or an established dementia reported that they had less social interaction than they would like. Depressed subjects reported having markedly less social interaction than the mentally healthy elderly, but did not complain that it was too little.

Henzi, H. (1984). Chronic Methanol Poisoning With The Clinical and PathologicAnatomical Features of Multiple Sclerosis. *Medical Hypotheses*; No. 18, pp. 63-75; 1984.

The details of two cases of chronic methanol poisoning are presented. Both patients initially developed clinical symptoms of multiple sclerosis: visual disturbances, intention tremor; reduced abdominal reflexes, impaired coordination and difficulties with walking. After the exposure to methanol had ceased, the multiple sclerosis symptoms persisted in one patient but disappeared gradually in patient 2. The results are discussed in connection with the theory that under certain circumstances multiple sclerosis itself is induced by formaldehyde stemming from the metabolism of methanol.

Hertzman, Clyde (1986). *The Health Context of Worklife Choices: A New Work Agenda for Canada*. Canadian Mental Health Association, Toronto Canada; 1986.

The relative average life expectancy rates of males and females widens around age 40 in response to gender differences in risk, including the differences between middle-aged men and women in their relationships to the world of work. There is, however, a lack of data and methodology to estimate the proportion of the differences in health status between men and women, and between socioeconomic groups, attributable to the physical, chemical, and biological hazards related to work. Evidence linking unemployment to poor mental health does exist and is stronger than that linking it to poor physical health.

Hess, C.T.; Weiffenback, C.V.; Norton, S.A. (1982). Variations of Airborne and Waterborne Radon-222 in Houses in Maine: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 59-66; 1982; Pergamon Press, Oxford and Toronto.

Concentrations of airborne radon ranging from 0.05 to 135 pCi/L were found in houses in Maine during the period October 1980 to May 1981. To investigate the association between elevated radon concentrations in well water and the indoor airborne radon concentrations, the radon in the water supplies of these houses was measured by liquid scintillation. Monitors of airborne radon, recording in intervals of 10 min for periods of 5 to 7 days, were used for dynamic studies in 18 houses, determining the component of airborne radon in bursts. House residents kept logs noting the time of major water uses. For some of the houses, ventilation rates ranging from 0.3 to 2 air changes per hour were determined. The component of airborne radon associated with water sources was found to vary inversely with ventilation rate and directly with waterborne radon concentration. According to the the data are pertinent to a study which revealed significant correlations between county averages, from the National Cancer Institute, or age-adjusted cancer mortality rates in Maine and average values of radon concentrations in water for the counties.

Hexter, Alfred C.; Goldsmith, John R. (1971). Carbon Monoxide: Association of Community Air Pollution with Mortality. Science, Vol. 172, pp. 265-267; April 16 1971.

Keywords: carbon monoxide; indoor air pollution; health hazards;

Hickey, Richard J.; Clelland, Richard C.; Bowers, Evelyn J.; Boyce, David E.; Alarie, Yves (1976). Health Effects of Atmospheric Sulfur Dioxide and Dietary Sulfites (and Rebuttal). Archives of Environmental Health, pp. 108-112; March/April 1976.

Some animal studies have shown that exposure to low levels of sulfur dioxide and dietary bisulfite is relatively harmless. To the contrary, sulfite oxidase deficiency is known in man, bisulfite is mutagenic for several test organisms, and the atmospheric sulfur dioxide level is

positively correlated with death rates for several chronic diseases among some human populations. The studies reporting harmlessness for animals may be misleading because of the fallacy of typology.

Hijazi, N.; Chai, R.; Bradstreet, J.W.; Duffee, R.A.; Astle, A.; Amster, M. (1983). Indoor Organic Contaminants in Energy Efficient Buildings. TRC Environmental Consultants, Inc., 800 Connecticut Blvd., East Hartford, CT, USA 06108, (203)-289-8631.

This private sector report describes the use of the TAGA gas-chromatograph mass-spectrometer (GCMS) equipment for measuring volatile contaminants in indoor or outdoor air. The discussion includes a listing of common chemicals found in trace amounts indoors.

Hildingson, Olov (1982). Radon Measurements in 12,000 Swedish Homes: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 67-70; 1982; Pergamon Press, Oxford and Toronto.

Radon daughter levels have been monitored in 12,000 Swedish homes during the last two years using two methods; a track-etch detector and a filter sampling technique with determination of ventilation rate. In 1979, the Swedish government introduced temporary limits for the radon daughter concentration in dwellings. For existing buildings, this limit is 400 Bq/m³ (0.11 WL). Close to 15% of the investigated houses have a radon daughter concentration higher than the limit. Almost 10% have a concentration about 1000 Bq/m³ (0.27 WL). The results from this study show that the two most important sources for radon in buildings are building materials and the ground.

Hirdes, John P.; Brown, K.S.; Vigoda, Debby S.; Forbes, W.F.; Crawford, Lawrence (1987). Health Effects of Cigarette Smoking: Data from the Ontario Longitudinal Study on Aging. Canadian Journal of Public Health, Vol. 78, pp 13 - 17; January/February, 1987.

The association between cigarette smoking and mortality is well established, but a similar association between cigarette smoking and morbidity is less well documented. This report describes an analysis of data obtained from the Ontario Longitudinal Study of Aging which represents a cohort of about 2,000 males of similar age, which have been followed over a 20 year period since 1959. Smokers were more likely to report poorer health and permanent disability. Drop-outs, because of death and lack of interest were significantly greater for smokers than for non-smokers.

Hirschman, Shalom Zarach; Feingold, Murray; Boylen, George (1963). Mercury in House Paint as a Cause of Acrodynia: Effect of Therapy with N-Acetyl-D,L-Penicillamine. The New England Journal of Medicine, Vol. 269, No. 17, pp. 889-893; Oct. 24 1963.

Keywords: mercury; building materials; air pollution; acrodynia;

Hoff, H. (1964). Mental Health Implications in the Peaceful Uses of Nuclear Energy. Social Implications of the Peaceful Uses of Nuclear Energy; UNESCO, pp 99 - 104; Klineberg, Otto, editor; 1964.

Emotional reactions to proposed and actual use of atomic energy for peaceful purposes include anxiety and fear — especially of automation, biological chain reactions, and poisoning of air, water and food. Such anxieties inevitably develop pathological defense mechanisms, such as denial, repression, and regression to infantile behaviour, that demand immediate gratification.

Holland, Jerry (1987). A Message From the President of One Voice. One Voice - La Voix, Vol. 1, No. 1, p 1; Ottawa Canada; July, 1987.

One Voice - The Canadian Seniors Network - is a national, non-profit organization which was founded in March, 1987 to help overcome the barriers that separate seniors from the government and to help governments understand seniors' perspectives on current issues, including health, housing, economic and other pressing social policy issues. It operates under the belief that Canadian's attitudes towards aging can be changed for the better. The newsletter, One Voice - La Voix, facilitates communication among individuals and groups

Holland, Walter W. (1974). Effects of Air Pollution on Children. Pediatrics, Vol. 53, No. 5, Part II, pp. 839-841; May 1974.

This article discusses the chronic and acute effects of exposure to high levels of air pollution, specifically impaired respiratory function.

Hollands, John (1984). Environmental Concerns in Offices and Homes: Conference Report. University of Toronto Staff Association Newsletter, Vol. XIV, No. VI, pp. 7-8; October 1984.

This is a report on a conference on Environmental Concerns in Offices and Homes, held at the University of Western Ontario's Occupational Health and Safety Resource Centre on September 24 and 25, 1984. Ministry of Labour representatives are reported as suggesting that the sick building syndrome symptoms of headaches, drowsiness, tired eyes and upper respiratory irritation, is caused by inadequate ventilation in modern air-tight buildings. VDT user fatigue, eye fatigue and radiation measurement studies were also discussed.

Hollenberg, C.H.; Siminovitch, L. (1986). Importance of National Centres of Research to Canadian Health Care and Medical Science. CMAJ, Vol. 135, pp 195 - 196; August, 1986.

A close relationship exists between the quality of a nation's health care services and the quality of its medical research programs. High-quality medical research endeavours can now be found in all Canadian medical schools and in some teaching hospitals. These efforts are based on the contributions of widely dispersed investigators usually working in small groups. While this approach to medical research has been successful in establishing the broad base of science required to support our educational institutions, it has become apparent that it is no longer adequate to meet all of the technologic and interdisciplinary demands of contemporary medical research. Many of the important research problems of our time can only be approached in a definitive manner by mobilizing the efforts of relatively large groups of scientists and providing them with often high-cost technology. Such concentrations are beyond the present capacities of universities, in spite of expanded facilities in Calgary, Toronto, London, and Winnipeg. The correction of the deficiencies will require a very significant change in the amount of money available for medical research and in the way the money is used. The federal government, acting in association with the private sector, could initiate a new program to create a group of major Canadian centres of research in proximity to those universities and health science centres that have developed a significant scientific base in relevant disciplines. Initiatives should be selected carefully, bearing mind the major thrusts of contemporary science, those diseases of great importance to Canadians in which a major research investment is likely to pay off and those research developments likely to spark commercial exploitation.

Hollowell, C.D.; Budnitz, R.J.; Traynor, G.W. (1976). Combustion-Generated Indoor Air Pollution. University of California, Lawrence Berkeley Laboratory, Energy and Environment Div., Technical Information Dept., Berkeley CA 94720; Dec. 1976.

Keywords: air pollution; heaters; carbon monoxide; nitrogen compounds; nitrogen oxides; sulfur compounds; sulfur oxides; health hazards; buildings; combustion products; cooking devices; space heaters;

Hollowell, Craig D.; Berk, James V.; Lin, Chin-I; Turiel, Isaac (1979). Indoor Air Quality in Energy-Efficient Buildings. University of California, Lawrence Berkeley Laboratory, Berkeley CA; March 1979.

This reference is one of a long series of competent articles containing experimental data regarding indoor air pollution, produced by the Lawrence Berkeley Laboratory of the University of California at Berkeley. The reader can obtain a listing by writing directly or by enquiring of the U.S. National Technical Information Service (NTIS), with which most of the papers are registered.

Hollowell, Craig; Berk, James V.; Traynor, Gregory, W. (1979). Impact of Reduced Infiltration and Ventilation on Indoor Air Quality in Residential Buildings. ASHRAE Transactions, Vol. 85, Pt 1, pp. 816-827; 1979.

Keywords: buildings; ventilation; energy conservation; air pollution; health care; nitrogen compounds; formaldehyde; indoor air quality;

Hollowell, C.D.; Berk, J.V.; Boegel, M.L.; Miksch, R.R.; Nazaroff, W.W.; Traynor, G.W. (1980). Building Ventilation and Indoor Air Quality. University of California, Lawrence Berkeley Laboratory, Energy & Environment Div., Berkeley CA 94720; January 1980.

Keywords: ventilation; indoor air pollution; carbon monoxide; nitrogen dioxide; formaldehyde; radon; energy efficient building;

Hollowell, Craig D.; Miksch, Robert R. (1981). Sources and Concentrations of Organic Compounds in Indoor Environments. Bulletin of the N.Y. Academy of Medicine, Vol. 57, No. 10, pp. 962-977; December 1981.

Keywords: human; building material; smoking; formaldehyde; indoor air pollution; health hazards; nitrogen dioxide; safety standards;

Holma, B.; Kjaer, G.; Stokholm, J. (1979). Air Pollution, Hygiene and Health of Danish Schoolchildren. The Science of the Total Environment, Vol. 12, pp. 251-286; Elsevier Scientific Publishing Co., Amsterdam; 1979.

A comprehensive investigation forming part of a joint European study under the auspices of "WHO Long-Term Air Pollution Programme" uses 7-13 year old school children as the target group. The study has included social, housing, hygienic and epidemic factors as well as family smoking habits. The results indicate that, at exposure to low levels of air pollution, these factors dominate as causes for the impairment of health, especially that of respiratory health.

Holt, John (1987). Watching Children Learn: Correcting Mistakes. Growing Without Schooling, No. 57, p 9; 1987.

Intellectual learning is facilitated best by allowing children to figure things out on their own because it fosters remembrance and confidence. Rather than correction of errors, we should provide them with acceptance and recognition at having performed an intellectual feat. Children learn as they breathe — it is in their very nature to take the world in with their senses, and to make sense of it, without being aware of how or why they are doing it. A great error is made when, by correcting them, we make them self-conscious of their learning, so that they begin to ask themselves if they are learning or not and so, through worry, turn off

their learning. When people see the world as a place of danger, from which they must protect themselves — in short, when they begin to live less freely and fully — that is when learning dies down.

Hooker, C.A. (1982). The Human Context for Science and Technology: Final Report. Social Sciences and Humanities Research Council of Canada; March, 1982.

As an area of study, the Human Context for Science and Technology will demand an integrated approach transcending disciplinary and institutional boundaries. It will also involve a fundamental re-integration of human values into the inquiry process. Existing Canadian research activity in this area is inadequate because it is insufficient, uneven, narrow, and socially incomplete and unrepresentative. All too often, many groups and constituencies like women, industrial workers, and those whose jobs may be eliminated by the spread of microelectronics, have legitimate and pressing concerns but are excluded from access to the resources needed and from opportunities to articulate their perspective and develop their own analysis of the issues.

Hopkirk, Gerald (1986). Mental Fitness in the Workplace: Worklife Choices for Professionals in a Medium Size School System. Work and Well-Being Quarterly, Issue 1; pp 8 - 11; Canadian Mental Health Association, Toronto ON; Fall, 1986.

This paper describes a school system, its philosophy of administration, and the worklife choices available to its 600 professional staff members. Described are policies for leaves of absence, job sharing, flexible part-time work, and transfers. The employer operates under the assumption that the provision of meaningful worklife choices for employees is advantageous to employer and employee alike and is well worth the resources and energy spent on it. Helping people find work situations that allow them to reach personal goals they define for themselves make the organization a dynamic, fulfilling place to work and creates the type of climate conducive to effective performance.

Hosein, H. Roland; Corey, Paul (1986). Domestic Air Pollution and Respiratory Function in a Group of Housewives. Canadian Journal of Public Health, Vol. 77, pp 44 - 50; January/February, 1986.

Air pollution exposure was determined to sulfur dioxide, nitrogen dioxide, and respirable suspended particulates for a group of symptomatic housewives and a matched control group using a specially designed domestic air pollution sampler. The nitrogen dioxide and respirable suspended particle levels were higher indoors than outdoors, whereas the sulfur dioxide levels were higher outdoors. The cases and controls showed the same air pollution exposures. The use of gas stoves resulted in

elevated levels of nitrogen dioxide which might account for the observed reduction in lung function of residents in homes with gas stoves. Cigarette smoking indoors resulted in elevated particulate levels.

Huggins, Hal. A. (1982). Mercury: A Factor in Mental Disease?. Journal of Orthomolecular Psychiatry, Vol. 11, No. 1, pp 3 - 16; 1982.

Many ramifications of physical and mental health need investigation in light of the potentially hazardous effects of mercury toxicity. Suggestions have been forwarded that mercury leaching out of dental amalgam filling can affect the peripheral nervous system, immune system, and cardiovascular system, and that mercury in a biological system appears to create or mimic many disorders in these three areas. The case of a young girl has been presented which has been thoroughly investigated through hospital and direct evaluation by psychiatrists, internists, osteopaths, chiropractors, psychologists, and clergy. She was put into a mental institution based on a vast array of physical and emotional problems. None of these problems were present before her fillings were placed, or after they were removed. Because it is common to refer patients like her with multifocal problems defying diagnosis to a psychiatrist or psychologist, it is important that professionals in these fields become more aware of the possibility of mercury toxicity.

Hume Hall, Ross (1987). Why The EPA Won't Work. Probe Post, Volume 10, No. 1, pp 29 - 31; Spring, 1987.

The author is of the opinion that the 1987 Environmental Protection Act (EPA) will be ineffectual. He feels that the Act is flawed because it is preoccupied with an outmoded, single chemical approach to dealing with toxic chemicals which focusses on deciding on each one's level of toxicity rather than the design of new policies to keep all chemicals out of the environment. This approach results in a very cumbersome, slow process which, he feels, does not keep up with the pace of technology. A second difficulty which Hall feels dooms the EPA is Environment Canada's relatively low position in the hierarchy of importance of government departments, a status which necessitated its liaison with a relatively stronger department, Health and Welfare, in order to pass the EPA. He claims that Health and Welfare lacks the motivation, expertise, and incentive to explore the broad questions of harmful effects of environmental toxicity. The initiative for a comprehensive approach to environmental cleanup and protection has to come from nongovernmental groups and provincial and municipal agencies, he feels.

Humiston, Karl (1987). Letter from Karl E. Humiston, M.D. Personal Correspondence, Karl H. Humiston, New York, NY; July 6, 1987..

The author, a physician practising environmental medicine for the last decade, notes that he has seen patients decline in health while

practising avoidance so strict that it occupied their whole energy and attention. He concludes from personal experience that high-energy places, people and activities can strengthen one's immune system, and found that if he and his wife are reasonably attentive to environmental factors and select what they involve themselves in, New York is a healthy place for them to live. He now advises ecology patients that seeking positive influences is just as important as avoiding negative ones. This includes good family and spiritual life, food, clothing, housing, reading, music art, and socializing. Money makes a difference, but is not as important as the spirit of seeking. He instructs patients not to let ecological illness stop them from living their lives.

Hutchinson, Peggy; Lord, John; Osborne, Way, Lynn (1986). Participating With People Who Have Directly Experienced the Mental Health System. Canadian Mental Health Association; November, 1986.

The hopes of people disabled by mental health problems are not unlike those of other Canadians. Their use of formal mental health services does not mean that other things in life, like making a contribution to family, friends, work and neighbourhood, are any less important. In essence, participation is about involving and supporting individuals with a disability to be part of the everyday things which many of us take for granted. To achieve the quality of life which we all desire, the lives of individuals who have experienced the mental health system must be more fully experienced in the community. Only as participants in this sphere of human activity will they achieve real citizenship status. This report deals with the issues arising from community participation of the mentally disabled, presenting barriers and illustrations from experiences across the country.

Hutchison, Peggy; Lord, John; Savage, Harvey; Schnarr, Anne (1985). Listening to the People Who Have Directly Experienced the Mental Health System: Building a Framework for Support. Canadian Mental Health Association; August, 1985.

This study presents a consumer perspective on the mental health system using the words and stories of those who have experienced it. Feelings of cynicism and powerlessness result from the overwhelming paternalism directed towards these clients of the system. Most of those interviewed expressed deep concern that traditional psychiatric labels make it almost impossible for them to be seen as human beings. These people express a desire for independence but also require a great deal of support to become re-integrated into community life. They may periodically need some form of shelter or refuge but they believe the imposing controls and practices of institutions and hospitals are generally an inappropriate intervention. The rehabilitation and welfare systems tend to trap many into a life of poverty and dependence which result in feelings of low self-worth. Lack of appropriate housing conditions and work opportunities contribute to lack of dignity, income, and security.

Hutchison, P.A. (1986). Health Care for Canadian Native People. Can Med Assoc J, Vol. 134, pp 305, 308; February, 1986.

The Regina Health Department has as permanent staff two native community health care workers who had initially trained under the Medical Services Branch of the Department of National Health and Welfare in addition to two other native health workers. The program's aim is to hire native health care workers, to motivate native people to avail themselves of the department's programs and to adopt healthy lifestyles, and to facilitate liaison with the native community and organizations. Evaluation is difficult, but some native people with health problems have been reached, there have positive results from immunization, and medical problems have been recognized and corrected.

Ibsen, Karsten Kaas; Valbjorn, Ole; Nielsen, Aksel (1981). The Indoor Environment and Symptoms of Disease. Assessment of the Conditions in Schools in Copenhagen. Ugeskrift For Laeger, No. 143, pp. 1919-1923; July 1981.

A comparison was made between the incidence of disease and symptoms and the design of schools, with particular attention to the age of the buildings, flooring and ventilation systems. The registration of conditions of health in 14,561 pupils in 88 schools made by the school health services and a questionnaire involving 1226 pupils in 14 schools in Copenhagen were used. Significantly more numerous cases of allergy (asthma and hay-fever) were found in schools with textiles as floor coverings and significantly more numerous cases of skin conditions in the schools which were less than five years old or which had been renovated within the past five years. 31% of the pupils and 40% of the teaching staff stated that they took headache medicine at least once per month. Under one third of these pupils and over half of these teachers took headache medicine at least once weekly. By comparison between data obtained in the indoor environmental investigation and the registrations made by school health services, it was found that pupils stated higher incidences than those registered by the health services people. More than one third of the pupils participating in the indoor environment investigation had complaints which they attributed to the indoor environment. The majority had complaints concerning heat and noise. In schools with textile floor coverings, static electricity was the most widespread inconvenience as 60% complained about this as compared with 5% in the remaining schools.

Illich, Ivan (1976). The Age of Disabling Professions. Health Promotion Through Designed Environments, pp 5 - 26; Health and Welfare Canada, Health Programs Branch; October, 1976.

A radical departure is necessary from a world of disabling professions to a world in which health is deinstitutionalized and personalized in such a manner that the individual may begin to assume personal responsibility for his or her health, with little or no assistance from the

health institutions or the various health professions. Specialist power to define issues in terms of problems creates problems. Medicine tends to engender helplessness and disease simply by taking out of the environment the opportunities for coping which constitute health. Our major institutions have acquired the uncanny power to remove society consistently from those same purposes for which they originally had been engineered and financed. Thousands of small groups are at present challenging professional dominance over themselves and over the socio-technical conditions in which they live.

Imbus, Harold R. (1982). Acute Effects of Exposure to Formaldehyde. Presented at the Conference: "Formaldehyde..the Facts", Toronto Canada, May 1 1982.

Formaldehyde can result whenever there is combustion and enters the air from auto exhaust, fuel burning, photooxidation reactions, cigarette smoking. Consumer products that also contain formaldehyde are fabrics and paper products as well as a wide range of building materials. It is an irritant and a sensitizer, the latter meaning that it is capable of producing allergic sensitization - a specific immune response in an individual who becomes allergic to a substance. A number of organs can be affected by formaldehyde, primarily the skin, eye, upper and lower respiratory tracts, and the gastrointestinal system.

Ingraham, Leon (1983). Electromagnetic Radiation and Student Off-Task Behavior. Alberta Education, Planning Services Branch; July 1983.

The effects of electromagnetic radiation emitted from fluorescent lights on the off-task behaviours of grade three school pupils was studied. The independent variable was the level of electromagnetic radiation which was eliminated by grounding and shielding the fluorescent light fixtures in the experimental classroom. The off-task behaviours were recorded by a reliable on-sight observer for two groups of pupils. For the intact classroom group) comprised of all pupils and heterogenous with respect to hyperactivity) the elimination of electromagnetic radiation decreased significantly the rate of off-task behaviours. For triad groups (groups of three pupils selected as being most "hyperactive) the results were mixed and inconclusive. Contrary to expectations, pupils from this latter group who were prone to hyperactivity, demonstrated no benefit from the elimination of electromagnetic radiation. Further studies are suggested.

Innes de Neufville, Judith (1981). Social Indicators of Basic Human Needs: Quantitative Data for Human Rights Policy. Institute of Urban and Regional Development, University of California, Berkeley CA; May, 1981.

Among the internationally recognized human rights are a series which deal with basic human needs — education, health, nutrition and income, and the right to an adequate standard of living. Discrimination against women in these areas is also of prime importance. U.S. human rights

policy has not generally taken these into account. If international policies are to encourage, assist, or pressure nations to achieve basic human needs, some definition and measurement of existing levels of need is essential. These social indicators are required if human rights policy on basic needs is to be implemented.

Innes de Neufville, Judith (1985). *Knowledge and Action: Making the Link*. Institute of Urban and Regional Development, University of California, Berkeley CA; November, 1985.

Central to the idea of planning is the systematic use of information to help shape decisions. In practice, however, though data are plentiful, the lament is all too familiar that a decision was made with inadequate information. This paper argues that much of the problem lies in the fact that explicit prescriptions about how to link knowledge to action are hampered by implicit assumptions about the nature of knowledge. These assumptions place a distance between knowledge providers and users and define knowledge as consisting of abstract principles and measurable facts. Knowledge could, instead, be defined as including a recognizable social reality and assumes researchers cannot be simply detached observers. Planners must embrace rather than deny ambiguity and uncertainty. Information can reduce uncertainty, but not impose certainty. Public debate over data can provide one acceptable way to decide how to handle ambiguity and risk.

Insel, Paul M. (1983a). *Environmental Variables and the Prevention of Mental Illness*. Lexington Books, 1983.

Mental health services cannot adequately respond to the needs of citizens unless those involved in the planning, organization, and delivery fully recognize the harmful effect that a variety of social, environmental, physical, psychological and biological factors can have. This book provides a thorough summary of the literature about environmental variables and their effects on behaviour. Extensive bibliographic references are included.

Insel, Paul M. (1983b). *Social Climate of Mental Health*. *Environmental Variables and the Prevention of Mental Illness*, chapter 2, pp 9 - 26; Lexington Books; 1983.

From the social-ecological perspective, mental health is intimately linked to the social environment so that in order to prevent mental illness, one must understand and assess the social environment. The probability of achieving an optimum environment is increased when planning of the environment is conducted by those who must function within the environment. Since it is the perceived climate that has the most important effect, perceptions appear to be more valuable than objective

assessments of an environment by outside planners. Involvement in planning one's own environment is vital to self-esteem and good mental health. Many investigators have suggested that humans have an active need for control over their own environment. Participatory planning and the use of systematic assessment can fulfill this human need while developing competence in individuals to change and control their own environments.

Irwin, John (1987). The Liberation of Males. Presented at 1987 International Men's Leaders' Conference, Hebron, Connecticut, USA; October 23-25, 1987.

The author describes in detail how male stereotyping has led men to take an oppressor role against women (sexism), forcing women into unhealthy situations. The dehumanization of men, particularly being treated as an expendable economic and military resource, has also led to unhealthy situations for men in our society. The threat of being called 'gay' or 'not manly' is used, from an early age, to enforce male stereotyping (e.g. to be strong, aggressive, violent, unfeeling, sexually compulsive, oppressive to others, and less able to establish close relationships than women). The author sees the elimination of male stereotyping as a process which must proceed in parallel with the elimination of sexist behaviour (oppression of women by men), for either process to be successful.

Ishikawa, Satoshi; Miyata, Mikio; Okuwaki, Kenichi; Namba, Tatsuto; Fukushima, Kazuya (1986). Analysis of HLA Antigens in Behcet's Disease - A Possible Implication of Environmental Chemicals. Clinical Ecology, Vol. IV, No. 2, pp. 81-87; 1986.

Environmental chemical exposure was postulated as a possible cause of Behcet's disease, a mucocutaneous-ocular syndrome leading to blindness, ulcers, and sometimes death. Most of the patients studied had a history of contamination with organochlorine compounds and/or organophosphorus pesticides. The number of patients in Japan with this disease declined following the banning of certain strong toxic chemicals, including DDT, BHC (benzene hexachloride) and parathion, with some time lag. The researchers postulated that Behcet's disease may be caused by environmental chemicals, which may cause supersensitivity of the tissues, possibly leading to abnormality in the immune system.

Israelson, David (1987). Federal Department Fails to Back Report on Lakes Pollution. Toronto Star, December 1, 1987.

Federal researchers Tom Muir and Anne Sudar are reported to have authored a report which states there is strong evidence to link Great Lakes pollution with health problems such as cancer and birth defects. Cancer rates and rates of birth defects were found to increase from west to east along the lakes, and are heaviest near highly-contaminated sections such as the Niagara and St. Lawrence Rivers. People living along

the lakes suffer higher rates of heart disease and strokes than other Canadians. The article also reported that the Federal Government will begin a followup in early 1988, and report on health effects of Great Lakes pollution in 1989.

Iyer, E.M.; Dikshit, M.B.; Suryanarayana, S. (1985). Effect of Exposure to Heat, Hypoxia, Cold, Acceleration, and Vibration Stress on the Total Blood Sulfhydryl Groups in Human Subjects. *Aviation, Space, and Environmental Medicine*, pp 1097 - 1101; November, 1985.

The percent utilization of total blood sulfhydryl (bl-SH) group was assessed in 100 healthy, unacclimated male Indian Air Force personnel who participated in various stress trials which included heat, cold pressor test, hypoxia, 70 degree head-up tilt, acceleration, and vibration. Exposure to a hot and dry environment for a period of 50 minutes showed 49.7% fall, while exposure to a hot and wet environment showed a greater fall of 61%. Hand immersion for 2 minutes in water at 4 degrees C showed a 42.7% fall. Acceleration stress showed a highly significant fall and low frequency sinusoidal vibration produced a 42.7% rise.

Izumi, K. (1968). *Psycho-Social Considerations of Environmental Design*. National Society of Interior Designers, Interior Environment Research Council, 315 East 62nd St., New York NY 10021; May 1968.

Keywords: environmental design;

JML (1987). The Ethics of Paternalism in Public Health. *Canadian Journal of Public Health*, Vol. 78, Jan./Feb. 1987; pp 3 - 4.

Those who practice public health believe implicitly that they know best what is in the public interest. This is an attitude which others might describe as paternalistic or gratuitous interference in other people's lives. Situations such as occur when parents' religious or philosophic beliefs prevent their children from having blood transfusions or being immunized are discussed. Efforts to control tobacco smoking and provide sex education to children also create awkward ethical questions. There should be a formal debate about these issues.

Jacko, M.G.; Ducharme, R.T. (1973). *Brake Emissions: Emission Measurements from Brake & Clutch Linings From Selected Mobile Sources: Final Report*. Environmental Protection Agency, Office of Air and Water Programs, Ann Arbor MI 48105; March 1973.

In order to define the extent of gas and particulate emissions from automotive brakes and clutches, a combination separation and storage collection system was devised. Unique emissions collectors for both disc and drum brakes and for a clutch were designed and built as the main

embodiment of this instrumentation. The hardware was installed on a vehicle which was then driven through various test cycles to determine extent and type of brake emissions generated at low and high operating temperatures. The particulates were processed and analyzed by a combination of optical and electron microscopy to ascertain the asbestos content and the particle size distribution. On the average, more than 99.7 percent of the asbestos is converted; the contribution to the atmosphere is 5060 pounds, or 3.2 percent of the total asbestos emissions.

Jaeger, R.J. (1981). Carbon Monoxide in Houses and Vehicles. Bull. N.Y. Acad. Med., Vol. 57, No. 10, pp. 860-872; December 1981.

Carbon monoxide is a common indoor air pollutant whose major toxic action from a public health standpoint appears to be associated with very subtle changes in neonatal growth, learning, and activity when low to moderate concentrations are tested. Such concentrations might occur after cigarette smoking or from living in poorly ventilated homes that use gas stoves for heating. Utilization of unvented internal combustion engines indoors should be discouraged, as should smoking. Carbon monoxide levels in blood range from less than 1% in nonsmokers to 5 to 10% in smokers. In fetuses of smoking mothers, as much as 7.6% of CO-HB was reported.

Jaffe, Louis S. (1967). The Biological Effects of Ozone on Man and Animals. American Industrial Hygiene Assoc. Journal, pp. 267-277; May-June 1967.

Keywords: ozone; health effects;

Jaffe, Peter; Wolfe, David A.; Wilson, Susan; Zak, Lydia (1986). Emotional and Physical Health Problems of Battered Women. Can. J. Psychiatry, Vol. 31, October 1986; pp 625-629.

The study focused on the emotional and physical health problems of battered women by comparing a sample of residents in shelters with a group of women in the community matched for family income, length of marriage, and number of children on the General Health Questionnaire. The results indicated that battered women report a significantly higher level of somatic complaints, anxiety, and depression. These effects tended to be associated with other life stressors and children with serious behaviour problems. The findings suggest that battered women represent a population that is at an elevated risk of developing pronounced mental health problems. The implications of the study are discussed in terms of assessing the needs of battered women and their children as well as being vigilant for family violence as an etiological factor for other problems. Future research with larger samples of women and children, as well as longitudinal studies to investigate changes in adult and child adjustment over time, is necessary to provide data that goes beyond these findings.

Jammet, H.; Bosnjakovic, B.F.M.; Czerski, P.; (1985). Occupational Hazards From Non-Ionizing Radiation. Occupational Safety and Health Series No. 53, International Labour Office, Geneva, Switzerland;.

This review provides an overview of the non-ionizing radiation field, including biological effects from extremely low-frequency fields (0-300 Hz). The Committee concluded that no specific pathology was found to be associated with exposure to electric fields around transmission lines, and that no significant health risks were identified on the basis of epidemiological studies of workers occupationally exposed to extremely low frequency radiation and power frequencies (50 Hz and 60 Hz).

They note that a low-frequency electric field does not penetrate the human body to any significant extent, in contrast to a magnetic field, and that there are no known specific symptoms for ELF electromagnetic field effects. With strong alternating electric fields three interactions occur:

- a) Small currents flow within the body due to the capacitive coupling to the field.
- b) Spark discharges occur when objects of significantly different potential are contacted.
- c) Large surface fields occur, particularly at sharply curved surfaces, and these local fields may produce superficial sensations of the field.

The magnitude of the internal currents is small in comparison with that of the currents that flow when contact is made with charged conductors. The internal electric field associated with the small currents is approximately one million times smaller than the applied external electric field. Magnetic ELF fields induce eddy currents in the body, that may produce heating. A magnetic field corresponding to a 10 kV/m electric field could induce a voltage of 1 mV in humans. The authors note that this is a small value with respect to the voltages needed to cause biological effects, and that only with field intensities of some tens of milli-Teslas (about ten thousand times that associated with power lines) have specific effects been reported.

There is some evidence that the nervous system may be affected by ELF electric fields that are far too weak to have a direct effect on brain synaptic function or membrane excitability through field influence on ionic motion through the cell membrane. Animal studies have also shown effects on calcium exchange from various brain tissues. The authors conclude, however, that none of the evidence indicates that even strong electric fields have effects that 'compromise man's ability to function or have long-lasting or permanent effects on neuro-physiological health.'

Jarvis, Bruce B. (1986). Potential Indoor Air Pollution Problems Associated with Macrocytic Trichothecene-Producing Fungi. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Four genera of fungi have been reported to be producers of macrocytic trichothecene toxins. A case study is presented which describes a family living in a brick house in Chicago. They had been, over the course of a 5 year period, subject to a variety of recurring maladies including cold and flu symptoms, sore throats, diarrhea, headaches, fatigue, dermatitis, and generalized malaise. The father experienced severe leg pains and the family was highly stressed. Air sampling indicated the presence of numerous fungal spores. This, and the family's symptoms, indicated testing for trichothecenes; testing proved positive. Building materials of high cellulose and low nitrogen content that become moist and are subjected to temperature fluctuations can provide ideal growth conditions for fungi production, specifically *S. atra*. The symptoms exhibited by the occupants of the Chicago house are consistent with those reported to result from macrocytic trichothecene toxins. The trichothecenes are also immunosuppressive agents, a role that is often overlooked.

Johnson, Byron (1983). Evacuation Techniques for Disabled Persons. National Research Council of Canada; March, 1983.

This is a study of techniques of preparing, lifting, and carrying persons with various disabilities in emergency evacuations of buildings. It is difficult for untrained rescuers to motivate certain people to evacuate. The tests with semi-ambulant persons showed that those who need assistance in a fire or other emergency were able to judge their needs quite well and may move quicker when aided, although persons prone to spasticity can move more surely with assistance. Without special planning, it is unlikely that non-ambulant disabled persons can participate in a total building evacuation.

Johnson, Kirk A.; Batts Young, Bambi (1982). Environmental Agents and Behavioral Aberration: A Neglected Public Health Dilemma. Center for Science in the Public Interest, Environment and Behavior Program, Washington DC 20009.

This paper gives a brief review of some of the pesticides, heavy metals, pollutants, foods, food additives, and drugs that have been linked with unexpected behavioral disturbances, including paranoia, hyperactivity, anxiety, and hallucinations. Unfortunately, high rates of misdiagnosis indicate that many health professionals are unaware that behavioral aberrations in a patient might be caused by exposure to such environmental factors. Many health professionals tend to attribute patients' complaints to mental illness, stress, or other non-physical agents. The authors find a clear need for more controlled, double-blind studies

of behavioral toxins. Federal regulatory agencies could provide a major impetus for such research by requiring that new and existing chemicals be tested for behavioral effects, and by considering behavioral effects when setting health and safety standards.

Johnson-Lussenburg, C.M. (1986). Viruses in Air and Their Potential for Infectivity as well as the Presence and Importance of Endotoxins and Legionella in Indoor Air: prepared December, 1985. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Certain viral infections may be transmitted indoors by the airborne route. The relative importance of airborne transmission in the spread of infections in comparison to other routes depends on the nature of the virus, the susceptibility of the host and the influence of the indoor environment. Although increased respiratory infections have been correlated with exposure to certain pollutants, the effect of those pollutants on the host susceptibility is difficult to evaluate. It is also difficult to identify the etiological agents involved in these infections and to determine whether the severity of the infection is a property of the infecting agent or a consequence of increased susceptibility of the host. An extensive bibliography is included.

Johnston, Anne (1983). Hazardous Substances & the Right to Know: The Public Health Perspective. Board of Health for the City of Toronto; October 1983.

This presentation discusses the various immediate and long-term health problems resulting from both low and high level exposures to a variety of toxic substances. It is pointed out that it is essential that hazards be anticipated and averted and that exposure be reduced or eliminated. We can best protect ourselves if we know where the toxic substances are located, therefore so-called "right-to-know" legislation is imperative.

Jonassen, Niels (1981). Radon Exhaling Properties of Building Materials. Presented at the 1981/2nd/European Conference on Building Materials, April 1981, Glasgow Scotland.

Keywords: radon; health hazards; building materials;

Kagawa, Jun; Toyama, Toshio (1975). Photochemical Air Pollution: Its Effects on Respiratory Function of Elementary School Children. Archives of Environmental Health, Vol. 30, pp. 117-122; March 1975.

The effects of photochemical air pollution on respiratory function of Tokyo elementary school children were investigated. Nine types of environmental factors were continuously recorded. Seven categories of respiratory function tests were performed on 20 normal 11-year old children

once a week from June to December 1972, as a general rule. The correlation coefficients between respiratory function measurements and each of the environmental factors were calculated. The maximum expiratory flow rate (V max) showed high correlation with the largest number of environmental factors. Among environmental factors, temperature highly affected various respiratory function tests. The ozone was significantly associated with airway resistance (Raw) or specific airway conductance (Gaw/Vtg), NO or NO₂ with V max, and temperature with Raw, Gaw/Vtg, and V max. Two subjects among all subjects were considered as the reactors to the environmental factors.

Kagawa, J.; Tsuru, K.; Doi, T.; Tsunoda, T.; Toyama, T.; Nakaza, M. (1980). Lung Function Studies on Intermittently Exercising High School Students Exposed to Air Pollution. Nitrogen Oxides and Their Effects on Health; pp. 333-342; Lee, S.D., editor; Ann Arbor Science Publishers Inc.; 1980.

The acute effects of air pollution on the lung function of 9 high school students with two hours intermittent exercise were investigated during 2 consecutive 5-day periods. Multiple regression analysis was performed for each subject, utilizing each lung function test as the dependent variable and each environmental factor as the independent variable. The evaluation of the square of the multiple correlation coefficient suggests that the significant contributions of the environmental factors to the total variance of lung function were seen in about half the subjects.

Kahn, Robert; Hein, Karen; House, James; Kasl, Stanislav; McLean, Alan (1983). Report on Stress in Organizational Settings: Analysis and Implications of Research/A Study by the Institute of Medicine, National Academy of Sciences. Stress and Human Health; chapter 5, pp 81; Elliott, Glen R.; Eisdorfer, Carl ed; Springer Publishing Company, New York; 1983.

Stress associated with organizational settings, such as work or school, constitutes a major part of the total stress experienced in people's lives. Organizational settings also are likely sources of stress because they provide the main context in which society makes demands on people to perform and to relate to a broad range of others in specified ways. But because organizations are ready-made mechanisms of social influences, power, and communication, they can and should be utilized in any large-scale effort to reduce stress or improve health.

Kailin, Eloise W.; Brooks, Clifton R. (1963). Systemic Toxic Reactions to Soft Plastic Food Containers. Medical Annals of the District of Columbia; Vol. XXXII, No. 1, pp 1 - 8; January, 1963.

Three patients with intolerance to multiple chemical factors in the environment are described. Double-blind testing established beyond reasonable doubt that certain persons react systemically and adversely to some factor which passes into food or water from ordinary polyethylene food containers.

Kailin, Eloise W.; Hastings, Alicia (1966). Cerebral Disturbances from Small Amounts of DDT: A Controlled Study. Medical Annals of the District of Columbia, Vol. 35, No. 10, pp 519 - 524; October, 1966.

Crystalline DDT dissolved in food oil at from 10 to 10,000 ppm when held 1 to 2 inches under the nostrils of highly sensitive subjects elicited in a few minutes visual disturbances, headache, weakness, perceptual abnormalities, and slowed mental and motor activities.

Kalnins, R.; and Gaudert, P.C. (1985). Formaldehyde Emissions from Typical ParticleBoard Applications and Assessment of Specific Abatement Measures: Sponsored by Air Pollution Control Association TT-7 Indoor Air Quality Committee, Health and Welfare Canada, National Research Council of Canada, Public Works Canada, Consumer and Corporate Affairs Canada, and Canada Mortgage and Housing Corporation. Presented at the International Speciality Conference on Indoor Air Quality in Cold Climates: Hazards and Abatement Measures; Ottawa, Ontario, April 29, 30 and May 1, 1985.

Formaldehyde emissions emanating from building materials made with formaldehyde-based resin binders have caused much concern to health authorities, government regulatory agencies, and affected homeowners. In particular, certain urea formaldehyde (UF) resin-bonded particle boards tend to have high emission potential because of the inherent hydrolytical instability of such resins. In modern home construction such board is used extensively in the built-in furniture and shelving in the kitchen, bathrooms, and closets. In some cases UF-bonded particleboard is also applied in the floor construction. Additional amounts of such board are introduced into the finished dwellings by furniture. A preliminary investigation in a senior citizens apartment indicated that installed particleboard was the major cause of the unacceptable formaldehyde levels. Concentrations in the air ranged from 0.2 to 0.8 ppm in the upper floor apartments, built with particleboard floor underlay. The present recommended limit is 0.1 ppm.

Kalymun, Mary; Combs, E. Radene; Hanzal-Kashi, Amy; Wilde, Vicki L.; Glunt, Eric K.; Welch, Polly; Robinson, John Deschamps (1985). Person/Environment Relationships and the Well-Being of Older Adults. Environmental Change/Social Change; EDRA 16; pp 354; Environmental Design Research Association; 1985.

As personal resources diminish, as often happens with aging, it appears critical that individual needs and environmental characteristics complement one another in order to maintain independence and personal control in one's life. One ecological perspective suggests that the goodness of fit that exists between the individual needs and preferences of older adults and environmental characteristics is preliminary to well-being. Both constituents must be considered simultaneously, as parallels, in examining the congruence between older adults and their residential settings. If environmental characteristics in residential settings

restrict behavior, elderly people are less likely to exercise control and independence. This situation is exacerbated by the limitations of space, physical design, and options to choose and maintain living arrangements that meet personal needs.

Kaminoff, Robert D.; Proshansky, Harold M. (1981). Stress as a Consequence of the Urban Built Environment. to appear in L. Goldberger & S. Breznitz, *Handbook of Stress*; New York: Free Press-Macmillan (In press, 1981).

Stress is defined as that pattern of psychological, behavioral, and physiological responses of the individual to demands of the physical and social environment that exceed his capacity to cope effectively, that is, carry out activities, realize goals, and experience satisfactions. Lack of fit between the properties of the physical environment and the requirements of the person may induce stress in that person by creating demands that exceed his or her ability to cope and still pursue other goals in the setting. To the extent of this stress, the individual may be motivated to reduce the discrepancy between negatively perceived aspects of the environment and personal requirements, either by adaptation to the environment, or exertion of control over it. In the context of an urban existence in a built environment, the nature and intensity of environmental stimulation including its predictability and the extent to which it can be controlled assumes considerable importance. A literature review suggests many relationships between physical features of the built environment and stressful outcomes for inhabitants.

Kasuga, Hitoshi; Hasebe, Akihisa; Osaka, Fumio; Matsuki, Hideaki (1979). Respiratory Symptoms in School Children and the Role of Passive Smoking. *Tokai J. Exp. Clin. Med.*, Vol. 4, No. 2, pp. 101-114; 1979.

A study of respiratory symptoms in 1,937 school children aged 6-11 years was carried out in Suginami, Tokyo and the response rate was 99.5%. The prevalence rate for symptoms was associated with their families' smoking habits and their residential conditions; the prevalence rate was the highest among heavy-smoker families living in the area along a main highway (within 50 meters), and the lowest in non-smoker families regardless of their residential areas. Furthermore, if they lived in an area well away from a main highway (over 100 meters), the prevalence rate was also the lowest regardless of the families' smoking habits. Relative risk in the highest group was 4.0. Previous studies have already suggested that there was a relationship between symptoms and passive smoking while some have insisted that there was no relationship between them. This study may be useful in solving such contradictory problems and in establishing the role of passive smoking in respiratory symptoms.

Kato, Terutaka; Kasuya, Minoru; Kagamimori, Sadanobu; Kozuka, Hiroshi; Hashimoto, Takejiro; Kawano, Shoichi (1981). Evaluation of Aerial Environmental Pollution by Means of Indicator Plants in a Slightly Polluted Area--Relationship Between the Vigor of Japanese Cedars and Prevalence of Respiratory Symptoms in School Children. *Journal of Pharmacobio-dynamics*, Vol. 4, No. 5, p. S-60; 1981.

The effects of low grade air pollution on human health and vegetation were studied in a Japanese rural area by three approaches: a field survey for estimating the degree of injury to Japanese Cedars; a biochemical study to clarify the mechanisms of inhibition of tannin biosynthesis in the Cedars; an epidemiological study of respiratory symptoms in children.

Kaye, Herbert (1987). Clinical and Research Applications of Quantitative Electrophysiology. Quantified Signal Imaging Inc., North York, Ontario; 1987.

A new computerized diagnostic system which provides physicians with a reliable and affordable window to the brain's electrical activity has recently been unveiled by an Ontario-based company. This advanced brain imaging system provides a high resolution colour image of the brain's electrical activity which accurately mirrors the brain's state of health. It can, both in the hospital clinic and clinical professional office, serve for assessing patients with head-trauma, substance abuse effects, cerebrovascular disease, stroke, epilepsy, genetic disorders, sensory-motor disabilities, diseases of the kidney and liver, and a variety of illnesses. Qualitative electrophysiological testing can also be used to evaluate persons who have been exposed to environmental chemicals at toxic levels. Such procedures have also provided support for the contention that there are specific changes in brain function that accompany cognitive or affective shifts in patients who are sensitive to exposure to common foods and odours.

Keating, John P. (1983). Environmental Stressors: Misplaced Emphasis. *Stress and Anxiety*, Chapter 3, pp 55 - 66; Sarason, Irwin G.; Spielberger, Charles D. ed; Hemisphere Publishing Corporation; 1983.

Although environmental factors may contribute to stress, stress-related behaviour will always result from the person and that person's cognitive structure of the environment.

Keller, Martin D.; Lanese, Richard R.; Mitchell, Ralph L.; Cote, Roger W. (1979). Respiratory Illness in Houses Using Gas and Electricity for Cooking. *Environmental Research*, No. 19, pp. 495-503; 1979; Academic Press, Inc.

Keywords: gas cooking appliances; indoor air pollution; respiratory illness; health hazards;

Kennedy, Donald A. (1977). Community Health and the Urban Environment: A Report of the Inter-University Board of Collaborators. The Effect of the Man-Made Environment on Health and Behavior, Chapter 2, pp 7 - 44; Centre for Disease Control, Public Health Service, U.S. Dept. of Health Education and Welfare, Atlanta GA; 1977; Hinkle, Lawrence E., Loring, William C., ed.

American cities are unhealthy places in which to live, work, play, or visit. All persons using the city suffer to some degree from the noise, air pollution, traffic congestion, crowding and lack of open areas of trees, lakes, grass, and rivers. Each person brings to the city a certain set of vulnerabilities and handicaps. If his or her residential area has a high crime rate, the work situation has high levels of toxic chemical exposure, and his or her children are denied an adequate education due to racial discrimination, civil disorders and inadequate tax revenues, then he or she operates under a burden of stresses and potentially hazardous conditions that are likely to affect his or her level of health and longevity. Self-directed change is possible with the open collaboration of three groups: community developer/managers, community resident-clients, and health researchers. A comprehensive bibliography is included.

Kerr, H.D.; Kulle, T.J.; McIlhany, M.L.; Swidersky, P. (1975). Effects of Ozone on Pulmonary Function in Normal Subjects: An Environmental-Chamber Study. American Review of Respiratory Disease, Vol. 111, pp. 763-773; 1975.

Keywords: ozone; pulmonary function; indoor air pollution; health hazards; NHW

Kerr, H.D.; Kulle, T.J.; McIlhany, M.L.; Swidersky, P. (1979). Effects of Nitrogen Dioxide on Pulmonary Function in Human Subjects: An Environmental Chamber Study. Environmental Research, No. 19, pp. 392-404; 1979; Academic Press.

Keywords: nitrogen dioxide; pulmonary function; indoor air pollution; health hazards;

Kerrebijn, K.F.; Mourmans, A.R.M.; Biersteker, K. (1975). Study on the Relationship of Air Pollution to Respiratory Disease in Schoolchildren. Environmental Research, No. 10, pp. 14-28; 1975; Academic Press, Inc.

Respiratory symptoms and ventilatory functions were measured in fourth and fifth grade school children living in two areas that differ in air pollution but are comparable with respect to social and demographic factors. Children living in the high polluted area showed a higher prevalence of cough during the day or at night, which is assumed to be due to the difference in pollution level. Mean ventilatory function in the children from both areas did not differ. In both areas mean PEF (peak

expiratory flow rate) and mean FEV (forced expiratory volume) of children with symptoms were lower than in children without. Depending on the criteria used, prevalence figures for chronic respiratory disease ranged between 5.3-12.7% in the high polluted and 3.3-9.8% in the low polluted area.

Kershner, John R.; Morton, L.L. (1984). Negative Air Ionization Improves Memory and Attention in Learning-Disabled and Mentally Retarded Children. *Journal of Abnormal Child Psychology*, Vol. 12, No. 2, pp. 353-366; 1984.

The effect of increased concentrations of ambient negative air ions on incidental visual memory for words and purposive auditory memory was investigated in 20 normal grade 4 children, 8 learning-disabled children, and 8 mildly mentally retarded children. All of the children breathing negatively ionized air were superior in incidental memory. The action of negative ions on the neurotransmitter, serotonin, may be the mechanism by which negative ions produce such behavioural effects.

Kershner, John; Stansfield, Michael; Kershner, Barbara; Hadfield, Audrey (1985). Air Quality in the Classroom Affects Memory in Four Learning Impaired Children. OISE; 1985.

The air in a special education classroom was modified alternatively over a three month period so that every two weeks the children were breathing air with a high ambient concentration of negative air ions. Under double-blind conditions, the four learning impaired children in this class were assessed on measures of immediate, memory, letter naming, text copying, math, spelling and a parent-teacher questionnaire. When breathing negatively ionized air, in comparison to a placebo-air condition consisting of regular classroom air, all of the children, irrespective of their unique psychoeducational characteristics, demonstrated a moderate improvement in short-term visual and auditory memory. One child showed a large improvement in classroom behaviour.

Kim, Yoon Shin; Spengler, John D.; Yanagisawa, Yukio (1985). NO₂ Concentrations in Offices With Kerosene Space Heaters and Electric Stoves. Air Pollution Control Association, 1985.

As part of a study to evaluate the impacts of use of stove or heater on indoor NO₂ concentrations, a pilot study on NO₂ measurements in 20 offices was conducted using diffusion tube and filter badge personal samplers during January and February of 1984 in Seoul, Korea. Offices with kerosene space heaters had average NO₂ levels approximately four times higher than the corresponding levels in offices with electric stoves. Average NO₂ concentrations above 50 ppb of the Korea ambient NO₂ standard were exceeded in approximately 71% of offices with kerosene heaters.

King, David S. (1981). Food and Chemical Sensitivities Can Produce Cognitive-Emotional Symptoms. *Nutrition and Behavior*; chapter 11; pp 119 - 130; Franklin Institute Press, Philadelphia PA; 1981; Miller, Sanford A., ed.

Exposure to allergens and other environmental substances has been clinically reported to provoke cognitive, emotional, and behavioral reactions in sensitive individuals. Controlled experiments directly testing such observations are limited. However, several studies provide evidence consistent with the possibility of allergic exposure provoking psychological reactions, although alternative explanations are possible. One double-blind experiment has tested whether sublingual allergy testing could provoke cognitive-emotional symptoms. In this study, most of the 30 allergy outpatients were abnormal on at least one MMPI scale. Cognitive-emotional symptoms were reported significantly more often on allergen trials than on placebo trials, as were psychosomatic symptoms, but not somatic symptoms. Heart rate change variability was greater on allergen trials. A follow-up MMPI found a significant improvement on the MMPI overall. Future research is recommended and suggestions are included.

King, E. (1982). Lead Poisoning. *Public Health Review*, Vol. 10, No. 1, pp. 49-76; Technosdar Ltd.— International Scientific Publications; 1982.

Despite the enormous research effort, as evidenced by the number of publications in recent years, there is still no clear understanding of the mechanism of lead poisoning or consensus of opinion as to the importance, or even existence in some cases, of the effects of lead below those of clinical lead poisoning. This article discusses how we have arrived at our present impasse in both the occupational and communal fields.

Kinney, Jennifer M.; Parris Stephens, Mary Ann; McNeer, Ann E.; Murphy, Michael R. (1985). Personalization of Private Spaces in Congregate Housing for Older People. *Environmental Change/Social Change*; EDRA 16; pp 184 - 188; Environmental Design Research Association; 1985.

As older people move from homes in the community to congregate housing, the housing environment becomes the residents' microcommunity. Similarly residents' apartments in the facility may assume the role of the homes they left behind in the community. This relocation may be stressful for older people, in that they are leaving a familiar home environment and meaningful social ties. It has been suggested that bringing familiar possessions from a past home will facilitate adaptation to the facility. Personalization brings a sense of control, and reinforces self-identity while communicating values to others, enabling social ties to develop. This study examines how older residents of congregate housing feel about and personalize their apartments, and identifies demographic and health

variables influencing the amount and type of personalization in which they engaged. Analyses indicate the importance of environmental factors in making a place feel homelike. Health variables account for more variance than age and gender in the amount of functional and decorative possessions appearing in residents' rooms. Residents with more decorative possessions tend to be more satisfied with their apartments. Results from this research have implications for planners and designers of congregate housing for older people.

Kinsman, Gary (1987). *The Regulation of Desire: Sexuality in Canada*. Black Rose Books, New York, 1987.

The author notes that as of March 1987, Quebec, Ontario and the Yukon Territory are the only Canadian Jurisdictions to have added "sexual orientation" protection to their human rights codes. He notes that a number of municipalities have instituted limited forms of protection and many unions have won such protection in their contracts. At the same time, he emphasizes that even today, governments, social agencies, businesses, and landlords continue to deny lesbian and gay individuals basic civil and human rights.

Kirsh, Sharon (1983). *Unemployment: Its Impact on Body and Soul: Questions and Answers Addressing the Human Costs of Unemployment*. Canadian Mental Health Association; 1983.

Responses to losing a job follow a general grieving process made up of shock; optimism and attempts to find employment; pessimism, depression and anxiety; and fatalism, accompanied by low job-seeking, apathy/despair, then adaptation to a lower standard of living. Factors which moderate the effects of unemployment include the availability of social, physical, material, and psychological support and externalization of blame onto the economy and government leading to action for dealing with one's anger. There is little research into the impact of unemployment on women workers. The stress of unemployment can also cause family breakdown, contribute to child abuse and adolescent crime, and create school-related problems for children. Older workers who suffer unemployment are victimized by age discrimination. Mental hospital admission rates vary according to economic changes, however job availability for the chronic psychiatric patient presents a special set of problems including the stigma of mental illness, the effects of some medications, gaps in resumes, lack of work incentives, and an entrenched sheltered-workshop mentality.

There is little research into the connection between unemployment and physical health. But there is a causal connection between chronic stress and several infectious diseases, cancer, heart disease, stroke, and ulcers. In addition, people are more likely to use addictive substances like alcohol and tobacco during periods of unemployment. Economic in-

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

stability and poverty also create poor nutritional levels, and decreased ability to utilize medical care facilities. Primary prevention techniques in both the areas of mental and physical health as they relate to unemployment are discussed, and a comprehensive bibliography is presented.

Kittler, Fred J.; Baldwin, Deane G. (1970). The Role of Allergic Factors in the Child with Minimal Brain Dysfunction. *Annals of Allergy*, Vol. 28, pp. 203-206; May, 1970.

Formal elementary education finds many children with learning disabilities and behavior problems. Many manifest some combination of the signs and symptoms of the minimal brain dysfunction syndrome. Allergy of the tension-fatigue syndrome general classification is present in a significant percent of children with minimal brain dysfunction syndrome and the authors believe allergy causes them to have findings of this syndrome.

Kleiman, Cindy (1987). Allergic Students Will Soon Breathe More Easily. *Toronto Star*, North Section, October 6, 1987.

Students of the York Region Separate School Board who suffer from severe allergies or hypersusceptibility will be housed in two ecological classrooms designed with high efficiency air filters and a minimum of synthetic materials. The classrooms are part of a two-part strategy to decrease the number of allergens in school buildings. New schools are being constructed with sensitivity to the average child's allergies.

Kon, S.H. (1978). Underestimation of Chronic Toxicities of Food Additives and Chemicals: The Bias of a Phantom Rule. *Medical Hypotheses*, No. 4, pp. 324-339; 1978.

Keywords: food additives; chemicals; toxicity;

Konopinski, Virgil J. (1983). Formaldehyde in Office and Commercial Environments. *American Industrial Hygiene Association Journal*, No. 44 No. 3, pp. 205- 208; March 1983.

Keywords: workers; office; formaldehyde; commercial buildings; indoor air pollution;

Korosec-Serfaty, Perla (1985). Experience and Use of the Dwelling. *Home Environments, Human Behavior and Environment*; Vol. 8; pp 65 - 86; Plenum Press New York; 1985; Altman, Irwin and Werner, Carol M., editors.

This study looks at how the relationship to home is experienced by the dweller, especially in relation to people's reactions to burglary as it ruptures the inside/outside boundaries created by one's home. In

this context, the author emphasizes the importance of comprehending the whole dwelling experience, including the modes of the relationships which are established within the home. Being burglarized is experienced as being defiled and causes a specific psychological impact on the person's relationships with others.

Kose, Satoshi; Naoi, Hideo; Uno, Hidetaka (1986). An In-depth Study of Accidental Injuries Associated With Building Features: Translating Research into Practice. CIB.86, Advancing Building Technology, Volume 7, pp 2877 - 2884; 1986.

A questionnaire survey was conducted on the incidence of building related accidents in dwellings. About a thousand questionnaire forms were received from the residents in the Tokyo metropolitan area, covering over 3,500 family members. The result was compared with other data. It is found that accident types with high injury rates are not necessarily the same as those leading to death; severity of injuries differs among age groups even for the same accident type. Some of the implications to the building characteristics, such as stair design, are discussed, in order to reduce the tragic outcome of domestic accidents.

Kose, Satoshi (1987). Survey of Domestic Stair Conditions and Safety in Japan. unpublished paper presented at EDRA-18, Ottawa Canada.

A mail questionnaire survey on the present conditions of domestic stair design and accident incidence on stairs was conducted for Japanese dwellings. Two hundred and twenty-eight samples were used for analysis. The results suggest that pitch of Japanese domestic stairs ranged between 45 and 55 degrees with very small tread goings, and they were not necessarily provided with handrails. Reported accident incidence among the samples was fairly low and it was difficult to pinpoint factors of design that determine the level of danger for any type of stairs. Further in-depth study on accidents is needed to reveal quantitative basis for safer stair design.

Kreiner, C. (1986). Address on Oppression of Working-Class Men. For Men, Number 11, pp. 4-5; London, England, Autumn 1986; Excerpts from an address to a Men's Workshop in Yorkshire, England, 1985.

The author describes the exhaustion, poisoning, pollution, manipulation, physical exploitation of working conditions for many working class men in Western society. "We are told that endurance of this is manly - rather than be encouraged to reclaim our power to change the conditions. Miners', factory and chemical workers' conditions damage them physically, but they are taught to feel ashamed of protesting or saying 'this hurts' or 'this is not right' because these conditions are manly to endure."

Kreiss, K.; Gonzalez, M.G.; Conright, K.L.; Scheere, A.R. (1982). Respiratory Irritation Due to Carpet Shampoos: from Indoor Air Pollution; Spengler, John—editor; pp. 337-341. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Dried detergent residue left in carpets after they were shampooed with underdiluted carpet shampoo caused respiratory irritation among most employees in an office building and among all staff members and most children in a day-care centre. Symptoms included cough, dry throat, difficulty in breathing, nasal congestion, and headache. Eye irritation was also noted by day-care centre staff members. Symptoms persisted for many weeks until the carpets were wet extracted. The major ingredient of the three shampoo products implicated in these two outbreaks and in a third similar report is sodium dodecyl sulfate, a respiratory irritant in mice. Detergent dust is a newly recognized example of indoor air pollution and should be considered when patients or employees complain of building-specific respiratory or eye irritation.

Krieger, Martin H. (1970). Six Propositions on the Poor and Pollution. Policy Sciences, No. 1, pp 311 - 124; 1970.

The effect of improving the environment may be greater inequities in our society. Current environmental programs maintain this inequity, proposed environmental programs may make things worse, and even if we do improve the environment, contentment may decrease. A political coalition of environmental and equity enthusiasts may provide a viable way out of these dilemmas.

Kruss, P.; Valeriote, L.M. (1979). Fluoride: from Controversial Chemicals, A Citizens' Guide. Multiscience Publications Limited, Montreal Canada; 1979; pp 92-98.

Industrial processes release fluoride into the environment, damaging plants and animals. There is a controversy over the fluoridation of drinking water and its effect on health. The chronic effect of long term exposure to fluoride has been blamed for an increased incidence of mongolism, a variety of symptoms such as headache, nausea and loss of weight, bone abnormalities such as osteosclerosis, and kidney and thyroid injury.

Kruss, P.; Valeriote, L.M. (1979). Asbestos. from Controversial Chemicals, A Citizen's Guide; pp. 17-30; Multiscience Publications Ltd., Montreal PQ; 1979.

Asbestos has long been known as a threat to health. It causes asbestosis—severe scarring of the lungs caused by continually inhaling asbestos over the course of many years. The symptoms begin with

shortness of breath, which develops into near paralysis that makes breathing and body movement increasingly difficult, leading to ultimate suffocation. It is now accepted that asbestos is also an agent for cancer of the abdomen, liver, gastro-intestinal tract and lung. The risk of lung cancer appears to be synergistic with smoking. It is widely used for filters in the chemical, food and beverage industries, resulting in asbestos fibres in most beers, wines and liquors, soft drinks, sugar and lard. It is also found in our drinking water.

Kruus, P.; Valeriote, I.M. (1979). Controversial Chemicals—a Citizen's Guide: Canada Science Series. Multiscience Publications Limited, 1253 McGill College, suite 175, Montreal Canada H3B 2Y5; 1979.

25 chemicals are listed in alphabetical order, a chapter devoted to each. Historical, technological, medical, economic and political backgrounds of these controversial chemicals are reported.

Kruus, P.; Valeriote, I.M. (1979). Lead: from Controversial Chemicals, A Citizen's Guide. Multiscience Publications Limited, Montreal Canada; 1979; pp. 128-136.

Lead is used in gasoline additives, and is therefore an ever-present pollutant in urban areas. Leaded gasoline is the largest single source of gaseous atmospheric contamination. The Toronto Refiners and Smelters and Canada Metal Co. cases are cited. Health effects in humans and effects on wildlife and vegetation are discussed. Lead affects three systems in people: the nervous system, the kidney and red blood synthesis. Government regulation in Canada is described.

Kruus, P.; Valeriote, I.M. (1979). Sulphur Dioxide: from Controversial Chemicals, A Citizen's Guide. Multiscience Publications Limited, Montreal Canada; 1979; pp 190-195.

Acidic precipitation is increasing due to the growing introduction of sulphur dioxide into the atmosphere. It has been linked with detrimental effects on fish, vegetation, buildings and people. Human health problems are related to irritation of the respiratory system.

Labelle, Huguette (1986). Nurses as a Social Force. Journal of Advanced Nursing, No. 11, pp. 247 - 253; 1986.

Nurses in every continent engage in social action. This has been a fact throughout the ages. The major social issues which contribute to health today are economic, energy, environmental and social welfare problems. Other major issues affecting health are population growth, poverty, education, clean water supply, and family planning. The author shows how individual nurses and nurses' associations can, and should, exercise

their influence and power in promoting health at local, national, and international levels, making primary health care a priority. By using their levers of power, together with technology and communications, nurses can help create a new world health order.

Lachtman, Dennis S. (1980). Human Data Associated with Diesel Exhaust. from *Health Effects of Diesel Engine Emissions*, Vol. 2; pp. 1100-1113; Pepelko, W.E.; Donner, R.M.; Clarke, N.A., editors; November 1980.

A review of the literature concerning data relative to human exposures from diesel engines is presented. Epidemiologic evaluations among workers exposed to diesel exhaust are discussed. Morbidity and mortality data are critically reviewed. No conclusive evidence suggesting an association between diesel exhaust exposures and adverse long term health effects was found, but more studies involving humans evaluating health effects are needed. Those constituents found in diesel exhaust that are of primary health concern are oxides of nitrogen, particulates and sulphur oxides.

Laseter, John L.; DeLeon, Idefonso R.; Rea, William J.; Butler, Joel R. (1983). Chlorinated Hydrocarbon Pesticides in Environmentally Sensitive Patients. *Clinical Ecology*, Vol. II, No. 1, pp. 3-12; Fall 1983.

The sensitive and selective analytical techniques of high-resolution gas chromatography and high-resolution gas chromatography-mass spectrometry were employed to characterize 16 different synthetic chlorinated hydrocarbon pesticides and common metabolites present in randomly selected environmentally sensitive patients. Of 200 patients initially screened, 99% had residues at or above the .05 ppb level in their sera.

Lassiter, Donald V.; Milby, Thomas H. (1978). Health Effects of Diesel Exhaust Emissions: A Comprehensive Literature Review, Evaluation and Research Gaps Analysis. Environmental Health Associates, Inc., 2150 Shattuck Ave., Berkeley CA 94704; January, 1978.

The purpose of the report was to provide a comprehensive literature review of the human health consequences of exposure to diesel exhaust emissions, especially as encountered in underground mines, and to define those gaps in current scientific knowledge which require further clarification through additional research. The toxicology of diesel exhaust both as a mixture and as individual toxicants was examined. In a properly ventilated mine, neither CO, SO₂, nor NO₂ were found to be major toxic problems. Polycyclic aromatic hydrocarbon (PAH) compounds were identified as the most potentially hazardous agents found in diesel engine exhaust. Although a properly maintained diesel engine would be likely to emit extremely small amounts of these agents, the possibility that they might be carcinogenic to humans singles them out for careful attention.

Last, John M. (1987). Scenarios and Methods to Support Long Term Health Planning: Conference Report. Canadian Journal of Public Health; Vol. 78, pp 11 - 12; January/ February, 1987.

There is a great necessity and urgency for government involvement in long range health planning for maximum flexibility, based on changing demographics of population.

Laties, Victor G.; Beard, Rodney R.; Dinman, Bertram D.; Schulte, John H. (1969). Behavioural Aspects of Carbon Monoxide Poisoning. from Effects of Chronic Exposure to Low Levels of Carbon Monoxide on Human Health, Behaviour, and Performance; pp. 32-37; US National Research Council, Washington DC; 1969.

It is apparent that there has been very little modern work on CO and behavior. Most investigators have noted symptoms of CO poisoning first when the level of CO in the blood was raised to between 10% and 20% but many symptoms don't appear until the concentration of CO in the blood increases beyond 25%. However, Schulte has shown positive correlations between CO concentrations in the blood and performance decrement in a number of tests. For example, his subjects were required to do 50 arithmetic problems, each involving five five-digit numbers. They took about 13 minutes to complete the task under control conditions but about 17 minutes when their COHb level was 20% and made more errors under the latter conditions.

Laufer, Robert S.; Wolfe, Maxine (1977). Privacy as a Concept and a Social Issue: Multidimensional Developmental Theory. to be published in the Journal of Social Issues, Winter 1977/78.

If we are to understand privacy as a future as well as a contemporary social issue, we must understand it as a concept. Individuals' concepts of privacy are tied to concrete situations in everyday life. These situations are described in terms of three dimensions: self-ego, environmental, and interpersonal. In combination with the dynamic of time, both developmental and sociohistorical, this analysis helps us to understand individual perceptions of privacy and privacy invasion, to predict potential privacy or invasion experiences, and to see the potential effects of the absence of certain privacy-related experiences.

Laurie, R. Dana; Boyes, William K.; Wessendarp, Thomas (1980). Behavioral Alterations Due to Diesel Exhaust Exposure. from Health Effects of Diesel Engine Emissions; pp. 698-712; Pepelko, W. E., Danner, R.M., Clarke, N.A., editors; US Environmental Protection Agency; November 1980.

Several experiments examining the effects of diesel exhaust on the behavior of rats are reported. Animals were exposed either as adults or neonates. The spontaneous locomotor activity (SLA), measured in stan-

dard running wheel cages, of adult rats exposed for 8 hours per day, 7 days per week was significantly less than that of controls. Experiments involving diesel exhaust exposure to neonatal rats indicated that adult rats, exposed to diesel exhaust during their neonatal lives, were significantly less active as measured by SLA. Adult rats, exposed to 20 hours of diesel per day as neonates, were placed in skinner boxes after the SLA experiment described above had been completed. The exhaust exposed animals showed significantly decreased acquisition of a food reinforcing bar pressing task. All animals that learned this task extinguished at the same rate. The results of the neonatal diesel exhaust experiments support the hypotheses that diesel exhaust exposure during development of an organism can lead to behavioural differences in adulthood.

Laverne, Albert A. (1970). Air Pollution, Healing, and Civilization. Behavioral Neuropsychiatry, Vol. 2, No. 3,4; pp 26 - 38; 1970.

Medical evidence has shown that inhaled air pollutants can cause and aggravate respiratory diseases and can affect the brain and central nervous system, body defense mechanisms, and other vital organ systems. Air pollution also has a deleterious effect on overall work efficiency, intellectual and emotional functioning, and behaviour. There is even evidence that the healing processes may be substantially impaired, prolonged, or actually prevented by air pollutants. Double blind studies performed on both psychiatric and infectious disease patients appear to confirm these observations. In a group of 250 patients with acute and chronic psychiatric disorders, improvement was significantly greater when pollution-free air was administered immediately post-treatment. During the course of any disease, tissue resistance is lowered, and tissue susceptibility to toxins is increased.

Layton, Jack (1987). Healthy Toronto 2000: A Discussion Paper. Toronto Board of Health; August, 1987.

Health does not result simply from the actions of professionals in the health care system, but from a complex interplay of biological, psychological, social, environmental and political factors acting on the individual and family through the local community and through the whole human ecosystem. Health promotion and disease prevention are better than cures. A multi-sectoral approach is necessary. Social interventions for the common health are appropriate and may take precedence over individual concerns. Community development is the proper approach to many health problems. The vision presented in the report includes: appropriate technology in waste management, transportation, food production, energy use, and manufacturing process; quality food, shelter, work, safety and education with the emphasis on self reliance rather than dependence; integrated caring neighbourhoods, built to human scale; support services for those with special needs through a strong social support network; a wellness system rather than an illness system; and involvement of the whole community. Public participation is requested.

Lebowitz, M.D. (1983). Health Effects of Indoor Pollutants. Annual Review of Public Health, Vol. 4, pp. 203-221; 1983.

Sources, concentrations, health risks and public health concerns associated with indoor pollutants are analyzed. Both outdoor and indoor sources of pollutants are responsible for indoor pollutant concentrations. The pollutants that are predominantly outdoors include sulfur oxides, ozone and related oxidants, most trace metals, halogen compounds, petrochemical compounds, pesticides, ultra-violet radiation, and pollen. Indoor-based pollutants include carbon monoxide, respirable particles, organic vapours, nitrogen dioxide, formaldehyde, radon, asbestos, mineral and synthetic fibers, carbon dioxide and viable organisms.

Lebowitz, M. (1986). Passive Smoking and Health Effects. Indoor Air, Vol. 6, 1986; Swedish Council for Building Research; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

The area of most difficulty in passive smoking is one of establishing exposure or dose. More research is required on statistical methods, and on measurement methods, methods of pulmonary function, more studies on chronic disease, and more research on carcinogens in smoke and lung cancer, especially on dose-response relationships to know of any association effects.

Lecuyer, Gerard; Aitken, Roy (1987). Report of the National Task Force on Environment and Economy. Canadian Council of Resource and Environment Ministers, Task Force on Environment and Economy; September, 1987.

Complete integration of the economy and environment is possible and necessary. Long-term economic growth depends on a healthy environment and affects the environment in many ways. Ensuring environmentally sound and sustainable economic development requires the technology and wealth that is generated by continued economic growth. A new cooperative initiative is proposed to integrate economic and environmental planning through the participation and debate of senior decision makers at the provincial and federal levels. Also recommended is the development of conservation strategies as a valuable multisectoral approach to defining and implementing sustainable economic development. Public participation and improved environmental education are also recommended.

Lee, A. John (1987). Recognizing the Importance of Ethnicity for Canada's Elderly: Short Report. Canadian Journal of Public Health, Vol. 78, p. 6; January/February, 1987.

It is imperative that our consideration of the health and social services and research on the needs of the elderly in Canada extend beyond the mainstream white middle class. Changing values, language capabilities and cultures exacerbate the already complex issue of aging.

Letourneau, E.G. (1985). Development of a Radon Standard for Canada. Air Pollution Control Association, 1985.

The jurisdictional responsibility for natural radioactivity in Canada is divided between the federal and provincial governments. Over the years various standards for radon daughters in homes have been derived for special circumstances involving both radon arising from man-made contamination and natural occurrence. Wide spread surveys have indicated that these initial standards were not appropriate for general use. The desire for uniform standards has resulted in the formation of a scientific group to advise a federal-provincial committee concerning the desirability of standards for exposure to natural radioactivity. The scientific committee could not agree either philosophically or practically on the control level for radon in homes. To resolve this impasse, it undertook a calculation designed to illustrate for Canada, the potential cancers saved each year compared to the actual lung cancer rate, with an estimate of the cost, for each limit derived. A second calculation reviewed the cost of modifying the building code and estimated the long term effect comparing avoided cancers and cost.

Letz, Richard; Ware, James H.; Ferris, Benjamin G.; Spengler, John D. (1982). Pulmonary Functions of Children and Indoor NO₂ Concentrations in Portage, Wisconsin: from Indoor Air Pollution: Proceedings of the International Symposium on Indoor Air Pollution, Health and Energy Conservation; Spengler, John—ed. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

A study of NO₂ concentrations inside 137 homes in Portage Wisconsin was made. An attempt was made to directly relate the observed lung functions of the children living in these homes to the observed NO₂ concentrations. This was not found to be possible in this study. Several explanations are put forward: 1) NO₂ may not be the offending agent. Gas stoves emit carbon monoxide, carbon dioxide, sulfur dioxide, nitrogen dioxide, formaldehyde, hydrogen cyanide, water vapour, carbon, and sulfate particles. 2) Perhaps only a sensitive sub-population is affected by the offending agent, and they may not have been well represented by this group. 3) Peak levels of NO₂ may cause these health effects and long-term averages are poorly related to those peaks in this study. 4) Perhaps such a relationship exists and can only be detected by a much larger sample size.

Levi, Lennart; Frankenhaeuser, Marianne; Gardell; Bertil (1983). Report on Work Stress Related to Social Structures and Processes: Analysis and Implications of Research/A Study by the Institute of Medicine, National Academy of Sciences. Stress and Human Health; chapter 6, pp 119 - 146; Elliott, Glen R.; Eisdorfer, Carl ed; Springer Publishing Company, New York; 1983.

Important stressful psychosocial situations fall into four general categories: uprooting; dehumanization of society institutions; psychosocial side-effects of the spread of innovations; and psychosocial factors as constraints on environmental health programs and activities. A substantial body of indirect evidence strongly suggests that such social structures and processes affect the prevalence of ill health, lack of well-being, and low quality of life. Evidence suggests that the following interventions should be researched as means for reducing workplace-related stress: increasing a worker's control of work arrangements; providing mechanisms for worker participation in decision making on the organization of work; avoiding monotonous, machine-paced, and short but frequent work actions; optimizing automation; helping workers see their specific task in relation to the total product; avoiding qualitative work overload or underload; facilitating communication and support systems among work mates and others. A comprehensive bibliography is included.

Lewis, Beatrice E. (1985). The Stigmatized House. Environmental Change/Social Change; EDRA 16; pp 184 - 188; Environmental Design Research Association; 1985.

When a child with severe chronic mobility problems becomes too heavy or too cumbersome to lift and to carry, parents must consider removing architectural barriers in the home. Practical and psychological difficulties frequently cause them to postpone making such changes until a crisis occurs. Practical difficulties include lack of resources, gaps in services, and a general disregard by medical personnel of how environments affect behavior. Psychological difficulties, which has so far received little attention, mainly stem from conflict between the parents' desire for an idealized house and their child's functional requirements, as well as from the parents' desire to appear as normal to their neighbors. Adapting the home means accepting the permanence of the disability. It also means making the family's disability public — stigmatizing the house, and thus its occupants. These conclusions are based upon interviews with six families raising children who have severe mobility problems as well as upon residential histories taken from parents of these families and upon drawings of one's ideal home made separately by parents and by the disabled child.

Lewis, Myrna (1985). Older Women and Health: An Overview. Women and Health...the Journal of Women's Health Care; No. 2/3; pp 1 - 16; 1985.

Older women's health issues are unique. There are more older women than ever before. They are living increasingly longer than men, yet they report more acute and chronic illness and disability than men. They are disproportionally represented in nursing homes, since many women are alone: 25% aged 70 or over have no living children and over 60% of older women are widowed, divorced, or single. Older women have fewer personal financial resources for health care than men. They face age and sex

discrimination on the part of many health care providers and are subject to a growing tendency to be seen as burdens and problems in the health care system.

Lidwell, O.M. (1979). Ventilation, Air Movement and the Spread of Bacteria in Buildings: from Indoor Climate: Effects on Human Comfort, Performance, and Health in Residential, Commercial, and Light-industry Buildings; Fanger, P.O. and Valbjorn, O.—editors; pp. 239-256. Danish Building Research Institute, Copenhagen Denmark; 1979.

Keywords: ventilation; bacteria; indoor air pollution; health hazards;

Lindstrom, Kari; Harkonen, Hannu; Hernberg, Sven (1976). Disturbances in Psychological Functions of Workers Occupationally Exposed to Styrene. Scand j work environ & health, No. 3, pp. 129-139; 1976.

Keywords: styrene exposure; polyester; psychological functions;

Liss, Gary M. (1987). Health Effects of Welding and Cutting Fumes and Gases. Occupational Health in Ontario, Vol. 8, No. 2, pp 69 - 78, Spring, 1987; Ontario Ministry of Labour, Occupational Health Branch, Toronto Canada.

The health effects, both acute and chronic, due to exposure to the products of welding and cutting processes are described. The welding process produces potential respiratory hazards with a particulate fraction, which is largely respirable metal oxide fume, and gases, of which ozone and oxides of nitrogen are the most dangerous. The welding environment may also contain contaminants that are extraneous to the welding process, such as gases and particulates. The available data do not permit the health effects to be correlated with cumulative duration of exposure, much less with levels of specific constituents that can be deemed safe or not safe, as they relate specifically to the welding environment. An extensive bibliography is included.

Lofgren, Inger (1981). Health and Safety in Buildings: R & D Programme, 1981-1984. Swedish National Council for Building Research.

This publication sets out the basis for the plan of action of the Swedish National Council for Building Research for the years 1981 to 1984. It notes that in order for further improvements in public health to be made, public health problems must be attacked by preventive measures such as identification of health hazards in the environment. Areas of concern are mould and other biological factors, thermal climate, radon, electromagnetic fields, air ions, lighting, noise, infrasound, vibration, chemical factors, addition and interaction effects, psychological effects, risk assessment, and epidemiological evaluations.

The research objectives of the group as stated are: to identify factors in buildings which may have a negative effect on the health of people; to analyze health hazards, taking into account biological effects, dose-effects and dose-response relationships, interactions, variations in the sensitivity of the population on exposure (with special reference to risk groups); and to develop appropriate technical and structural methods whereby health hazards can be reduced.

Lund, S.M.; Dowdle, E.B. (1977). The Effect of Prolonged Isolation from Environmental Allergens on the Clinical and Laboratory Manifestations of the Allergic State: Observation on Members of the South African Antarctic Expedition. SA Medical Journal, No. 52, pp. 556-561; September 24 1977.

Keywords: environmental allergens; treatment;

Lundqvist, G.R.; Iversen, Martin; Korsgaard, Jens (1982). Indoor Climate in Low-Ventilated Day-Care Institutions: from Indoor Air Pollution; Spengler, John—editor; pp. 139-142. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Measurements of indoor atmospheric environment were carried out in a day nursery and a kindergarten, in which natural air infiltration supplied the rooms with only 1-2 m³ fresh air per person and hour under the prescribed occupancy. The exposure situation from the field was duplicated under controlled conditions in a climate chamber. The results indicate unacceptably high concentrations of carbon dioxide, and emphasize the need for more elaborate building design in airtight buildings.

Lykke, A.W.J.; Stewart, B.W. (1977). Fibrosing Alveolitis (Pulmonary Interstitial Fibrosis) Evoked by Experimental Inhalation of Gasoline Vapours: Specialia. Experientia, Vol. 34, No. 4, p. 498; August 1977.

Keywords: gasoline vapours; health hazards; air pollution; lung function;

MacLeod, Kathryn E. (1981). Polychlorinated Biphenyls in Indoor Air. Environmental Science and Technology, Vol. 15, No. 8, pp. 926-928; May 1981; American Chemical Society.

PCBs have been recognized as environmental contaminants since the mid-60s. Until recently, however, little work has been done on identifying levels of PCBs in the indoor air. This paper describes a method for the analysis of PCBs utilizing low-volume indoor air sampling. The method uses polyurethane foam as a collector. This study shows that indoor air whether in commercial, industrial, or residential buildings, contains levels of PCBs at least 1 order of magnitude higher than outdoor levels. Defective fluorescent light ballasts are also shown to emit PCBs and to be an important source of indoor atmospheric contamination.

MacLennan, J.G. (1985). Hypersensitivity Reaction to Indoor Air Contamination in the Susceptible Individual. unpublished conference address.

The response of any sensitive person to environmental incitants is not predictable, and will vary according to that individual's degree of sensitivity and their own physical peculiarities, physiologically and biochemically. Usually there is a multiorgan response in those who have exhibited chronic symptomatology for many years. Occasionally a sudden massive chemical exposure, or some other severe traumatic experience may release a reservoir of latent or hidden sensitivities. This frequently results in a cascade of severe varied symptomatology which occasionally is incapacitating and prevents the individual from functioning normally.

There is no direct correlation between any particular exposure and a specific organ response. Frequently several organs may respond to a single environmental incitant. Successful ecologic management is based on an accurate diagnosis by establishing a cause and effect relationship between an environmental exposure and a symptomatic response. Then a program of avoidance, change in life style, and therapy as indicated, can be formulated.

MacLennan, John G. (1985). The Impact of Indoor Air Pollution on the Behaviour and Learning Ability of School Children: Sponsored by Air Pollution Control Association TT-7 Indoor Air Quality Committee, Health and Welfare Canada, National Research Council of Canada, Public Works Canada, Consumer and Corporate Affairs Canada, and Canada Mortgage and Housing Corporation. Presented at the International Speciality Conference on Indoor Air Quality in Cold Climates: Hazards and Abatement Measures; Ottawa, Ontario, April 29, 30 and May 1, 1985.

The author states that increasing numbers of the Canadian population are inheriting the ability to acquire hypersensitivities to different environmental exposures. The commonest indoor contaminants found in school are particulate inhalants and chemical fumes, odours and smokes. There are also hazards for the food sensitive person, but these can usually be managed by ecological control of lunches and supervision by school personnel. Important chemical exposures are derived from tobacco smoke, perfume, fossil fuels and their derivatives, plant terpenes, photocopiers, duplicators, cleaning and maintenance materials, craft and other office and school supplies. In the majority of cases, the brain and central nervous system are the major areas of hypersensitive reaction which is reflected in the child's behaviour and ability to learn.

Macpherson, A.S. (1984). Letter to the Minister of Labour re Tobacco Smoke. City of Toronto Dept. of Public Health; December 1984.

This is a letter recommending that the Minister of Labour make tobacco smoke a designated substance under the Occupational Health & Safety Act, and that a standard for tobacco smoke in the workplace be developed.

Macpherson, A.S. (1987). Environmental Health Effects of Waste Incineration in the City of Toronto. City of Toronto Department of Public Health; October, 1987.

The environmental fate and persistence of potentially hazardous chemicals in waste incineration emissions as well as their potential health effects are reviewed in this report. It is difficult to determine the contribution of chemicals emitted from incinerators compared to other loadings to the ecosystem. Also, data on potential health effects of many chemicals are incomplete, particularly for long-term, low-level exposures. A risk assessment was undertaken to predict the degree of the hazard to human health resulting from the emissions from two sewage sludge incinerators. In addition, a risk assessment was conducted on a hypothetical solid waste incinerator equipped with the best available technology. The results showed that emissions from one incinerator do not pose any significant risks to human health, but that the emissions from the other do. Some of the emissions from the hypothetical incinerator could also pose significant risks to human health. Further research is needed. Technical and operational measures need to be implemented to reduce emissions of chemicals and questions need to be asked about siting of new incinerators and acceptable levels of health and environmental risk.

Mahaffey, Kathryn R.; Vanderveen, John E. (1979). Nutrient-Toxicant Interactions: Susceptible Populations. Environmental Health Perspectives, Vol. 29, pp. 81-87; 1979.

Nutritional status can substantially modify the toxicity of environmental pollutants. Investigations with experimental animals and epidemiological observations on humans have established the role of nutrition in altering susceptibility to a variety of pollutants including pesticides and heavy metals. The degree of nutritional deficiency that alters susceptibility need not be severe. Frequently only biochemical indications of nutritional deficiency can be associated with changes in the dose-response of an animal or person to a toxic compound.

Makower, Joel (1981). Office Hazards: How Your Job Can Make You Sick. Tilden Press, 1737 DeSales St., NW, Washington DC 20036; 1981.

Energy efficiency, design deficiencies, combined with hazardous working materials, have helped to make air pollution in offices a bigger health problem than air pollution outdoors. A list of contaminants is provided. Fluorescent light, video display terminals and other sources of radiation join stress as being some of the main offenders.

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

Makower, Joel (1981). There's Something in the Air. from Office Hazards, chapter 1, pp. 9-37; Tilden Press; 1981.

Energy efficiency and design deficiencies have helped to make air pollution in offices a bigger health problem than air pollution outdoors. Substances such as ammonia, asbestos, benzene, cadmium, carbon monoxide, ethanol, fiberglass, formaldehyde, methanol, nitropyrenes, ozone, PCBs, particulates, radon, tobacco smoke, toluene, trichlorethane, trichloroethylene, trinitrofluorenone and vinyl chloride are all present in the typical office environment and all are possibly hazardous. Some are potential or proven carcinogens, while a number, such as trichloroethylene, carbon monoxide and benzene have been implicated in central nervous system damage, dizziness, and other symptoms that could cause learning problems.

Mallov, Joseph S. (1976). MBK Neuropathy Among Spray Painters. Journal of the American Medical Assoc., Vol. 235, No. 14, pp. 1455-1457; April 5 1976.

Keywords: painting; occupational health and safety;

Mandell, Marshall; Waller Scanlon, Lynne (1979). Physical and Mental Allergies in Children. from Dr. Mandell's 5-Day Allergy Relief System, chapter 5, pp. 133-187; Thomas Y, Crowell Publishers; 1979.

Many types of poor behaviour in schools, from hyperactivity, irritability, violence, fatigue, restlessness, to poor academic performance, can be caused by food and chemical allergies. In many cases, the nervous system is totally incapable of functioning normally because it is under allergic distress. The child who reads well once in a while and poorly at other times; the child whose handwriting is poor one day and fine the next; the child who can sit still in the morning, but is impossible to control in the afternoon - some of these children may be demonstrating cerebral allergy. Many case studies are provided from the author's practice.

Marchant, R.,; Wnesley, B.; Frel, D.; Bartram, J.; Yoshida, K. (1985). A Community Indoor Air Pollution Program. Air Pollution Control Association, 1985.

The Saskatoon Community Health Unit Indoor Air Pollution Program is a collaboration of specialist community resources. Trained Public Health Inspectors field all requests for appropriate follow-up. Through the definition of the problem, preliminary measurements (humidity, temperature, formaldehyde, CO 2) and a short history, the appropriate specialist team members are then involved. A suggested referral by the family doctor to a specialist in environmental medicine provides a thorough health assessment. If a more detailed environmental study or a study to define the pollutant is required, an Industrial-Hygienist-Aerosol-Scientist will

do a 24-48 hour study of the problem under an individual contract. Ventilation Engineers from the National Research Council provide consultation on remedial environmental changes and pollution source identification. The team meets on a regular basis for in-service training and feedback and review of current issues.

Marchant, R.; Yoshida, K.; Figley, D.A. (1985). Mobile Home Residents' Exposure to a High Concentration of Formaldehyde — A Case History. Air Pollution Control Association, 1985.

A series of health complaints by an elderly couple soon after they occupied a new mobile home in suburban Saskatoon Canada prompted this investigation. Their symptoms included nausea, vomiting, dizziness, headaches, sleep disturbances, shortness of breath, burning eyes, running nose, and chronic fatigue. There was an initial indoor formaldehyde concentration of 0.7 ppm, far exceeding the recommended limit for residential buildings of 0.1 ppm. Testing revealed that the wall panelling (approx. 2-3 ppm) had a significantly higher emission rate than the cupboards (approx. 1 ppm) or subfloor (approx. 0.4 ppm). The occupants' medical history and physical examination showed only symptoms related to chronic ailments. Since occupying an ordinary house, their symptoms characteristic of formaldehyde have subsided.

Margulis, Stephen T. (1981). Building Accessibility in Relation to Door Hardware, Door Users and Door Use. Architectural and Transportation Barriers Compliance Board, 33 C St., SW Washington DC 20201; January 1981.

Keywords: architectural barriers; building accessibility; codes and standards; disability; doors; ergonomics; handicap;

Marha, Karel (1982). The State of Knowledge Concerning Radiations from Video Display Terminals. Canadian Centre for Occupational Health and Safety; 1982.

Keywords: video display terminals; occupational health and safety; offices;

Marha, Karel (1983). Low Frequency Emission from VDT's and Possible Adverse Biological Effects. Canadian Centre for Occupational Health and Safety; April 29 1983.

Keywords: video display terminals; occupational health and safety; offices;

Marha, Karel (1983). Very Low Frequency Fields Near VDT's and An Example of Their Removal. Canadian Centre for Occupational Health & Safety; September 1983.

Keywords: video display terminals; vlf fields; occupational health and safety; shielding;

Markham, John W.; Kirkbride, John; Pelmeur, Peter L. (1986). Health Surveillance Data. Occupational Health in Ontario, Vol. 7, No. 4, pp 192 - 204; Fall, 1986; Ontario Ministry of Labour, Occupational Health and Safety Division, Toronto ON.

An adequate health information system is an essential part of worker health surveillance. In Canada, as in most western industrialized countries, the necessary systems have not yet been developed. This paper enumerates the principles and factors that should be considered, as well as the benefits that may be derived from good information systems. The desirable types of information include worker identification, worker disease susceptibility, relevant clinical tests, biological monitoring, morbidity, mortality, specific hazards or exposures in the workplace, exposure measurements, occupation, population-at-risk records, and reproductive information. An extensive bibliography is included.

Marks, James G. (1981). Allergic Contact Dermatitis from Carbonless Copy Paper. Journal of the American Medical Assoc., Vol. 245, No. 22, pp. 2331-2332; June 12 1981.

Allergic contact dermatitis to carbonless copy paper is documented. The offending agent proved to be a coating on the paper - a colour-forming chemical composed of paratoluene sulfinate of Michler's hydrol (PTSMH). It apparently has a low sensitizing capacity.

Marks, James G., Jr.; Trautlein, Joseph J.; Zwillich, Clifford W.; Demers, Laurence M. (1984). Contact Urticaria and Airway Obstruction From Carbonless Copy Paper. Journal of the American Medical Association, pp. 1038-1040; August 1984.

A 27-year-old woman experienced pruritus, eye and throat irritation, hoarseness, shortness of breath, and fatigue within half an hour of exposure to carbonless copy paper. On two separate occasions, she was purposely challenged in a controlled-blinded fashion with portions of the carbonless copy paper. This resulted on both occasions in contact urticaria of the hand that held the paper and changes in pulmonary function flow-volume loops characteristic of upper airway obstruction. It was concluded that the cutaneous and respiratory symptoms induced by the carbonless copy paper were probably related to prostaglandin release.

Matas, M; el-Guebaly, N; Harper, D.; Green, M.; Peterkin, A. (1986). Mental Illness and the Media: Part II. Content Analysis of Press Coverage of Mental Health Topics. Can. J. Psychiatry, Volume 31, June 1986; pp 431 - 433.

The public image of psychiatry has been tarnished in recent years. In order to determine the extent to which press coverage has contributed to negative attitudes towards psychiatry, a content analysis was conducted of a random selection of newspaper articles which appeared over a twenty-

year period in two different newspapers. It was found that although there had been some minor changes over the years, such as more appropriate headlines and more direct quotes from psychiatric experts, on the whole, content and attitudes had changed very little. An accuracy check of media reporting of forensic cases over a 20-year period revealed that when reporters have access to written material, the accuracy levels are greatly improved.

Maulfair, Conrad G., Jr. (1987). Letter to Bruce M. Small & Associates Limited, re Study on Healthy Environments. unpublished.

The single most controllable ecological irritant is chemical. Although the immune system can tolerate and adapt to chemical exposures to a point, that point is often exceeded, and the immune system fails, resulting in symptoms and disease.

May, Charles D. (1976). Objective Clinical and Laboratory Studies of Immediate Hypersensitivity Reactions to Foods in Asthmatic Children. *J. Allergy Clin. Immunol.*, Vol. 58, No. 4, pp. 500-515; October 1976.

Keywords: food hypersensitivity; asthma; children;

Mayhew, D.R.; Donelson, A.C.; Beirness, D.J.; Simpson, H.M. (1986). Youth, Alcohol and Relative Risk of Crash Involvement. *Accid. Anal. & Prev.*, Vol. 18, No. 4, pp 273 - 287; 1986.

Deaths and injuries due to road-crash involvement are a major health and safety problem, especially among youth. Numerous factors can account for the overrepresentation of young drivers in road crashes and one of these — alcohol — has received renewed attention. This paper examines evidence pertaining to drinking and driving among youth to determine the extent to which alcohol has special significance for crashes involving young drivers. Findings show that frequent and heavy alcohol consumption among teenagers and young adults is not unusual, although they are less likely than older age groups to drive after drinking. Nonetheless, those young people who drive after drinking have a greater risk of crash involvement than older drinking drivers at all blood alcohol concentrations. Further investigation into the personal and social characteristics of young people who drive after drinking would provide a sounder empirical basis for policies and programmes to reduce crash involvement among youth.

McCauley, Gary (1987). Letter to Bruce M. Small and Associates Limited re Healthy Environments. unpublished; September, 1987.

Although surveys show Canadians place the issue of child health high on the national agenda, people at the Canadian Institute of Child Health often feel that children are Canada's forgotten people.

McCluskey-Fawcett, Kathleen A; Meck, Nancy; Harris, Marybeth (1986). Prevention During Prenatal and Infant Development: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 43 - 73.

Because of the great vulnerability of the embryo and fetus to environmental assault, the prenatal period is the most critical stage of human development. Many adverse outcomes during gestation can be prevented by good obstetrical management and maternal compliance. Prenatal examples discussed: in the area of maternal disease such as diabetes and sexually transmitted disease; in the area of maternal habits such as social, prescription and illicit drug use; and in the area of maternal characteristics such as age. Problems during the birth period discussed: analgesia and anesthesia, Casearean delivery, home delivery. Postnatal prevention issues discussed: Down's Syndrome, Spina Bifida.

Improved prenatal care and expanding medical technology have permitted babies to be born and kept alive that would never have survived two decades ago. This achievement has not been matched in the area of decreasing morbidity, such that more infants may survive with severe handicaps than in previous years. In addition, actual practice of preventive services is not widespread and the whole issue of cost/benefit is not clearly resolved.

McDonald, J. Corbett (1984). Investigation of Employee Health Complaints at Les Terrasses de la Chaudiere: Final Report to TB/PSAC Steering Committee. Treasury Board of Canada, July 1984.

Research is described into the extent and nature of Federal employee health complaints at a three-tower government office complex opened in 1977. The investigation had three components; a questionnaire survey of a random sample of employees, a telephone inquiry of a selected group of employees, and an environmental survey on selected floors and individual work locations. It was found that most of the employees had, since moving into the building, suffered from upper respiratory tract and eye irritation, together with a variety of other complaints — typically headache, drowsiness, exhaustion, sleeplessness and irritability — and sometimes skin dryness and irritation. People who work in cubicles tend to suffer more frequently than those in open areas or closed offices. The investigation failed to identify the cause. However, some floors and work location had less than optimal ventilation and control of temperature and humidity, partially due to the fact that the building was designed for predominantly open plan use and ventilated accordingly but was partitioned or divided into closed spaces which interferes with air movement.

McDowall, M.E. (1986). Mortality of Persons Resident in the Vicinity of Electricity Transmission Facilities. *British Journal of Cancer*, Volume 53, Number 2, 1986; pp. 271-279.

The study examined whether the electricity transmission system presents a long term hazard to public health, by analyzing mortality data among nearly 8,000 persons who lived in the vicinity of transmission lines. Overall mortality was lower than expected and no evidence of major health hazards emerged. The study did not support previously reported associations of exposure to electro-magnetic fields with acute myeloid leukemia, other lymphatic cancers, and suicide. The only statistically significant excess mortality was for lung cancer (in women, overall, and in persons living closest to the sub-stations); this result is difficult to interpret in the absence of smoking data, and is not supported by other evidence but does not appear to be due to the social class distribution of the study group. The idea that electromagnetic fields might act as weak cancer promoters, rather than initiators, is discussed.

McGee, Charles T. (1979). Health, Light, and the Electromagnetic Spectrum. *How To Survive Modern Technology*, chapter 5, pp. 108-116; Keats Publishing, Inc., 1979.

Studies by researchers such as John Ott show that electromagnetic radiation can have negative effects on both plant life and people. Human muscle strength is weakened under exposure to light that lacks part of the natural spectrum. Classroom experiments indicate that student performance, behaviour and hyperactivity improve when regular fluorescent lights are replaced with full spectrum fluorescent lights. Microwaves & X rays have also been discovered to be health hazards.

McGrath, T.; Paolini, R.; Wright, G.R.; Kusiak, R. (1987). Smoking Survey of Ministry of Labour Employees. *Occupational Health in Ontario*, Vol. 8, No. 3, pp 102 - 111; Summer, 1987; Ontario Ministry of Labour, Occupational Health Branch, Toronto Canada.

The results of a survey of Ministry of Labour employees' viewpoints on smoking within their workplace are presented. Employees were of the opinion that smoking should be restricted to designated areas that are ventilated so that exposure to secondhand smoke generated within Ministry premises does not occur.

McGregor, R.G.; Vasudev, P.; Letourneau, E.G.; McCullough, R.S.; Pranti, F.A.; Taniguchi, H. (1980). Background Concentrations of Radon and Radon Daughters in Canadian Homes. *Health Physics*, Vol. 39, pp. 285-289; August 1980; Pergamon Press Ltd.

Keywords: radon; residential; Canadian homes; health effects;

McKillop Farlow, D'Arcy (1987). Achieving Health For All: The Epp Report. Healthsharing, pp 9 - 10; Summer, 1987.

The Epp Report, Achieving Health For All, indicates some government movement to incorporate health promotion into overall health strategies and may provide a stepping stone to increasing budgets for health promotion activities. But it overemphasizes, according to this researcher, individual self-care without recognition of the polluted society within which we live. This emphasis may undercut the value of self-care models and deflect attention away from environment and socio-economic causes of ill health. It also pushes people to assume impossible levels of individual responsibility without recognizing that they have decreasing control over the environment in which they live.

McKnight, John L. (1986). Regenerating Community. From Consumer to Citizen: Building a Framework for Support; Chapter 2, pp 13 - 22; Canadian Mental Health Association; May, 1986.

Community represents unique social tools that are unlike the social tool represented by a managed institution. For example, the structure of institutions is a design established to create control of people. On the other hand, the structure of associations is the result of people acting through consent. The roles of citizen and community are often traded in for the right to clienthood and consumer status. While we have reached the limits of institutional problem solving, we are only at the beginning of exploring the possibility of a new vision for community. This is a vision of regeneration, a vision of reassociating the exiled people who are the mentally disabled.

McLaughlin, Don E.; Kagen, Herbert P. (1974). The Effects of Sulfur Dioxide on Learning and Activity in the Rat: Psychology and Education Section. Proceedings of the West Virginia Academy of Science-CISTI, Vol. 45, No. 4 pp. 439-444; 1974.

The investigation was an attempt to ascertain the behavioural effects of chronic exposure to sulfur dioxide, a major atmospheric pollutant. The behaviours of interest were learning in simple and complex learning tasks and activity. In addition, a progress record of each subject's weight was kept. The subjects consisted of forty-three male rats which were assigned to one of two control conditions or one of four experimental conditions. The results indicate that sulfur dioxide alone had no effect on performance. There was, however, a significant trials effect and the interaction between SO₂ and the number of trials in the maze also proved to be significant. Activity was found to be increased as the SO₂ concentrations increased. Finally, it was found that the rats living in the higher concentrations tended to be heavier than those in the lower concentrations.

McLeod, Linda (1987). Battered But Not Beaten: Preventing Wife Battering in Canada. Canadian Advisory Council on the Status of Women; June 1987.

Almost 1 million Canadian women from all walks of life are physically, sexually and emotionally abused by their husbands, boyfriends, and former spouses. For these women, this means depression, isolation, physical injury, reduced life options, and sometimes death through homicide or suicide. For the children involved, it means terror, pain, behaviour problems during childhood, the likelihood that the cycle of violence will be repeated in their future families, and a higher risk of becoming violent outside the family. The typical battered woman is trapped in a cycle of poverty from which there has traditionally been little chance of escape. It is recommended that the government increase funding for shelters and staff trained to deal with the problem.

McVittie, Douglas J. (1986). The Control of Occupational Health Hazards in Construction. Occupational Health in Ontario, Vol. 7, No. 4, pp 205 - 212; Fall, 1986; Ontario Ministry of Labour, Occupational Health and Safety Division, Toronto ON.

Construction workers may be exposed to a wide variety of toxic materials about which they know very little and over which they have little control. The traditional solutions in fixed industry, using substitution, ventilation, process modification and administrative controls, are inappropriate for construction sites, and control by procedure is more appropriate. The detailed and uniform specifications make it easy for regulatory agencies to assess compliance.

McWhinney, Ian R. (1987). Fine Words, But Will He Deliver?. CMAJ, Vol. 136, pp 473 - 474; March 1987.

The "Achieving Health for All" report by Minister of National Health and Welfare Jake Epp is welcome but does not convey a sense of the massive changes that will have to occur. Regional and social class inequities, as well as health problems due to substance abuse, industrial hazards, environmental pollution, unemployment and social isolation, are challenges for public policy. Although they are very important for medicine and other health care professions, the record to date is not very good. "Health for All" is impossible because as our environment is constantly changing, so are the challenges to health.

Government policies work against the policy in many ways. For instance, family medicine holds the most potential for health promotion, but there is little economic incentive for physicians to practise it. Most of the financial rewards go to those who have procedures to perform, either diagnostic or therapeutic. There is little incentive for a family physician to work with nurses in health promotion.

The belief that there is a technologic solution for every problem is not covered by the report. This is combined with the belief that if things go wrong somebody is to blame and must pay the price, the fragmentation of the profession, and the declining influence of the generalist in all fields of medicine. These are trends on which government can not be expected to have much influence.

Medical Services Division staff, Workers' Compensation Board (1984).
Occupational Diseases. Workers' Compensation Board, Ontario, Canada; 1984.

This publication is designed as a reference to occupational hazards and diseases which are recognized and for which claims may be allowed under the terms of Ontario legislation. It includes a list by occupation of the hazards to which workers in a given trade may be exposed. In the past, many occupational diseases were named after the specific trades or occupations in which they were first observed. A few diseases still remain quite specific, but many hazards are now so widespread that they may be encountered in jobs in a wide variety of industries. Also listed are particular biological, chemical and physical hazards, their effects on the human body and the occupational or industry groups which may encounter these hazards. Cigarette smoking is discouraged in all occupations due to the possibility of ingestions of toxic substances through the act of smoking.

Meggs, W. (1987). Interview with Dr. William Meggs, Bethesda, MD. Personal communication, July 1987.

In this interview, Dr. Meggs commented that he regularly sees the health and injury consequences of the abuse of illicit drugs and alcohol, as well as the consequences of physical violence. He also cites the widespread use of cigarettes and caffeinated beverages such as coffee as dangerous to public health.

Melia, R.J.W.; Florey, C. du V.; Chinn, S.; Goldstein, B.D.; Brooks, A.G. F.; John, H.H.; Clark, John D.; Craighead, I.B.; Webster, X. (1980). The Relation Between Indoor Air Pollution from Nitrogen Dioxide and Respiratory Illness in Primary Schoolchildren: "Rabka Symposium". Bulletin Europeen de Physiopathologie Respiratoire; Vol. 16, No. 1, pp. 7-8; 1980.

In the UK an association between respiratory illness in primary school-children and the use of gas for cooking has been found in two separate national samples. These findings appeared to be independent of interfering factors such as age, social class and smoking in the home. The association has been found in one large study in the USA, but not in three other studies in the same country. The authors suggest that nitrogen dioxide arising in the emissions of gas combustion might be the cause of the association.

Mendelsohn, Robert S. (1981). *Male Practice: How Doctors Manipulate Women.* Contemporary Books, Inc., Chicago IL; 1981.

Chauvinistic male physicians subject their female patients to medical procedures that are degrading, unnecessary, and often dangerous. So thoroughly institutionalized is medicine's condescending attitude toward women that many doctors are unaware of their negative feelings toward female patients. Thus, they order lab tests and x-rays more indiscriminately for women than for men, over-prescribe for their female patients, treat childbirth almost as a disease, and intervene surgically when sometimes unnecessary, especially with hysterectomies and radical mastectomies.

Mes, Jos; Davies, David J.; Turton, Davida; Sun, Wing-Fung (1986). Levels and Trends of Chlorinated Hydrocarbon Contaminants in the Breast Milk of Canadian Women. *Food Additives and Contaminants*, Volume 3, Number 4, 1986; pp 313 - 322.

A total of 210 breast milk samples from 5 different regions across Canada were analyzed for PCBs and other chlorinated hydrocarbons. There is an apparent increase in PCB residue levels since 1970, an increase should be viewed with caution and may be attributed to better sampling and analytical techniques. There was no evidence for the presence of hexachloro-1,3-butadiene, Mirex, octachlorostyrene, chlorinated naphthalenes, or tetrachlorobenzenes in breast milk, but there were residues of other compounds. Residue levels of breast milk in Canada were similar to those found in other industrial nations. Regional differences in residue levels appeared to be minimal.

Meyer, C. Beat; Hartley, Robert (1981). *Inventory of Current Indoor Air Quality-Related Research.* US Govt.; NTIS PB82-127952; April 1981.

Keywords: pollution monitoring; public health; radon; nitrogen oxides; carbon monoxide; research inventory; formaldehyde; asbestos; particles; indoor air pollution;

Meyer, Beat (1983). *Indoor Air Quality.* Addison-Wesley Publishing Company, Inc.

This author describes some of the factors that determine indoor air quality and reviews the status of knowledge in the field. He supplies a complete bibliography on indoor air quality problems and measurement techniques.

Miksch, R.R.; Hollowell, C.D.; Schmidt, H.E. (1982). Trace Organic Chemical Contaminants in Office Spaces: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1-6, pp. 129-137; 1982; Pergamon Press Ltd., Oxford, Toronto.

Data is presented suggesting that workers in offices are exposed to a broad spectrum of solvent-related organic chemical contaminants present in very low concentrations relative to promulgated or recommended industrial hygiene exposure levels, but in high concentrations relative to outdoor air. With the aid of simple modeling, working hypotheses about various contaminant sources — new and aged building materials, wet-process photocopiers, tobacco smoke, and building maintenance products — are made with respect to the composition, amounts, and generation patterns of their emissions. The results show that effective control strategies can be implemented that do not compromise energy efficiency.

Millar, L.B.; Cooney, P.A. (1982). Urban Lead—A Study of Environmental Lead and its Significance to School Children in the Vicinity of a Major Trunk Road. Atmospheric Environment, Vol. 16, No. 3, pp. 615-620; Pergamon Press Ltd. 1982.

Measurements of children's blood lead levels were ascertained in relation to lead in air and lead in dust attributed to an Inner London arterial highway carrying about 35,000 vehicles per day. The contribution of lead from deteriorating paintwork was also examined. The authors suggest that the findings indicate that lead from vehicles is not contributing measurably to the blood lead levels of children living or being educated in the area.

Millar, Wayne J.; Wigle, Donald T. (1986). Socioeconomic Disparities in Risk Factors for Cardiovascular Disease. Can Med Assoc Journal, Vol. 134, pp 127 - 132; January, 1986.

Despite a general decline in mortality rates in recent decades, these rates are substantially higher among lower socioeconomic groups. To determine target groups for preventive health promotion programs, the prevalence of risk factors for cardiovascular disease by socioeconomic group in Canadian adults aged 20 to 69 years was examined through comparison of estimates from the 1978-79 Canada Health Survey, the 1981 Canada Fitness Survey and the labour force smoking surveys of 1975 and 1983. Level of education was used as a measure of socioeconomic status. The risk factors considered were cigarette smoking, obesity, elevated diastolic blood pressure, physical inactivity, excessive alcohol consumption, elevated serum cholesterol level, diabetes mellitus and the conjoint use of oral contraceptives and cigarettes. The prevalence of the risk factors tended to be higher among men and women with a low level of education. The results were consistent with those of recent Canadian studies showing that both men and women in lower socioeconomic groups are more likely to die from cardiovascular disease.

Miller, C. Arden; Coulter, Elizabeth J.; Schorr, Lisbeth B.; Fine, Amy; Adams-Taylor, Sharon (1985). The World Economic Crisis and the Children; United States Case Study: Special Report on the World Economic Crisis and Health. International Journal of Health Services, Volume 15, No. 1, pp 95 - 135; 1985.

This is a review of the United States experience with issues of child health and services as they relate to changes in economic trends. No existing data systems are entirely adequate for reporting on the current health status of children. An important consideration for the monitoring of children's health is the status of subgroups such as those who are disadvantaged for reasons of poverty, discrimination, or geographic isolation. Ample evidence confirms that children living in poverty suffer adverse health consequences and the proportion of children living in poverty has increased steadily since 1975. When either local or widespread economic reversals are anticipated, health services and social supports for children need to be expanded rather than contracted.

Miller, Claudia S. (1979). Mass Psychogenic Illness or Chemically-Induced Hypersusceptibility?. U.S. Dept. of Health, Education, and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health; 1979.

There are literally millions of workers who are wondering why they never seem to feel good, why they are irritable when they get home, sleepy all day long, and exhausted when they get up in the morning. They are not hypochondriacs or hysterics. Most are simply susceptible to chemicals with which they are working, or to which they are exposed. It has been shown that so-called subjective symptoms of mass hysteria are identical in every respect to symptoms experienced by persons tested for susceptibility to various petrochemicals. It can be objectively demonstrated by testing affected individuals with suspected chemical incitants.

Miller, Joseph B. (1978). Hidden Food Ingredients, Chemical Food Additives and Incomplete Food Labels. Annals of Allergy, Vol. 41, No. 2, pp. 93-98; August 1978.

Keywords: food additives; product labelling;

Miller, Suzanne M. (1980). Why Having Control Reduces Stress: If I Can Stop the Roller Coaster, I Don't Want to Get Off. Human Helplessness; Academic Press; Garber, J., Seligman, Martin E.P., editors; pp 71 - 95; 1980.

Having control reduces the stress generated by an aversive event by providing individuals with an guaranteed upper limit or by enabling them to match their internal state with external events.

Moch-Sibony, A. (1981). The Effects of Prolonged Exposure to Noise on Certain Psycho Motor Intellectual and Personality Aspects of Children—Comparison Between A Soundproof and a Nonsoundproof School. *Travail Humain*, Vol. 44, No. 1, pp. 169-178; 1981.

The effects were studied of prolonged exposition to noise on certain psychomotor, intellectual and personality aspects of children. The authors present the results obtained in a field study undertaken in a zone highly exposed to aviation noise. A comparison made between a sound-proofed and a non sound-proofed school brought out the following facts: the children did not express any feeling of disturbances as to the frequency of airplane passage; yet, after prolonged exposure to the auditory stimulus, perturbations of certain intellectual, psychomotor and personality aspects were observed. The authors estimate that the numerous airplane passages are at the origin of attentional troubles which would explain to a large extent the negative effects observed.

Mohr, Richard D. (1985). Invisible Minorities, Civil Rights, Democracy: Three Arguments for Gay Rights. *The Philosophical Forum*, Vol. XVII, No. 1, pp. 1-24; Fall 1985.

The author puts forth an exacting which establishes that civil rights for lesbians and gay men as necessary preconditions for lesbians and gay men having equitable access to civic rights, i.e. the rights to the impartial administration of civil and criminal laws in defense of property and person. An invisible minority historically subjected to widespread social discrimination has reasonably guaranteed access to these rights only when the minority is guaranteed non-discrimination in employment, housing, and public services. He notes further that a society which forces minority views into invisibility, loses the ability to scrutinize and bring about repeal of undesirable legislation, thereby seriously curtailing the operation of those political processes ordinarily to be relied upon to protect minorities. The social prejudice against gays virtually eclipses their political rights, including freedom of speech, freedom of press, freedom of assembly, freedom to petition for the redress of grievances, and freedom to join with and be identified with other persons for common political goals. Virtually all the rights discusses involve public actions, which are impossible for a person who must remain invisible, hidden and secreted, in respect of his or her minority status, as a condition for maintaining his or her livelihood.

Molhave, L. (1979). Indoor Air Pollution Due to Building Materials: from Indoor Climate: Effects on Human Comfort, Performance, and Health in Residential, Commercial, and Light-industry Buildings; Fanger, P.O. and Valbjorn, O.—editors; pp. 89-110. Danish Building Research Institute, Copenhagen; 1979.

Keywords: building materials; indoor air pollution; Danish research;

Molhave, Lars (1982). Indoor Air Pollution Due To Organic Gases and Vapours of Solvents in Building Materials: from Indoor Air Pollution; Spengler, John—editor; pp. 117-127. Environment International, vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

The emission of organic gases and vapours of solvent type from 42 commonly used building materials was measured under standard conditions. An average of 22 compounds was found in the air around each building material, and the total concentration of gases and vapours was from 0.01 to 1410 mg/m³. The risks of health effects due to the compounds identified were investigated, and criteria for future air quality standards were discussed. It is concluded that the possibility of negative health effects cannot be neglected, especially not for the more sensitive minority of the general population.

Molhave, Lars; Bach, Bodil; Pedersen, Ole F. (1986). Dose-Response Relation of Volatile Organic Compounds in the Sick Building Syndrome. Clinical Ecology, Volume IV, No. 2, Summer 1986.

A survey of the literature indicates that houses with indoor climate problems have higher concentrations in the air of Volatile Organic Compounds (VOC) than houses without problems. In an attempt to establish a dose-response relationship, this study exposed a group of 62 human subjects in a double blind, controlled experiment to a mixture of 22 volatile organic compounds known to be indoor air pollutants. The subjects were all healthy and without diagnosed asthma, allergy or chronic bronchitis, but claimed to suffer from dry mucous membranes in eyes, nose or upper airways. A significant effect of exposure was found in relation to general air quality, odor, ability to concentrate and mucous membrane irritation. Below 0.6 mg/m³ no irritation by VOC can be expected. Above 2 mg/m³ VOC are expected to cause irritation. Above 5 mg/m³ irritation and decreased mental performance may occur.

Monahan, John T.; Vaux, Alan (1983). Macroenvironment and Community Mental Health. Environmental Variables and the Prevention of Mental Illness, chapter 3, pp 27 - 41; Lexington Books; 1983.

Aspects of the physical environment can create serious mental health problems. Those which have received the most research attention have been noise and density. In addition, economic status, unemployment, and economic change have been found to have an effect on community mental health variables. This research should lead community mental health professionals to pay greater attention to physical aspects of their own particular areas, including noise, crowding, architectural design, as well as community economy, particularly economic downturns, plant closings, and local unemployment rates. Such information should allow both the anticipation of service needs and the development of methods for preventing psychosocial problems and for enhancing human functioning.

Whenever environmental stressors may be prevented or diminished, mental health professionals should make every effort to do so. Participation in the environmental impact assessment process is one likely medium for such activity. When the macroenvironmental stressors are not preventable, the major role of the community mental health professional is to mitigate their effects through such techniques as anticipatory guidance and attribution therapy, carried out individually and through the media.

Moschandreas, D.J.; Rector, H.E. (1982). Indoor Radon Concentrations: from Indoor Air Pollution; Spengler, John—editor; pp. 77-82. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

The indoor air of 60 residences in and around a Maryland suburb of Washington DC was monitored in a pilot study to determine residential radon concentrations. In each residence, a radon grab sample was acquired in the living room, and if possible, in the basement. Over 60% of the residences sampled show air infiltration rates below 0.6 air changes per hour. Approximately 55% of all surveyed basements and 30% of all surveyed living areas displayed radon concentrations in excess of 4.0 nCi m⁻³. Assuming an equilibrium factor of 0.5, these radon levels may lead to working levels above the annual guidelines suggested by EPA for Florida homes built on land reclaimed from phosphate mining.

Moschandreas, D. (1986). Formaldehyde: Sources, Methods of Analysis, Exposure and Health Effects. Indoor Air, Vol. 6, 1986; Swedish Council for Building Research; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

Elevated formaldehyde concentrations, higher than 100 ppb, are measured in many indoor environments. They relate positively with temperature and decrease with the age of the source. Urea formaldehyde bonded products can be improved to such a degree that indoor levels can be reduced to ambient levels if the materials are properly installed and used. Formaldehyde is not a potent carcinogen for humans. Pulmonary functions are similar in control and sample populations. Nasal and skin symptoms are higher in populations exposed to elevated formaldehyde concentrations.

Mueller Associates (1981). Wood Combustion: State-of-knowledge Survey of Environmental, Health, and Safety Aspects. US Govt.; NTIS DE83005070; October 1981.

Keywords: wood combustion; air pollution; sulfur dioxide; nitrogen oxides; hydrocarbons; carbon monoxide;

Muittari, Antero; Veneskoski, Tapani (1978). Natural and Synthetic Fibers as Causes of Asthma and Rhinitis: "a communication from Finland". Annals of Allergy, Vol. 41, pp. 48-50; July 1978.

Keywords: asthma; rhinitis; fibers; indoor air pollution;

Murray, T. Keith (1987). Diet and Health: The Canadian Experience. Am J Clin Nutr, No. 45; pp 1390 - 1393; 1987.

The development of Canada's food selection guidelines is described and perceived errors in the process are noted. The guidelines were derived from the report of an expert committee, convened, not to develop guidelines, but to advise the government on the relationship between diet and cardiovascular disease and to identify the advice that should be given to the public. The request for such a study came from the margarine industry, which wanted to know what health claims could be made in advertisements. The committee contained no nutritionists nor educators. Dietary guidelines should be readily understandable by the public in order for them to be used fully, meaning that guidelines should be expressed in terms of food.

Mustard, Fraser J. (1987). Achieving Health for All: Implications for Canadian Health and Social Policies. MCAJ, Vol. 136; pp 471 - 473; March 1987.

The broad philosophical concepts of health have swung in emphasis over the centuries between the important role of social and economic factors and individual choice in determining health status and the effectiveness of determining the cause of disease and developing specific therapeutic measures. Based on the government's policy of "Health for All", the individual choice concept (or the ecology of health) is again becoming paramount, with implications for the role of medicine in society and for groups of health care professionals. For example, achieving a solution to the first of the paper's health challenges, to reducing inequities in the health of low versus high income groups, is a formidable socioeconomic task which requires communication between those concerned with our health and social problems and those concerned with our economic future.

Response to these challenges has to be met with limited resources; thus, there are implications for the level and distribution of funding among the different sectors responsible for health care. If too rapid, the changes necessitated by the emergence of the ecology of health as the dominant theory will be complex and at times will create conflict. One change already evident is that more and more physicians are being placed within managed health care services, with resultant impact on their professional freedom.

Nagda, Niren L.; Rector, Harry E. (1983). Guidelines for Monitoring Indoor Air Quality. US Govt.; NTIS PB83-264465; September 1983.

This document provides guidelines for designing programs to measure indoor air quality. Brief summaries of past and current research and descriptions of indoor contaminants provide a background for developing the monitoring design. Factors that influence air quality are discussed with the aid of mass balance models. An extensive review of measurement systems, including a listing of numerous instruments with their performance specifications, is presented.

Nagira, T.; Hisashige, A.; Kume, Y.; Ueno, M.; Yamamoto, M.; Aoyama, H.; Kurumatani, N. (1981). Health Hazards Among Children in Air Polluted Districts, Report 2: Exposure to Automobile Exhaust and Prevalence of Subjective Symptoms Among School Children. *Nippon Eiseigaku Zasshi*, Vol. 36, No. 3, pp. 596-612; 1981.

Recent rapid motorization in Japan has posed the serious problem of air pollution caused by automobile exhaust. Route 43 has the highest congestions of vehicular traffic in Japan and runs through densely residential areas. Children in schools located along this highway have been found to have increasing rates of respiratory disorders. NO₂ concentrations near Route 43 have exceeded the short-term exposure limit (0.10-0.17 ppm) proposed as the guideline for the protection of public health by WHO Task Group, which also recommended that this limit be stricter for sensitive subjects such as children. In 1979, the authors polled children concerning respiratory, rhinopharyngeal and eye disorders in two elementary schools facing the highway and one school in a non-polluted area, using a self-administered questionnaire. Results were as follows: children attending the two polluted schools had higher rates of disorders relating to bronchial asthma, recurrent respiratory infection and allergies; a close relationship was found between the prevalence of such conditions and distance from the highway and this gradient corresponded to the level of nitrogen oxides; air conditioners furnished at schools appeared to have some remedying effect on these disorders; children who lived south of the highway had higher rates of disease due to higher concentration of exhaust carried at night by northerly winds.

Nathanson, Constance A. (1975). Illness and the Feminine Role; A Theoretical Review. *Soc Sci Med*, Vol 9, No 2; Pergamon Press; 1975.

Women in the Western world live longer than men and have lower mortality rates for most causes of death. There is strong evidence that these differences are due to women's constitutionally greater resistance to infectious and degenerative disease. However, in sharp contrast to their favorable mortality, women report more physical and mental illness than men, and utilize health services at substantially higher rates. These latter differences are documented with health survey data from the United States and Great Britain. Three explanatory models to account for sex differences in illness experience are considered: 1) women report more illness than men because it is culturally more acceptable for them to be ill; 2) the sick role is more compatible with women's other role responsibilities; and 3) women have more illness than men because their assigned social roles are more stressful. Evaluation of these alternative models is based on a review of data from studies of variations in illness among women, and it is concluded that a model looking at illness behaviour as a function of the number and character of other role obligations offers the most promise for future research.

National Commission for the Protection of Human Subjects of Biomedical and Behavioral Research (1977). Research Involving Children: Deliberations and Conclusions: Report and Recommendations. US Government, DHEW; 1977; pp 123 - 154.

The Commission identified three ethical principles that should underlie the conduct of biomedical and behavioral research involving human subjects: beneficence, respect for persons, and justice. In the case of research involving children, the challenge is to find a proper balance in applying these principles and to establish priorities among the principles when they appear to be in conflict. In order to promote the health of both children and adults, the participation of children in research is needed but those who conduct or sponsor such research must protect children from harm by limiting the risk to which they may be exposed as research subjects. Guidelines are given for research which presents more than minimal risk but no immediate prospect of direct benefit to the individual children involved. The threat of an epidemic that could be offset by developing a safe and effective vaccine might justify research involving risk greater than otherwise acceptable to establish safety, efficacy and dosage levels for children of different ages.

The incapacity of children to consent to involvement in research is one aspect of a more general condition of dependence on adults who are responsible for their care. There must, however, be respect for the child's developing autonomy. Any child capable of some degree of understanding (generally, a child of seven or older) should participate in research only if he or she assents. When parental permission cannot be relied upon as a protective mechanism, alternative mechanisms should be set in place to protect the health and welfare of the children.

National Program to Reduce Tobacco Use, Consultation, Planning and Implementation Committee (1987). Directional Paper of the National Program to Reduce Tobacco Use in Canada: Break Free. National Program to Reduce Tobacco Use; June 1987.

This directional paper outlines a national strategy aimed at achieving a generation of non-smokers in Canada by the year 2000. It was developed as a cooperative, collaborative effort of the federal, provincial, and territorial governments and national health organizations. Three goals are: 1) to protect the health and rights of non-smokers; 2) to help non-smokers to stay smoke-free; and 3) to encourage and help those who want to quit smoking to do so. For each goal, quantified short and long term objectives have been developed. These objectives ensure that both individual and environmental perspectives of the problem will be addressed. The following strategic directions have been identified: legislation; access to information; availability of services and programs; message promotion; support for citizen action; intersectoral policy coordination; and research and knowledge development.

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

Tobacco use represents the number one preventable health issue in Canada. More than 30,000 Canadians die each year from smoking-related illnesses. While overall tobacco use has declined in Canada, a number of disturbing trends persist: smoking onset among 12 to 14 year olds has remained high; the prevalence of smoking among young women (20 to 29) is increasing; and regional and socioeconomic differences in smoking rates continue to exist.

McNall, Preston E. (1975). Practical Methods of Reducing Airborne Contaminants in Interior Spaces. Arch Environ Health, Vol. 30, pp. 552-556; Nov. 1975.

Keywords: indoor air pollution;

Needleman, Herbert L; Gunnoe, Charles; Leviton, Alan; Reed, Robert; Peresie, Henry; Maher, Cornelius; Barrett, Peter (1979). Deficits in Psychologic and Classroom Performance of Children With Elevated Dentine Lead Levels. The New England Journal of Medicine, Vol. 300, No. 13, pp. 689-695; Mass. Medical Society; March 29, 1979.

To measure the neuropsychologic effects of unidentified childhood exposure to lead, the performance of 58 children with high and 100 low dentine lead levels was compared. Children with high lead levels scored significantly less well on the Wechsler intelligence Scale for Children than those with low lead levels. This difference was also apparent on verbal subtests, on three other measures of auditory or speech processing and on a measure of attention. Analysis of variance showed that none of these differences could be explained by any of the 39 other variables studied. Also evaluated by a teachers' questionnaire was the classroom behavior of all 2146 children whose teeth were analyzed. The frequency of non-adaptive classroom behavior increased in a dose-related fashion to dentine lead level. Lead exposure, at doses below those producing symptoms severe enough to be diagnosed clinically, appears to be associated with neuropsychologic deficits that may interfere with classroom performance.

Needleman, Herbert L. (1983). Lead at Low Dose and the Behaviour of Children. NeuroToxicology, Vol. 4, No. 3, pp. 121-133; 1983.

Lead can be a useful paradigm with which to deepen our understanding of the health significance of hazardous substances in general, and to clarify the necessary steps and barriers to their removal from the human environment. Epidemiologic data demonstrate a clear effect of lead at low doses. When taken with animal studies which show disturbed learning at low dose in the rodent and non-human primate, altered development of synapses in the developing rodent brain and delayed appearance of brain cytochromes in the immature rodent, the case seems strong that lead at low doses is an important and widely distributed neurotoxin.

Nero, A.V. (1986). Radon in Dwellings; Exposure and Risk Analysis. Indoor Air, Vol. 6, 1986; Swedish Council for Building Research; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

A very broad range of concentrations of radon is observed inside buildings, particularly in homes, and even the average concentration has an estimated risk that is large by comparison with most environmental risks. There is estimated to be a chance of about one in a thousand for the average member of the population to contract lung cancer from indoor radon daughters. And significant numbers of people in many countries receive much higher exposures, corresponding to individual risks of lung cancer exceeding one percent. Control measures take the following pattern, at least in homes with high levels; source control has the greatest influence, ventilation has lesser influence, and air cleaning has questionable effect. Considerable progress is being made in developing strategies to locate and reduce excessive concentrations in existing residences and to prevent such levels in new buildings. These include guidelines for allocation of responsibility and for acceptability of indoor levels, as well as specification of means for identification and control of cases that may have concentrations.

Nethercott, James R. (1982). Health Effects of Formaldehyde. Corpus.

This paper describes the health effects of formaldehyde. These include irritation to the skin, eye, lung and mucosal surfaces; sensitization; alteration in irritancy threshold; mutagenicity; carcinogenicity. Although some of these effects have not yet been proven in humans, the author feels that it is prudent to regard formaldehyde as a possible cause because of animal data. Detailed studies of persons who believe they are being intoxicated by low level formaldehyde exposure is necessary.

Neus, Hermann; Godderz, Werner; Otten, Heinz; Ruddel, Heinz; von Eiff, August-Wilhelm (1985). Family History of Hypertension and Cardiovascular Reactivity to Mental Stress — Effects of Stimulus Intensity and Environment. Journal of Hypertension, Vol. 3, No. 1, pp 31 - 37; Gower Medical Publishing Ltd.; 1985.

This study examines the hypothesis that normotensives with hypertensive parents exhibit enhanced cardiovascular reactions to mental stress. Normally healthy male subjects were examined and, under laboratory conditions, administered an intensified standard stress test. Blood pressure, heart rate and stroke volume were measured. Subjects with at least one hypertensive parent exhibited enhanced responses of systolic and diastolic blood pressure and heart rate. The conclusion was made that cardiovascular hyper-reactivity in subjects with hypertensive parents is only apparent using sufficiently intense stimuli but there are familial trends in cardiovascular reactivity to mental stress.

New York Academy of Medicine (1981). Proceedings of the Symposium on Health Aspects of Indoor Air Pollution. Bulletin of the New York Academy of Medicine, Vol. 57, No. 10; December 1981.

Keywords: indoor air pollution; conference proceedings;

Newball, Harold H.; Brahim, Sami A. (1976). Respiratory Response to Domestic Fibrous Glass Exposure. Environmental Research, No. 12, pp. 201-207; 1976; Academic Press, Inc.

Keywords: fibrous glass exposure; respiratory response; health hazards;

Nieder, Joseph (1986). Evaluation and Treatment of Allergic Disorders in Children: A Psychiatric Perspective. Psychobiological Aspects of Allergic Disorders; chapter 10, pp 233 - 249; Praeger Publishers; 1986; Young, Stuart H., Rubin, James M., Daman, Harlan R., ed.

There is a complex interplay between asthmatic symptoms and emotional factors. Emotional factors can trigger or worsen asthma in a physiologically or biochemically predisposed individual. The disease process has a very definite emotional toll on the individual with asthma, as well as on other members of the family. The adaptation to illness, the alterations in lifestyle, and the family and personal interactions, all require complex psychological adaptations if one is to achieve the optimal degree of functioning with one's illness. Emotional factors and psychological problems which accompany or result from having asthma can be a major impediment to progress toward health.

Nietzel, Michael T.; Himelein, Melissa J. (1986). Prevention of Crime and Delinquency: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 195 - 221.

Five areas are offered as crime-prevention basics. These are reduction of domestic violence; enhancement of various cognitive and behavioral competencies in children, adolescents and young adults; development of effective discipline techniques within families; promotion of safer environments and protected victims; and diversion of predelinquents and delinquents from formal contact with the criminal justice system. Citizens groups should be organized because fear of crime is greatest in communities that lack the power to regulate themselves or perceive that they lack such power.

Nitta, H.; Maeda, K. (1982). Personal Exposure Monitoring of Nitrogen Dioxide: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 243-248; 1982; Pergamon Press, Oxford, Toronto.

Measurement of personal exposure to nitrogen dioxide for short and long term was made with a sensitive NO₂ passive sampler to volunteer housewives and office workers in different seasons. These measurements were compared with the simultaneous measurement of outdoor and indoor concentration of the participants. A common result over all the measurements is the potential effect of using an unvented space heater to increase personal exposure. Mean personal exposure and indoor concentration are higher than outdoor levels elevated by the samples exposed to pollutant produced from the heater. A time-weighted indoor/outdoor activity model gives modestly improved estimates of personal exposure over those predicted from measured indoor concentrations alone.

Norris-Baker, Carolyn (1985). Perceived Importance of Community Settings in Daily Lives of Frail Elderly in Small Towns. *Environmental Change/Social Change*; EDRA 16; pp 216 - 222; Environmental Design Research Association; 1985.

A substantial number of older people living in rural areas experience the compound problems of aging and coping with physical disabilities, yet they continue to reside independently or with spouses in the community. For these frail individuals, the social and physical environment plays a vital role in maintaining the balance between their needs for autonomy and their needs for security. The present research explored the roles of attachment to and participation in small town settings by the frail elderly. Descriptions of settings that frail persons perceived as important in daily life, and their participation in settings in the community, were compared with the perceptions and participation of healthier, more active older persons. Findings indicate that the settings perceived as important in older peoples' lives varied widely and included many nonservice settings such as homes and social groups. The frail and active groups perceived similar numbers of settings as important, but the frail elderly were more limited in their use of those settings, particularly those that were more public and service-oriented. The frail elderly also identified a number of less accessible and less public settings which were important in their lives, but never frequented. When participation in community settings was examined within the context of personal characteristics such as age and health, findings suggested that the perceived importance of settings, and particularly their publicness, does play a modest role in community involvement. An extensive bibliography is included.

Novick, Marvyn (1986). Work and Well-Being: Social Choices for a Healthy Society: A New Work Agenda for Canada. Canadian Mental Health Association, Toronto Canada; 1986.

The employment crisis in the next decade threatens to undermine social and economic foundations of family life. A new work agenda is needed. The basic social choice facing Canadians is whether we re-affirm universal employment as essential to a healthy society, or withdraw into the

view that maintenance income is a sufficient condition of basic well-being. The absence of good jobs for those in the 35 to 45 year old category will have a profound impact on personal health and the well-being of families. This will include vulnerabilities to illness and death from acute economic stress, particularly for men in their forties, as well as the re-emergence of child poverty. New approaches to full employment would rest on two key propositions: access to decent jobs remains a primary life aspiration for the vast majority of the adult population; and there are destructive rigidities in present distributions of work among Canadians which have severe health impacts on the unemployed, and are leading to growing stress for many. A new approach would include elements such as special tax rates to cushion annual job hour reductions, shelter policies to stabilize housing costs, national income credits for parents, universal leave programs, enriched pension entitlements, and tuition rebates and interest free loans for students of all ages.

Nussbaum, Karen (1981). Warning: Health Hazards for Office Workers. Working Women Education Fund, 1224 Huron Rd., Cleveland OH 44115; April 1981.

There are many hazards present in a modern office building. These include those resulting from improper ventilation and "tight building syndrome", hazardous office products such as liquid eraser products (which contain trichloroethylene and tetrachloroethylene, known carcinogens) and cleaning solvents (containing benzene, a carcinogen and toluene, a powerful narcotic), ozone from photocopiers, formaldehyde and passive cigarette smoke.

O'Banion, Dan R. (1981). An Ecological and Nutritional Approach to Behavioral Medicine. Charles C Thomas, Springfield IL; 1981.

This book presents a history and analysis of research and descriptive evidence indicating that food, chemical and inhalant substances play a major role in determining human health. The full range of psychological and physical health disorders, including learning disabilities and childhood hyperactivity are related to susceptibility and reactions to ecological and nutritional events. Extensive case reports are included.

O'Banion, Dan R. (1981). The Ecological and Nutritional Treatment of Health Disorders. Charles C Thomas, Springfield IL; 1981.

This book ties together the physiological and behavioral effects, both direct and indirect, of increased sensitivity to various ecological agents including common foods and chemicals used prevalently in our society. Individual susceptibility to specific food, chemical, and inhalant substances may contribute to a wide range of psychological and physical health problems, including learning disabilities. The effects of food and chemical sensitivities, stages of adaptation, levels of reaction, diagnosis and ecological treatment are all discussed.

O'Banion, Dan R. (1981). Skill Development and Maintenance. from The Ecological and Nutritional Treatment of Health Disorders, pp. 168-179; Charles C. Thomas, Springfield IL; 1981.

Many people who are food and chemically sensitive have trouble developing and then maintaining appropriate social and academic skills. When a person is stressed physiologically, learning problems and skill deficits result and typically are treated solely from a psychological standpoint. Behaviors which are related to sensitivities and which affect learning are increased activation, an increased probability of frustration, decreased attention span, perceptual problems, and memory deficits. Most frequently chemically and food sensitive children experience problems in spelling, reading comprehension and auditory memory skills. Deficiencies in these areas may be indicative of malfunctioning neurological systems that predispose the individual to have difficulty learning certain types of tasks. Specific physiological systems may be malfunctioning, causing specific learning deficiencies. If an individual does have skill and learning deficiencies in specific areas due to an inability to attend to stimuli presented in a certain mode, special techniques and testing methods can be used. An improvement in the student's environment, removing the offending substances, will be the best long-term solution.

O'Banion, Dan R. (1981). The Chemical Problem: Chemical Contamination in the Work and School Environments. from An Ecological and Nutritional Approach to Behavioral Medicine, pp 69 -164; Charles C. Thomas, Springfield IL; 1981.

One of the most important settings that influences a child's life is the school environment. This environment should be kept relatively uncontaminated especially for chemically sensitive children whose learning may be greatly affected by reactions to chemicals; however, even less chemically sensitive individuals who appear not to have any dramatic problem learning may be affected by such exposures. Slight fatigue, sleepiness, and uncooperativeness are often produced by chemical exposure and easily corrected by cleaning up the chemical environment in the school setting. Yet, the school environment is one of the most chemically contaminated environments in our society. A wide variety of indoor air pollutant sources in schools is described, including the use of pesticides, motor exhaust, chemicals in clothing, and lighting. How these affect learning and behavior in children is documented.

O'Banion, Dan R.; Peek, Leon A.; Butler, Joel R. (1981). Behaviors Characteristic of Learning Disabilities Related to Specific Foods in an Adolescent: Case Report 4. from An Ecological and Nutritional Approach to Behavioral Medicine, pp. 147-164; Charles C. Thomas Publisher, Springfield IL; 1981.

A fifteen year old male adolescent with an extensive history of behavior and learning disorders had undergone traditional psychological, educational, and medical treatments to no avail. A food testing program in-

volving the ingestion of single foods of organic or less chemically contaminated sources was established to ascertain relationships between specific foods and the individual's physical, educational, and behavioral irregularities. It was found that extreme aggression and hyperactivity each occurred within minutes after the ingestion of a specific food. The food related to the hyperactive state was tested a second time under more closely observed conditions with a resulting decrease in performance on a symbol-digit coding task and the occurrence of a number reversal on one of the coding forms when the reaction was in progress.

O'Brien, I.M., Newman-Taylor, A.J.; Burge, P.S.; Harries, M.G.; Fawcett, I.W. and Pepys, J. (1979). Toluene Di-isocyanate-induced Asthma: II. Inhalation challenge tests and bronchial reactivity studies. *Clinical Allergy*, Vol. 9, pp. 7-15; 1979; Blackwell Scientific Publications.

Keywords: toluene diisocyanate; asthma; health hazards;

O'Hara, Bruce (1987). Work Well: A Guide to Creating Worktime Options Resource Centres. Canadian Mental Health Association.

Rather than having a scarcity of work, our society has a need to find ways to redistribute work to even out the extremes of the overworked and unemployed. This book is intended as a practical manual for promoting new work schedules. An excellent resource section is included, listing useful books, related reading, and contacts.

O'Quinn, Silas E.; Kennedy, C. Barrett (1965). Contact Dermatitis Due to Formaldehyde in Clothing Textiles. *Journal of the American Medical Assoc.*, Vol. 194, No. 6; Nov. 8 1965.

Keywords: formaldehyde; clothing; contact dermatitis; health effects;

Olafsson, Olafur; Svensson, Per-Gunnar (1986). Unemployment-Related Lifestyle Changes and Health Disturbances in Adolescents and Children in the Western Countries. *Soc. Sci. Med.*, Vol. 22, No. 11, pp 1105 - 1113; Pergamon Journals Ltd.; 1986.

According to official statistics, 11 million people under 25 in the 12 OECD member states are unemployed at any given time, a figure which in-depth studies show to be at least 40 to 50% higher. Unemployment hits mainly adolescents, school leavers, young unskilled adults, immigrants, and those who are in need of familial and social support such as the frail, sick, disabled children, and old people. Unemployment is endangering the socio-economic status of people, in spite of short-term unemployment benefits and is creating inequalities in health and serious social misfits.

The majority of young people do not learn to cope with unemployment. It fosters isolation, loss of self-esteem, frustration, and hopelessness. Unemployed school leavers are prone to a destructive lifestyle such as drug, tobacco and alcohol abuse and respond less to health promotion, family planning, hygienic and nutritional programmes. Children and family members of the unemployed suffer from various somatic and psychosomatic disorders. Loss of job or the mere prospect of becoming jobless have been found to cause elevated blood pressure and serum cholesterol, increased concentration of blood catecholamine and elimination of noradrenaline, an increase in the frequency of stress and psychosomatic diseases. After regaining employment, these values normalize. Unemployment is therefore considered as a real source of chronic stress which is considered to be a major contributor to cardiovascular diseases, ulcers, and asthma. The jobless exhibit a significantly higher prevalence of mental disorders, higher admission rates to mental hospitals, parasuicides and depression, and visits to doctors.

Olsen, Jorgen H.; Dossing, Martin (1982). Formaldehyde Induced Symptoms in Day Care Centers. *Am. Ind. Hyg. Assoc. Journal*, Vol. 23, pp. 366-370; May 1982.

A questionnaire study was performed among 70 employees at seven mobile day care centers where urea formaldehyde glued particle board has been used for indoor paneling and among 34 employees at three control institutions selected at random where no particle board has been used as building materials. Responses showed a significantly higher frequency of the following symptoms among the staff at the mobile institutions: mucous membrane irritation, headache, abnormal tiredness, menstrual irregularities and use of analgetics. The median concentration of formaldehyde in the mobile institutions was 0.43 mg/m³ in contrast to a concentration in the control institutions of about 0.08 mg/m³.

Ontario Federation of Labour (1982). *Occupational Health and Safety: A Training Manual*. Copp Clark Pitman, Toronto Canada; 1982.

This manual offers a comprehensive information base, enabling workers to understand their working environment. It covers a wide range of toxic substances and pollution sources, detailing how they affect health. Also covered are stress, material handling, making machinery and equipment safe, principles of ventilation, personal protective equipment and legislation in Ontario.

Ontario Mental Health Foundation (1987). *The Ontario Mental Health Foundation Annual Report*. The Ontario Mental Health Foundation; 1987.

The Foundation's major task is funding research investigations which have a bearing upon mental health and mental disorders. Some projects

include testing ways of teaching social skills to mentally handicapped adults; examining the effects of social support and personality variables on maternal adaptation to the stress of a hearing impaired child; examining the effects of depression in a mother upon the children in a family.

Ontario Ministry of Energy (1987). Airing Out the Sick-Building Syndrome. Ministry of Energy, Municipal and Commercial Programs, Toronto Canada; 1987.

The sick-building syndrome is described with specific reference to Les Terrasses de la Chaudiere in Hull Quebec. General suggestions for improving indoor air quality and for follow-up help and information are included.

Ontario Ministry of Health (1987). Health Effects of Extremely Low Frequency Electromagnetic Fields: A Review of Clinical and Epidemiological Studies. Disease Control and Epidemiology Service, Public Health Branch, Ontario Ministry of Health; July, 1987.

This critical review of the literature was carried out to document the quality of scientific evidence regarding the possible public health risk of installation of extra high voltage line corridors. Literature on human health effects of exposure to extra low frequency fields which are created from many electrical sources, including transmission of electricity at high voltage, was examined. Adverse health effects studied in the reports examined include general health effects such as headaches and dizziness, adverse reproductive outcomes, and cancer. This literature fails to provide conclusive or convincing evidence that there is a significant public health risk associated with ELF fields. General agreement exists among scientific researchers that the ambiguities present in the current literature require clarification. A bibliography is included.

Orehek, J.; Massari, J.P.; Gayraud, P.; Grimaud, C.; Charpin, J. (1976). Effect of Short-Term, Low-Level Nitrogen Dioxide Exposure on Bronchial Sensitivity of Asthmatic Patients. *Journal of Clinical Investigations*, Vol. 57, February 1976, pp 301-307.

A study was undertaken to determine whether exposure to a realistic concentration of nitrogen dioxide could increase the bronchial sensitivity of asthmatic patients to bronchoconstrictor agents. Dose-response curves were established for changes in specific airway resistance in response to aerosolized carbachol in 20 asthmatics after each had spent 1 hour in an exposure chamber breathing both polluted and unpolluted air. Nitrogen dioxide induced a slight but significant increase in specific airway resistance and enhanced the bronchoconstrictor effect of carbachol in 13 subjects. Although the mechanisms underlying the nitrogen dioxide effect remain controversial, the present results demonstrate that very low levels of the substance can adversely affect some asthmatics.

Ostapovich, L.K. (1975). Conditions of Respiratory Route Exposure to Sulfur Dioxide and Formaldehyde and Subsequent Sensitization. *Gigiena I Sanitariia*, No. 2, pp. 9-13; 1975.

The effects of concentration and conditions of exposure on the development of toxic and allergic reactions to sulfur dioxide and formaldehyde were investigated. It was concluded that sulfur dioxide has toxic effects which are manifested as an alteration in the chronaxy ratio of antagonistic muscles, as an increase in whole blood cholinesterase activity, and as changes in blood hemoglobin concentration and erythrocyte counts. Exposure to concentrations of formaldehyde in concentrations of 7 or 2 mg/m³ leads to sensitization. The possibility of an early onset of allergy in response to low doses of the agents in question points to the need for conducting specific tests indicative of allergy—as highly sensitive indicators of adverse effects—in establishing hygienic norms for air pollution.

Ott, John (1973). Health And Light: The Effects of Natural and Artificial Light on Man and Other Living Things. Pocket Books/Simon & Schuster, 1973.

In 1973 a study was conducted by the Environmental Health and Light Research Institute in Sarasota Florida, showing dramatic reactions in hyperactive children. In two first-grade classrooms, the standard cool white fluorescent tubes and fixtures with solid plastic diffusers remained unchanged. The plastic diffusers stopped the transmission of any trace of long-wavelength ultraviolet. In two other classrooms, the tubes were replaced with full-spectrum fluorescent tubes that more closely duplicated natural daylight. By means of hidden time-lapse cameras, student behaviour was observed and recorded. Under the standard cool white lighting, some children demonstrated nervous fatigue, irritability, lapses of attention, and hyperactive behaviour. Within a week of the new lights being installed, a marked improvement in their behaviour appeared and overall classroom performance improved. Experiments are also referred to in which the effect of light and darkness on the pineal gland has been studied, as well as studies concerning low-frequency non-ionizing radiation, hyperactivity and learning disabilities. A connection is also suggested between the hyperactive reaction to radiation from unshielded fluorescent tubes and the same symptoms triggered by artificial food flavours and colourings.

Ott, John; Mayron, Lewis W. (1976). Influence of Fluorescent Lights on Hyperactivity and Learning Disabilities. *Journal of Learning Disabilities*, Vol. 9, No. 7 (Aug/Sep 1976), pp. 417-422.

This study describes in detail the study of the Environmental Health and Light Institute, in which four regular elementary windowless classrooms, two with standard type fluorescent cool-white tubes and two with full spectrum, radiation-shielded fluorescent fixtures,

were compared for their effects on hyperactive behaviour. Time-lapse photography was used to produce a permanent record of the children's actions. The study concluded that cool-white unshielded fluorescent lights could contribute to hyperactive behaviour.

Ott. W.; Flachsbart, P. (1982). Measurement of Carbon Monoxide Concentrations in Indoor and Outdoor Locations Using Personal Exposure Monitors: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 295-304; 1982; Pergamon Press Ltd., Oxford, Toronto.

On 15 dates, 5000 measurements of carbon monoxide were made in downtown commercial settings in 4 California cities, using personal exposure monitoring (PEM) instruments. The data indicate that most commercial settings experience CO concentrations above zero indoors, because CO tends to seep into buildings from vehicular emissions outside. Levels in these locations usually are not above 5 ppm and seldom are higher than the U.S. health-related ambient air quality standards for CO. However, indoor garages and buildings with attached indoor parking areas are exceptions and can experience relatively high CO concentrations.

Ozkaynak, H.; Ryan, P.B.; Allen, G.A.; Turner, W.A. (1982). Indoor Air Quality Modeling; Compartmental Approach with Reactive Chemistry: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1-6, pp. 461-471; 1982; Pergamon Press, Oxford, Toronto.

Data on indoor/outdoor pollutant and tracer concentrations were collected during different periods in 1981 at a residence in Newton MA. Special studies within the kitchen were conducted to determine the vertical and horizontal variability of pollutant and tracer gas concentrations. A reactive chemistry model incorporating simplified NO_x chemistry was developed to simulate pollutant concentrations indoors. Further monitoring and modeling studies to investigate the critical aspects of the short-term dynamics of the reactive pollutants inside homes with gas cooking stoves are recommended.

Ozkaynak, Haluk; Spengler, John D. (1985). Analysis of Health Effects Resulting from Population Exposures to Acid Precipitation Precursors. Environmental Health Perspectives; Vol. 63, pp. 45 - 55; 1985.

Types of available studies relevant to the quantification of air pollution health effects and their principal limitations are discussed. Assessments are provided based on review and re-analysis of previously reported data bases, synthesis of published findings, and original analysis of health data sets using new methods or recent size-specific particle mass measurements. Interim results from ongoing research activities on airborne particle health effects are presented. It is shown that preliminary results obtained from cross-sectional and time-series mortality studies appear to be consistent, indicating that

particulate air pollution, even at current levels, could be of concern for public health. For future air pollution health effects studies, combinations of both improved exposure and health measures are needed. In particular, exposure and response data on acidic aerosols and oxidants involved in the formation of acid precipitation are vitally needed. Also needed are more prospective health studies to enable characterization of chronic and acute health effects. Also needed are better estimates of personal exposures to particles, acid aerosols, ozone, and nitrogen oxides, including information on indoor/outdoor particle exposures by source and chemical composition. Other research priorities are detailed.

Pape, Bonnie; Church, Kathryn (1987). Community Reinvestment: Balancing the Use of Resources to Support People With Mental Disabilities. Canadian Mental Health Association; June, 1987.

Our tendency to segregate mentally disabled persons into expensive formal health care and rehabilitation systems or discharge them with limited support into unprepared communities has created obvious problems for those who administer provincial mental health systems. It is the premise of this paper that in order to resolve this demand/supply dilemma, a strategy which emphasizes a reallocation of existing resources will have to employ community reinvestment as its guiding principle. The idea of community reinvestment promotes the belief that resources can be more effectively used from both an economic and quality of care point of view, by building the capacity of communities to assist mentally disabled persons to live with maximum reliance on the resources found or created within their natural environments, and with minimal dependence on formal service systems.

Paroski, Paul A. (1987). Health Care Delivery and the Concerns of Gay and Lesbian Adolescents. *Journal of Adolescent Health Care*, Vol. 8, pp. 188-192; 1987.

Homosexual adolescents in New York City were questioned concerning their perceived needs and health care requirements. Many had developed a stereotypic view of homosexuality and its associated lifestyle. The study revealed that a homosexual adolescent is often placed at a greater distance from his or her parents than his heterosexual counterpart. For those who reveal their sexual orientation, the family response is often not supportive. In the absence of a support group, the adolescent feels alone and isolated. Both male and females were concerned with receiving non-judgmental health care.

The authors conclude that the typical health care system does not meet the needs of such adolescents for support and guidance. The authors propose that health care providers need to be aware of the sense of isolation, the process of hiding one's homosexuality, and the conflicts that homosexual adolescents have regarding their lifestyle. They should

be knowledgeable about the specific and unique medical and biopsychosocial concerns of the homosexual adolescent. They stress that it is important not to make the assumption that all persons are heterosexual, thereby not allowing for the homosexual adolescent.

Pauls, Jake (1982). Stair Use and Design for Safety. unpublished; January 26 1982.

Keywords: falling accidents; stair safety; building design;

Paulson, George W. (1977). Environmental Effects on the Central Nervous System. Environmental Health Perspectives, Vol. 20, pp 75 - 96; 1977.

The central nervous system (CNS) is designed to respond to the environment and is peculiarly vulnerable to many of the influences found in the environment. Utilizing an anatomical classification, major toxins and stresses are reviewed with selections from recent references. Selective vulnerability of certain areas to particular toxins is apparent at all levels of the CNS, although the amount of damage produced by an noxious agent depends on the age and genetic substrate of the subject. It is apparent that the effect of certain well known environmental toxins such as lead, mercury, etc., deserve continued surveillance. In addition, the overwhelming impact on the CNS of social damages such as trauma, alcohol, and tobacco cannot be ignored by environmentalists. The effect of the hospital and therapeutic environment has become apparent in view of increased awareness of iatrogenic disorders. The need for particular laboratory tests, for example examination of CSF and nerve conduction toxicity studies, is suggested. Epidemics such as recent solvent neuropathies suggest a need for continued animal studies that are chronic, as well as acute, evaluations when predicting the potential toxic effects of industrial compounds. An extensive bibliography is included.

Pearse, L.H.; Crocker, L.H. (1947). The Peckham Experiment: A Study in the Living Structure of Society. Yale University Press, New Haven, Connecticut USA, 1947.

This reference describes an extensive experiment in alternative lifestyles, conducted more than forty years ago. It yields insights as to the difficulties involved in setting up experimental situations for evaluating alternative social environments.

Pedersen, Niels Bang (1980). Occupational Hand Eczema From Formaldehyde in Price Labels. Contact Dermatitis, No. 6, pp. 57-58; 1980; Munksgaard Copenhagen.

Keywords: occupational health and safety; eczema; formaldehyde; health hazards;

Pengelly, L.D.; Goldsmith, C.H.; Kerigan, A.T.; Toplack, S.A.; Furlong, W.J. (1982). The Hamilton Study: Relationships Between Indoor and Outdoor Levels of Sulphur Dioxide, Nitrogen Dioxide and Particulates in Elementary Schools. Air Pollution Control Association, Pittsburgh PA; 1982.

Some preliminary analyses of indoor and outdoor air pollution levels in 12 Hamilton, Ontario schools was carried out. Analysis of the results of SO₂, NO₂ and particulate measurement was carried out to determine the influence of school age and outdoor concentration on the measured indoor level. The design and construction of a building has an influence on the indoor level of NO₂ over a range of outdoor NO₂ levels. We can speculate that the principal factor which influences this relationship is the infiltration or ventilation rate. Newer schools tend to have positive ventilation systems which draw in "fresh air" from the outside; old schools did not have methods for providing for ventilation aside from opening windows.

Pengelly, L. David; Kerigan, Antony T.; Goldsmith, Charles H.; Inman, Elizabeth E. (1983). The Hamilton Study: Distribution of Factors Confounding the Relationship Between Air Quality and Respiratory Health. Air Pollution Control Association; June 1983.

To clarify the role of ambient air quality among the many factors which contribute to chronic lung disease, the respiratory health of a cohort of elementary school children in Hamilton, Ontario was studied. A specific issue in this study was that of the distribution of the covariables of: tobacco smoking in the home, the use of gas for cooking, respiratory symptoms in other family members, respiratory disease in infancy, crowding in the home, income and social/occupational status as well as the distribution of air quality. The study found that the prevalence or magnitude of these covariables was not uniformly distributed among the diverse areas of the city studied, nor were the characteristics of air quality which were measured. In addition, the patterns of prevalence which were found among the areas tended to occur in the same direction, such that the effects of the covariables tended to reinforce each other. For instance, the children studied in the industrial core had the highest percentage of prevalence of all the covariables.

Pepelko, W.E.; Danner, R.M.; Clarke, N.A. (1980). Health Effects of Diesel Engine Emissions: Proceedings of an International Symposium, Volume 11. US Environmental Protection Agency, November 1980.

The purpose of the symposium was to bring together scientists and engineers from the public and private sectors to discuss their research findings on the health effects of diesel engine emissions and to conclude with a discussion of health risk assessment of diesel exhaust. The Proceedings are organized into eight main sections corresponding to the format of the symposium and addressing physical and chemical character-

istics, in vitro carcinogenic and mutagenic effects, biochemical and metabolic effects, toxicological effects of inhaled emissions, mutagenic and carcinogenic potency of extracts of diesel and related environmental emissions, mutagenicity of inhaled diesel emissions, carcinogenic effects of exposure to diesel emissions, epidemiological studies, and lastly a panel discussion on health risk assessment of diesel emissions.

Pepys, J. (1982). Chemical Dusts, Vapours, and Fumes Causing Asthma: special issue of *Environment International*. Indoor Air Pollution; *Environment International*, Vol. 8, No. 1 - 6; pp 321 - 325; 1982.

Allergic respiratory disease due to common allergens of organic origin is well known. Less familiar, but of increasing importance, are chemicals of organic and inorganic nature, met as dusts, vapours, and fumes. Their relevance is shown in occupational respiratory allergic disorders. Confirmation is given by "real life", simulated, occupational-type provocation tests. Controlled exposure with minute amounts for brief periods, thus closely simulating allergic sensitivity, can precisely identify etiological causes in often complex exposures. The capacity of these widely different agents in different forms to elicit the different patterns of asthmatic reactions is a pointed example of their potential role, as well as the role of chemical agents in general. The introduction into indoor environments of such materials demands consideration of their possible allergenic effects and of the need to recognize the various forms of allergic respiratory reaction they may cause.

Petit, Ted L.; Alfano, Dennis P.; LeBoutillier, Janelle C. (1983). Early Lead Exposure and the Hippocampus: A Review and Recent Advances. *NeuroToxicology*, Vol.4, No. 1, pp. 79-94; 1983.

This report summarizes recent experiments conducted both in the authors' laboratory and in others' examining the effects of early lead exposure on the development of a specific portion of the human brain - the hippocampus, and behaviours characteristic of hippocampal dysfunction. Following postnatal lead exposure, marked reductions are seen in general hippocampal development. Observations support the suggesting that the hippocampus may play a critical role in mediating many of the behavioural changes observed following early lead exposure. However, as other brain areas are also clearly effected by lead, alternative explanations for these lead induced behavioural changes are also discussed.

Pfeifer, J.; Richter, J.; Keveova, E.; Kral, V. (1983). Alterations of Selected Indicators of Local Immunity in Children Having Been Exposed to the Action of Formaldehyde of a Long Duration. *Cs. Pediat.*, Vol. 38, No. 5, pp. 274-277; 1983.

A group of schoolchildren who had undergone long-term exposure to formaldehyde as well as an non-exposed control group were

studied. This long-term study (3 years) looked at 159 children. Decreased values of salivary SIgA and of lysozym were repeatedly observed in exposed children and after elimination of the exposure, a prompt normalization of the findings was seen. A statistically significant higher occurrence of abnormal findings in the studied immunity indicators could also be demonstrated. On the basis of data, the authors consider buildings built of prefabricated woodchip board made with the use of a formaldehyde glue to be unfit for the long-term use of children.

Pfeiffer, Carl C. (1979). The Schizophrenias—At Least Three Types: Cerebral Allergy. from *Mental and Elemental Nutrients*, chapter 40, pp. 415-417; Keats Publishing; 1979.

Allergic children can have an allergy-tension-fatigue syndrome which results in disinterest in learning and thus decreases learning ability. Such symptoms as specific learning disabilities, perceptual-motor deficits, coordination deficits, hyperkinesis, emotional lability, short attention span, and abnormal electroencephalograms have been cured or eased by diagnosis and the offending allergen being removed.

Phalen, Robert F.; Reischl, Peter; Faeder, Edward J.; Cavender, Finis L. (1981). Response of the Respiratory Tract to Inhaled Pollutants: from *Aerosols, Airways & Asthma*; Trautlein, Joseph J., MD—editor; pp. 125-139. ? ;1981.

Keywords: aerosols; inhaled pollutants; health effects; air pollution;

Philpott, William H.; Kalita, Dwight K. (1980). Human Ecology and Mental Health. from *Brain Allergies: The Psycho-Nutrient Connection*, chapter 3, pp 15- 27; Keats Publishing, Inc., 1980.

An individual's ability to handle toxins, pollens, foods and chemicals contracted from the environment differs considerably according to his or her unique chemical makeup. The more defective his or her ability, by inheritance, enzyme deficiency, malnutrition, harbored infection or otherwise, the more likely a person is to develop maladaptive symptoms on exposure to food and environmental contacts. A group of 250 emotionally disturbed patients developed major symptoms on exposure to their commonly consumed foods and frequently encountered chemicals. Symptoms included psychosis, schizophrenia, blurred vision, anxiety, dissociation, delusions, headaches, dizziness, inability to read or write, hyperactivity, and a whole range of gastro-intestinal problems.

Philpott, William H.; Philpott, Katherine; Khaleeluddin, Khaja (1981). The Roles of Bio-Ecologic Diagnosis and Treatment of Organic Factors in Mental Disorders. The Institute for Bio-Ecologic Medicine; January 1981.

Increasing pollution of the environment is beginning to be recognized as potential cause of nervous system reactions. Not only do we have these obvious pollutants to which people can react, but of even more basic importance is the observation that maladaptive central nervous system reactions can occur in susceptible people to otherwise nutritious food and/or chemicals.

Pim, Linda R. (1981). *The Invisible Additives: Environmental Contaminants in Our Food.* Doubleday Canada Limited, Toronto Canada; 1981.

Environmental additives are chemicals which inadvertently contaminate our food. Such substances as chlordane (a pesticide residue in milk), aflatoxin (a poisonous mould on peanuts) and sulfonamide (a drug residue in pork) contaminate the food supply and present health hazards.

Planek, Thomas W. (1982). *Home Accidents: A Continuing Social Problem.* *Accid. Anal. & Prev.*, Vol. 14, No. 2, pp. 107-120; 1982.

Keywords: home accidents; residential safety;

Plumlee, Lawrence; Coerr, Stanton (1979). *Panel Discussion: Role of High Risk Groups in the Derivation of Environmental Health Standards.* *Environmental Health Perspectives*, Vol. 29, p. 155-159; 1979.

There are a variety of syndromes which lead to hypersusceptibility, including exposure to high levels of chemicals, nutritional deficiency, and genetic predisposal to environmental chemical sensitivities. In setting standards for environmental contaminants, there must be the awareness that high risk groups are not a small portion of the population, but include virtually everyone from time to time, due to differences in susceptibility for each pollutant, nutritional factors, etc.

Poole, Ron (1987). *Environmental Health Alliance.* unpublished.

An organization, The Environmental Health Alliance, has been set up among local health and environmental organizations. The first task is to address concerns to various levels of government regarding a proposed garbage incinerator in Detroit which will not be equipped with pollution reduction equipment. There is a concern that the toxic, carcinogenic fumes will pollute the air in Windsor.

Poyner, Barry (1980). *Personal Factors in Domestic Accidents: Prevention Through Product and Environmental Design.* British Dept. of Trade, Consumer Safety Unit, Millbank Tower, London SW1P 4QU England; April 1980.

Keywords: residential safety; accidents;

Pressman, Norman E.P. (1982). Priorities for Planning Livable Cities. Creating Livable Cities, pp 1 - 9; Pressman, Norman E.P. ed; 1982.

The improvement of the urban environment is required to remove causes of danger; maximize opportunities for contact between individuals at block, neighbourhood and town levels, both among the residents themselves and with the various types of institutions; add economic value; and enhance the visual and aesthetic qualities of the urban environment as one means of promoting a favourable change in individual psychological behaviour.

Priesnitz, Wendy K. (1987). School Free. Village Books, Unionville ON; 1987.

There is a growing community of families in Canada who chose to help their children learn without attendance at school. These deschoolers value autonomy as the full development of a child's capacity for independent reflection, judgement, decision-making, and action. If autonomy is seen as the link between intellect and responsible action, it cannot be fostered in an atmosphere of coercion. Compulsory attendance laws negate the right to self-reliance and autonomous action. Children suffer greatly from lack of respect and autonomy in schools. The conventional view of children in the educational system is that they are objects to be manipulated. This is apparent in the use of behavioural psychology to create classroom behaviour acceptable to the teacher as well as in the top-down style of curriculum design which views children as empty vessels into which knowledge is poured by the teacher. A non-compulsory, more broadly-based system of community education is proposed in which people of all ages would participate in an on-going, self-generated process.

Priesnitz, Wendy (1987). Moving Away From Assembly-Line Education. Toronto Star, September, 1987.

A severe isolation from real-life experiences in the education system has led to a separation from the fact that learning is an activity which is natural to human beings. Children have ceased to be authentic participants in the life of society and would learn better in an environment free from compulsory schooling and rigid age segregation.

Proctor, N.H.; Hughes, J.P. (1978). Chemical Hazards in the Workplace: The Chemical Hazards: Portland Cement; Propane; Ozone; Formaldehyde; Formic Acid; Arsenic; Ammonia; Carbon Dioxide. J.B. Lippincott Co., Philadelphia PA; 1978.

Keywords: chemical hazards; workplace; portland cement; propane; ozone; formaldehyde; formic acid; arsenic; ammonia; carbon dioxide;

Proshansky, Harold M. (1975). The Environmental Crisis In Dignity. *Journal of Social Issues*, Vol. 29, No. 4; pp 1 - 19; 1975.

The environmental crisis in human dignity lies not just in overuse, the misuse, and the decay of physical settings, but far more significantly in how we conceive of the individual in relation to any such setting. In the design and organization of physical settings, the human properties of the individual are ignored, oversimplified, or implicitly assumed. Spaces and places are improperly designed not only in physical terms; designs overlook human needs for privacy, territoriality and freedom of choice, and the conceiving of the individual as a simple "machine man". Unintended consequences are often ignored and no attempt is made to evaluate just how well the setting actually works. The danger is that the person will adjust and at the price of a continuing erosion of the properties that make him distinctively human. It is imperative that as behavioral scientists turn to the systematic study of man/environment problems they recognize the need to maintain the contextual reality and integrity of any such problem as it evolves, develops, and becomes modified in the time framework of a complex society.

Proshansky, Harold M. (1985). Preface — Environmental Change/Social Change. *Environmental Change/Social Change*, pp iii - iv; Klein, S., Wener, R., Lehman, S, editors; Environmental Design Research Association; June 1985.

Environmental settings and social settings are each expressions of the nature and meaning of the other. Attempts at environmental change to affect social change must begin with understanding: knowing fully the nature, meaning, and functions of the environmental setting we are seeking to change. A planned and rationally determined change in physical setting carries with it unintended as well as intended consequences. Seeking and achieving scientific understanding of a human problem is one thing; applying that understanding and resolving that problem is another. In a democratic society only those who populate the society can make those solutions realities.

Proshansky, Harold M. (1986). Psychological Aspects of the Quality of Urban Life. *The Quality of Urban Life*; Chapter 2, pp 19 - 29; Walter de Gruyter & Co.; Berlin, New York; 1986.

There are many definitions and meanings to the term "quality of urban life". The concept of quality is multidimensional one, so the author suggests a more answerable question: What kinds of quality, for what kinds of people, in what kinds of places? Urban life is more than the organization and integration of a variety of physical settings. It not only consists of people who engage in a wide range of activities in these various physical contexts, but those individuals and activities are in turn organized and defined by larger social structures such as relationships between individuals, groups, and social institutions.

There must be a match or coherence between the person and his/her physical setting which must be examined over the life cycle of both. The quality of urban physical environments is, in the end, also rooted in many types of environmental factors such as freedom from intergroup conflicts, full employment, maintaining democratic freedoms, integrity in public life, etc.

Prudham, D.; Evans, J. Grimley (1981). Factors Associated with Falls in the Elderly: A Community Study. Age and Ageing, Vol. 10, No. 3, pp 141-146; August 1981.

Keywords: accidents; age factors; aged cerebrovascular disorders; female; wound and injuries; etiology;

Pryor, Gordon T.; Uyeno, Edward T.; Tilson, Hugh A.; Mitchell, Clifford L. (1983). Assessment of Chemicals Using a Battery of Neurobehavioral Tests: A Comparative Study. Neurobehavioral Toxicology and Teratology, Vol. 5, pp. 91-117; 1983.

Eight chemicals (acrylamide, methyl mercury, chlordecone, tetraethyl tin, triethyl lead, lead acetate, arsenic, monosodium salicylate) were compared with reference to their toxic effect on the central nervous system. The battery of tests used may have utility in the assessment of the potential neurobehavioral toxicity of various chemicals because the results compared favourably with what is known about the chemicals from other experiments with animals and from human experience.

Pryor, Paul; Reno, Stanley J. (1981). Health Hazard Evaluation Report, HETA 81-305-961, Aurora Schools. National Institute for Occupational Safety & Health, Hazard Evaluation and Technical Assistance Branch, Division of Surveillance, Hazard Evaluations, and Field Studies, Cincinnati OH.

At the request of the Aurora Schools, Aurora Colorado, an environmental evaluation was conducted at the East Middle School by NIOSH on June 19, 1981. The request concerned the potential exposure to teacher aides from methyl alcohol during the use of spirit duplicators. All of the teacher aides experienced some adverse health effects, e.g. blurred vision, headaches, burning of the nose, sluggishness, dizziness, sore throat, dermatitis, chest tightness, and depression—symptoms characteristic of toxic exposure to methyl alcohol. On the basis of the environmental and medical data, NIOSH determined that a health hazard from excessive methyl alcohol existed to the teacher aides. This exposure exists to those operating the duplicator, as well as those persons present in the room while this process is in operation. Recommendations on elimination or controlling the health hazard are included.

Raab, Karl H. (1985). *Appropriate Technology for Residential Air Quality*. Air Pollution Control Association, 1985.

In the absence of meaningful monitoring of residences, modest control approaches are appropriate. A general strategy should deal with particulate matter reduction because ample evidence indicates that particulate matter causes the major health risks in residential air. Some common household technologies can be easily modified to moderate their effect on indoor air.

A properly designed forced air heating system can produce better air quality than a hot water system if a medium efficiency extended surface fabric filter is used. If continuous low-speed fan operation is added, the system achieves temperature equalization, humidity distribution, and local odour dispersal as well. Vacuum cleaner studies indicate resuspension of precipitated particles and up to tenfold increases in suspended microbiological material. For bare floors, a manual carpet sweeper and damp mop are more appropriate. The high moisture generation from showering can be substantially reduced with a European-style hand-held hose unit. Bathroom ventilation technology has evolved to oversized, noisy exhaust fans which are infrequently used. Kitchen exhaust fans attached to an unvented hood introduces an increased health risk, possibly as a result of the recycled air stream being directed toward the user.

Raloff, J. (1987). Kids Leukemia From Parents' Exposures?. *Science News*, Vol. 132, pp 38 - 39; July 18, 1987.

A new study, first reported in the July issue of *Journal of the National Cancer Institute*, by Peters, John M. et al, says that a parent's workplace exposure to any of several classes of chemicals — particularly chlorinated solvents — or use of incense or pesticides around the home may increase children's risk of developing leukemia. The study surveyed 123 pairs of Los Angeles County families. Each pair contained one family with a leukemic child under age 10 and one family with a healthy child. A father's workplace exposure to chlorinated solvents increased his child's risk of developing leukemia, and the risk increased with frequency of exposure — to 8 times the expected rate when fathers encountered the solvents at least weekly. The authors surmise that fathers may have brought home traces of the chemical on their clothes or breath. Similar exposures to spray paint, cutting oil, methyl ethyl ketone, and dyes or pigments also showed signs of increasing a child's risk of developing leukemia. There were also notably increased risks, during nursing and pregnancy, associated with a parent's use of either incense or household and garden pesticides. A follow-up survey is being conducted.

Rand, George (1979). Caution: The Office Environment May be Hazardous to Your Health: The Need for an Ecological Approach to its Design. *AIA Journal*, pp. 38-41, 78; October 1979.

Keywords: occupational health and safety; indoor air pollution; offices; health hazards;

Rand, George (1985). Examining 'Sick' Buildings. *Architecture*, January 1985, pp. 80-83.

The author summarizes the results of a November 1984 seminar on indoor air quality sponsored by the American Institute of Architects and the California Council/AIA in response to growing concern over the "sick building syndrome". He quotes Harry Jacobs, president of the California Council: "So many different types of people use our buildings, from infants to the frail elderly in nursing homes, as well as the hardy people in between who may not be at all sensitive to polluted air. Research on health needs to be taken as seriously as the structural integrity of buildings, and it may mean adding a new type of consultant to the already long list of specialists needed to design a complex building. The architect may now have to master the languages of chemistry, biology, and medicine in order to competently orchestrate the health aspects of buildings."

Randolph, Theron G. (1976). Air Pollution. from *Human Ecology and Susceptibility to the Chemical Environment*, fifth printing; part 111, pp 35-62; Charles C. Thomas; 1976.

Environmental pollutants, both indoor and outdoor, can have debilitating mental and physical effects, especially on highly susceptible persons. The major sources of air pollution in homes and public places, such as schools, are: fuels, solvents and their combustion products; refrigerants and spray containers; insecticides; sponge rubber; plastics; mechanical devices; automobiles; and scented cleaning products. Indoor chemical air pollution of schools as a contributing cause of poor scholastic performance of susceptible children and the dizziness and confusion of susceptible teachers is rarely diagnosed correctly. The problem presented by indoor air pollution is sufficiently acute to warrant the transfer of certain highly susceptible students and teachers to more satisfactorily located, constructed, equipped, heated and ventilated classrooms or schools.

Randolph, Theron G. (1976). *Human Ecology and Susceptibility to the Chemical Environment*. Charles C. Thomas; 1976.

The author describes a wide range of clinical manifestations of maladaptation to the chemical environment—physical and mental, chronic and acute. The major chemical incitants are described and the tendency for susceptibility to spread to related materials to which cumulative exposures exist is emphasized. The diagnostic routine of comprehensive environmental control is explained. The author shows how finding and avoiding the inciting causes of illness are superior to treating the effects of illnesses.

Randolph, Theron G.; Moss, Ralph W. (1980). An Alternative Approach to Allergies: The New Field of Clinical Ecology Unravels the Environmental Causes of Mental and Physical Ills. Fitzhenry & Whiteside Limited, Toronto Canada; 1980.

Many physical and mental illnesses are caused by our increasingly contaminated environment. For those who are susceptible, everyday chemicals (gas from the kitchen range, supermarket food additives, cleaning compounds, automobile exhaust, etc.) can be a cause of mental and physical disease. Use of chemicals in schools can contribute to the overall chemical and food problem to cause poor performance by both children and teachers. Poorly designed heating and cooking systems are a major source of trouble, as are janitorial chemicals, office and art supplies. Studies of indoor air pollution in schools have been conducted with resulting anecdotal proof of cause and remedy.

Rapoport, Amos (1985). Thinking About Home Environments: A Conceptual Framework. Home Environments, Human Behavior and Environment; Vol. 8; pp. 255 - 286; Plenum Press; New York; 1985; Altman, Irwin and Werner, Carol M., editors.

Choice (preference) is important in people's interaction with all environments and is central regarding home environments; an imposed setting is unlikely to be a home environment. Constraints limiting and distorting choice may be: lack of resources and market choice; inability to cope or plan; degree of willingness and ability to move; knowledge of alternatives; external constraints like prejudice, discrimination.

Rapp, Doris (1979). Allergies and the Hyperactive Child. Sovereign Books, Simon & Schuster; 1979.

Medical reports from 1908 to the present indicate that a relationship between allergy and hyperactivity might exist. Anecdotal examples of the author's patients whose hyperactivity and learning difficulties were helped by diet modification are presented. Such things as easy distractibility, impulsiveness, emotional problems, poor coordination; perception problems, learning problems related to reading and spelling and short attention span are included under the category Minimal Brain Dysfunction. Hyperactivity, fatigue and behaviour and learning problems are often caused by medical problems such as allergies, anemia, hypoglycemia and lead poisoning.

Rapp, Doris J.; Bamberg, Dorothy (1986). The Impossible Child: A Guide for Caring Teachers and Parents. Practical Allergy Research Foundation, Buffalo NY; 1986.

Children who exhibit learning or behaviour problems may have an unsuspected or unrecognized allergy. This book is designed to help

educators and parents recognize which children have allergies, or food or chemical sensitivities interfering with their ability to learn and behave normally, and provides suggestions for helping these children. Graphic examples are presented which illustrate health and cognitive reactions by children to chemicals in the environment.

Ray, Sammy M.; Trieff, Norman M. (1980). Bioaccumulation of Anthropogenic Toxins in the Ecosystem. from *Environment & Health*, pp. 93-120; Norman M. Trieff, editor; Ann Arbor Science Publishers, Inc.

Many natural and synthetic compounds have entered the ecosystem in the past 40 years. The adverse effects of many of these were unanticipated, in that these compounds are highly resistant to degradation by natural processes and there is often a pervasive buildup on a worldwide basis. There are two types of biological endpoints in the development of water quality criteria: nonthreshold and threshold effects. In the case of carcinogens, mutagens and teratogens, thresholds cannot be established because even extremely small doses must be assumed to result in a finite increase in the incidence of response. Thus, safe levels cannot be established.

Raymer, Warren J. (1986). FDA Issues Sulfite Ban. *American Academy of Environmental Medicine Newsletter*, Vol. 21, No. 3, p 5; Winter, 1986.

The American Food and Drug Administration has banned the use of sulfites as preservatives in fresh fruits and vegetables. The agency cites a growing number of adverse reactions to the compounds, including at least 13 deaths. The ban applies to retail sales of fresh produce by food stores and restaurants. In addition, within six months, the FDA will require new labeling of processed foods that contain detectable levels of the compounds.

Rea, William J.; Bell, Iris R.; Suits, Charles W.; Smiley, Ralph E. (1978). Food and Chemical Susceptibility after Environmental Chemical Overexposure: Case Histories: Case Report. *Annals of Allergy*, Vol. 41, No. 2, pp. 101-110; August 1978.

Twelve patients with overexposure to commonly used environmental chemicals were studied. After the overexposure all developed recurrent signs and symptoms of inflammatory type diseases that were the result of ambient chemical fumes in the air and home environments. A period of time in a relatively fume-free and particle-free environment cleared the majority of symptoms and signs without the use of medication. Double-blind rechallenge with ambient dose levels of synthetic chemicals reproduced most of the symptomatology. Laboratory findings included abnormalities in complement, T-lymphocytes, eosinophils and IgG. The data suggest that some chemicals can trigger and propagate certain non-malignant inflammatory diseases, and re-emphasize the seriousness

of exposures to levels of some chemicals in our environment which were previously considered safe.

Once an individual is sensitized to a solitary chemical it is apparent that continued exposures result in a spreading phenomenon. Once this spreading occurs, reactions then proceed upon minute exposures. Spreading was demonstrated clearly in the patients in this series. After a massive exposure to one type of chemical the individual then became intolerant to ambient concentrations of many others. The mechanism of this spreading phenomenon is unclear at the present time.

Rea, William J. (1978). Environmentally Triggered Cardiac Disease. *Annals of Allergy*, Vol. 40, No. 4, pp. 243-251; April 1978.

Twelve consecutive, highly selected patients with non-arteriosclerotic cardiac arrhythmias and/or chest pain refractory to medication, having various associated symptoms relating to smooth muscle sensitization, were studied in a rigidly controlled, relatively fume- and particle-free environment. The majority of signs and symptoms were cleared in 10 patients without medication while under environmental control and all arrhythmias were reproduced with controlled, repeated, individual blind and double-blind, incitant challenges in 10 out of 12 patients. The incitants were common foods and chemicals to which the individual has been exposed frequently. Double-blind challenges with ambient doses of inhaled chemicals also reproduced the spectrum of arrhythmias (natural gas, odour of cigarette smoke, chlorine, perfume, pine-scented floor wash, ethyl alcohol, formaldehyde, phenol and pesticides). This myocardial involvement appeared to be part of a more generalized sensitization of smooth muscle. The distorted homeostatic mechanism included abnormal positive C-reactive protein, serum complements and T-B cell interactions in individual patients.

Rea, William J. (1979). The Environmental Aspects of Ear, Nose and Throat Disease: Part 1. *J.C.E.O.R.L. & Allergy*, Vol. 41, No. 7, pp. 41-56; July 1979; Medical Digest.

Ear, nose and throat involvement with environmental triggering factors appears to be the main early warning sign of environmentally triggered disease, according to this author. Sensitivity to odours seems to be the most common, many persons being intolerant of the odours of such substances as car exhaust, perfume, cigarette smoke, aerosol sprays, formaldehyde, alcohol, phenol, food, mould and dusts. Often rhinitis occurs. Recurrent sinusitis as well as severe otitis, various forms of vertigo (including Meniere's disease), and laryngeal edema are frequent presentations of the problem. Any portion of the respiratory system can be involved, resulting in such inflammatory diseases as recurrent bronchitis, bronchiectasis, and asthma. A masking phenomenon is also described which initially hides an adverse reaction.

Rea, William J. (1979). The Environmental Aspects of Ear, Nose and Throat Disease: Part II. J.C.E.O.R.L. & Allergy Digest, Vol. 41, No. 8/9, pp. 41-54; August/September 1979.

This article reviews many immune parameters that can be measured to help the clinician define changes in the body homeostatic mechanisms that occur on exposure to offending environmental agents. The researcher suggests that allergic diseases can be prevented from developing into measurable end-stage inflammatory disease, if treated early. It is being shown that patients may do well for a few months to several years with injection therapy for pollen, dust, mould, and food sensitivities, and then suddenly start deteriorating. They may lose their food sensitivity but develop a chemical sensitivity problem. It is not certain why this spreading phenomenon occurs, but it appears to be due to an overload of synthetic chemicals. Treatment of chemical sensitivity is complex, because most treatment consists of avoidance.

Rea, William J. (1979). Diagnosing Food and Chemical Susceptibility. Continuing Education, pp. 47-48, 52-53, 57-59; September 1979.

The author describes clinical experience in which a spectrum of disorders affecting smooth muscles, mucous membranes, and collagen in the respiratory, gastrointestinal, genitourinary, and vascular systems are manifested. The variety of symptoms presented are often mistaken for hypochondriasis, but actually are due to reactions to foods and chemicals found in the patient's home and work environments. Careful clinical histories should alert the physician, who can confirm suspicions by eliminating the potentially offending agents, and challenging the patient with them under controlled circumstances.

Rea, William J.; Butler, Joel R.; Laseter, John L.; DeLeon, Idefonso R. (1984). Pesticides & Brain-function Changes in a Controlled Environment. Clinical Ecology, Vol. 11, No. 3, pp. 145-150; Summer 1984.

The purpose of the present study was to determine: the extent to which chlorinated hydrocarbon pesticides in environmentally sensitive patients would be reduced by Environmental Control Unit (ECU) treatment programs; and the correlation of brain-function/psychological test results with treatment effect. Before and after therapy studies were performed on 40 ECU patients with proven levels of blood pesticides. There was a significant decrease in magnitude of blood pesticides and a significant increase in performance on brain-function/psychological tests after ECU treatment. There was also a corresponding decrease in symptoms overall. It was concluded that rigid environmental controls in treatment strongly contributed to the decrease in blood pesticide levels. Further, a more serious psychological profile was associated with these patients who showed improvement consistent with treatment.

Rea, William J.; Johnson, Alfred R.; Youdim, Said; Fenyves, Ervin J.; Samadi, N. (1986). T & B Lymphocyte Parameters Measured in Chemically Sensitive Patients and Controls. *Clinical Ecology*, Volume IV, No. 1, pp 11 - 14; 1986.

The T & B lymphocytes and T subsets of 70 proven chemically sensitive patients (symptoms including vascular dysfunction) with immune abnormalities were measured and suppression of the T8 suppressor cells was observed to the $p = .0001$ significance level versus the control subjects. The 27 asthmatics had no significant change over the controls.

Rea, William J. (1987). Environmental Electromagnetic Assessment Under Environmentally Controlled Conditions. Unpublished presentation to the 5th Annual International Symposium on Man and His Environment in Health and Disease, February 26/27, 1987, Dallas, Texas, USA;.

This researcher reported a case history of an individual known to be sensitive to low levels of chemical exposures. Blood tests for pesticide content indicated high levels, which went down over six weeks in a controlled, chemically-less-contaminated environment. The patient was improved somewhat in a new, pesticide-free home, then became ill again. Tests showed improvement in symptoms when the lights, refrigerator and stove were off; controlled experiments with exposure to 60 Hz fields for 5-10 minutes produced symptoms. Similar exposure to 50 Hz fields eliminated the symptoms. The researcher concludes that massive pesticide exposures may render an individual not only chemically sensitive, but electromagnetically sensitive as well.

Rea, William J.; Dawkins Brown, Ollie (1987). Cardiovascular Disease in Response to Chemicals and Foods. *Food Allergy and Intolerance*, Section G, Chapter 42, pp 737 - 753; Brostoff, Jonathan and Challacombe, Stephen J. editors; Balliere Tindall, Publishers; 1987.

A concept and method have now been established for the scientific definition of chemical and food triggering agents for inflammatory cardiovascular diseases including spastic vascular phenomena such as migraines and other vascular headaches, angina due to coronary spasm, Raynaud's disease, etc., many auto-immune vasculitides such as lupus, rheumatoid and other early collagen vasculitis, in addition to small and large vessel vasculitis, cardiac arrhythmias and non-traumatic phlebitis. There are now many articles in the scientific literature supporting the view that cardiovascular diseases can be caused by reactions to food and environmental irritants.

Rea, William J. (1987). Personal Interview, Dr. William Rea, Ottawa, Ontario. Personal Interview During the Canadian Society for Clinical Ecology and Environmental Medicine Conference, April 3, 1987.

The author suggests that the greatest benefit would be gained in the fight against building-related illness, by changing from the modern sealed building concept to buildings that allow open windows and greater individual control over fresh air.

Rees, David C.; Knisely, Janet S.; Balster, Robert L.; Jordan, Stephen; Breen, Timothy J. (1987). Pentobarbital-Like Discriminative Stimulus Properties of Halothane, 1,1,1-Trichloroethane, Isoamyl Nitrite, Fluorothyl and Oxazepam in Mice. *Journal of Pharmacology and Experimental Therapeutics*, Vol 241, No.2, pp. 507-515; 1987.

Volatile inhalants represent a diverse group of chemicals which pose a public health problem because of their abuse potential and neurobehavioural toxicity. They share certain properties with classic central nervous system depressants. Using drug discrimination procedures, the research showed that toluene, halothane, and 1,1,1-trichloroethane have pentobarbital-like discriminative effects.

Reible, Danny D.; Yonts, Paul; Shair, Fred H. (1985). The Effect of the Return of Exhausted Building Air on Indoor Air Quality. *Air Pollution Control Association*, 1985.

Reduced ventilation in order to improve the energy efficiency of buildings has resulted in the development of significant indoor air quality problems. The need for a balance between the competing objectives of low energy costs and good air quality are perhaps nowhere more evident than in chemical laboratories, where the potential for exposure to hazardous pollutants is very great. All too often, this need has been met through reductions in the overall ventilation of the building and a reliance upon individual laboratory fumehoods to achieve acceptable air quality. This approach is inadequate, in large measure due to the re-entry of as much as 10% of the exhausted pollutants into the building.

Reid, Lynne M. (1979). Session on Disease Conditions Predisposing Afflicted Individuals to the Toxic Effects of Pollutants: Introductory Remarks. *Environmental Health Perspectives*, Vol. 29, pp. 127-129; 1979.

Children with bronchiolitis, with cystic fibrosis, or children who have had asthma during childhood or have developed hyperlucency in the radiograph after childhood infection, are all those in whom exposure to industrial irritants or to an inclement environment may be serious. The hazard of tobacco smoking is an extremely significant factor in the susceptibility of an individual to a pollutant.

Reinhardt, Charles F. (1978). Chemical Hypersusceptibility: Original Articles. *Journal of Occupational Medicine*, Vol. 20, No. 5, pp. 319-322; May 1978.

Occupational medicine is concerned with workers who react to certain substances at low levels, below the threshold concentration associated with injury or definite discomfort. These particular workers are distinctly more susceptible than the majority and may be inadequately protected by control procedures that will suffice quite well for most employees. Their hypersusceptibility may be inherited or acquired—if the latter, it may be primarily environmental, or it may be an interaction of several factors. Nutritional hypersusceptibility is indirectly traced to the environment.

Remafedi, Gary J. (1985). Adolescent Homosexuality: Issues for Physicians. *Clinical Pediatrics*, Vol. 24, No. 9, pp. 481-485; September 1985;.

Homosexuality identity formation is a potentially tumultuous process that begins in childhood and extends through adulthood. The adolescent's experiences may ultimately contribute to a variety of physical and mental health problems. Negative reactions of family and peers toward a person's homosexuality, damaged self-esteem, the desire to be with gay-identified peers, and exposure to street life are suggested as important factors predisposing homosexuals to a higher risk of involvement with prostitution, alcohol abuse and drug abuse. There is potential for intense family discord, and physical and/or emotional abuse when homosexuality is first revealed.

The authors offer guidelines for physicians caring for homosexual youths, and stress that information about sexual decision-making, personal hygiene, prevention of sexually transmitted diseases, and substance use, can literally be life-saving for the young homosexual. They caution the physician that it is advisable to avoid anecdotal information and interjection of personal biases when counseling young people and their families on the issue of adolescent homosexuality.

Repace, James L.; Lowrey, Alfred H. (1980). Indoor Air Pollution, Tobacco Smoke, and Public Health. *Science*, Vol. 208, pp. 464-472; May 2 1980; American Association for the Advancement of Science.

Indoor air pollution from tobacco smoke is pandemic. In the presence of tobacco smoke, many normal nonsmokers experience eye and throat irritation, headache, rhinitis and coughing; allergic persons report wheezing, sneezing and nausea as well. Particularly acute symptoms may be found in infants, children, persons with cardiovascular or respiratory disease and wearers of contact lenses. The authors undertook a systematic study of the levels of respirable suspended particulates in several common indoor environments. They showed that under the practical range of ventilation conditions and building occupation densities, the respirable suspended particulates levels generated by smokers overwhelm the effects of ventilation and inflict significant air pollution burdens. Smoking indoors may be incompatible with the goal of maintaining a high level of indoor air quality.

Repace, James L.; Ott, Wayne R.; Wallace, Lance A. (1980). Total Human Exposure to Air Pollution. Air Pollution Control Association.

A number of studies have shown that fixed monitoring stations do not accurately reflect the exposure of the population to outdoor pollutants in the indoor environment. Furthermore, various studies have indicated that indoor exposures to air pollutants can be very important contributors to air pollution burdens of the population. Factors affecting an individual's "exposure" are explored by modeling and measurement, and research needs and implications for regulatory action are discussed.

Repace, James L. (1981). The Problem of Passive Smoking. Bulletin of the New York Academy of Medicine, Vol. 57, No. 10, pp. 936- 946; December 1981.

Substantial new evidence concerning the adverse health effects of passive smoking have recently emerged. This new evidence indicates that well-known health effects of smoking may be suffered by nonsmokers who breathe tobacco-smoke-contaminated air. Concentrations of tobacco smoke indoors are directly proportional to the smoker density and inversely proportional to the effective ventilation rate. Attempts to control smoking by ventilation are futile, requiring ventilation rates far in excess of what is economical, and are contrary to the current trend toward energy conservation in buildings. However, alternative measures which reduce the source have been proved effective.

Repace, J.L.; Lowrey, A.H. (1982). Tobacco Smoke, Ventilation, and Indoor Air Quality. ASHRAE Transactions 88, part 1, pp. 895-914; American Society of Heating, Refrigerating and Air Conditioning Engineers, Inc., 1791 Tullie Circle NE, Atlanta GA 30329; 1982.

Cigarette smoking has been found to be causally related to cancer, cardiovascular disease, and pulmonary disease. Based on a 1,078,894 person prospective epidemiological study covering 25 states, statistically significant dose-response relationships between exposure to tobacco smoke and loss of life expectancy have been calculated. From examining the data according to mortality by degree of inhalation of tobacco smoke, it can be shown that a limited amount of exposure to tobacco smoke carries a greatly increased risk of premature mortality, a fact that has great relevance for the problem of passive smoking. Breathing of ambient tobacco smoke by nonsmokers carries a significantly increased risk of morbidity and mortality.

Repace, James L. (1983). Effect of Ventilation on Passive Smoking Risk in a Model Workplace. US Environmental Protection Agency, Washington DC 20460.

An estimate of the variation of nonsmokers' lung cancer risk from so-called passive or involuntary smoking is given as a function of ven-

tilation rate in a typical office, at an occupancy of 7 persons/100 m² (per 1000 ft²) as specified under ASHRAE Standard 62-1981, Ventilation For Acceptable Indoor Air Quality. Assuming one third of the office workers are smokers who smoke at the average rate of two cigarettes per hour, the nonsmoking office workers exposed to environmental tobacco smoke for a 40-year period under the ventilation rate recommended by the standard, would be subject to an estimated lifetime involuntary risk of about 250 per 100,000. This risk is 250 times the maximum lifetime value considered acceptable under commonly used environmental criteria for carcinogenic contaminants in air, water or food. Practical increases in makeup air or assists by air cleaning, although helpful, cannot reduce the risk to an acceptable level.

Repace, James L. (1983). Risks of Passive Smoking. University of Maryland, Center for Philosophy and Public Policy, College Park, Maryland 20742 USA; August 1983.

The risks to nonsmokers from breathing indoor air pollution from tobacco smoke are discussed from medical, legal, social, historical and philosophical points of view. Problems encountered by nonsmokers who are hypersensitive to tobacco smoke are contrasted with chronic risks to normal nonsmokers. Ambient tobacco smoke is compared to hazardous outdoor air pollutants and the products of other human activities which are regulated by society, and it is argued that failure to regulate indoor smoking will invite increasing confrontation and social dissension between nonsmokers, smokers and employers, since the nonsmoker's claim on clean indoor air is morally superior to the smoker's right to pollute.

Repace, James L.; Lowrey, Alfred H. (1983). Modeling Exposure of Nonsmokers to Ambient Tobacco Smoke. Air Pollution Control Association; August 1983.

The authors have modeled the exposure of the nonsmoking US population to the particulate phase of tobacco smoke from indoor air pollution in buildings. It was estimated that the average US nonsmoker of working age receives an exposure of 1.4 mg per day, a weighted average taken over exposures encountered both at home and at work. Estimates show that the ratio of workplace dose to the exposure received at home is nearly 4:1, indicating that, on the average, the workplace is a more important source of exposure than the home environment. Such estimates of exposure should prove useful both for assessing exposure in epidemiological studies of passive smoking, and for carcinogenic risk assessment, as well as for estimating total population exposure in epidemiological studies of the health effects of particulate air pollution.

Repace, James L.; Lowrey, Alfred H. (1984). A Proposed Indoor Air Quality Standard for Ambient Tobacco Smoke: from Indoor Air, Berglund, Lindvall, Sundell, editors. Swedish Council for Building Research, Stockholm Sweden; pp 235-239; 1984.

The authors propose an indoor air quality standard for ambient tobacco smoke particles for a typical US workplace, based upon limiting the involuntary carcinogenic risk to a nonsmoker from passive smoking to 3×10^{-7} annually, corresponding to a 1 per 100,000 risk for a 40-year working life. To achieve this level of risk, the maximum permissible annual average of the daily workplace concentrations must be 0.75 micrograms per cubic meter. At typical smoking occupancies for an office, achieving the standard would require impractical amounts of ventilation or prohibitive costs for air cleaning equipment. It appears that complete physical separation of smokers and nonsmokers or prohibition of workplace smoking are the only practical control measures.

Repace, James (1984). Tobacco Smoke: The Double Standard. QQ—Report from the Center for Philosophy & Public Policy, University of Maryland, College Park, Maryland 20742; Vol. 4, No. 1, pp 6-8; Winter 1984.

A research scientist blasts the double standard that treats air pollution more leniently when emitted from a lit cigarette than from a factory smokestack.

Repetti, Rena L. (1987). Individual and Common Components of the Social Environment At Work and Psychological Well-Being. *Journal of Personality and Social Psychology*, Vol. 52, No. 4, pp. 710 - 720; 1987.

The social environment at work is related to the mental health of employees. Two components of a social environment were measured in a study: a common social environment, the social climate shared by employees in the same work setting, and an individual social environment, the social space surrounding one individual in the setting. The study averaged co-workers' ratings and individuals' own ratings of the social environment to individuals' self-reported psychological well-being. A group of 37 bank branches represented work environment and nonmanagerial personnel in the branches served as participants. Aggregate co-worker ratings of the common social environment were significantly correlated with individual depression and anxiety. However, an individual's perceptions appeared to mediate the social environment's impact. As hypothesized, well-being was more closely tied to the proximal individual social environment than to the common social environment.

Resin, Torjman, Sherri (1978). Mental Health in the Workplace. Canadian Mental Health Association; November 1978.

The material was selected so as to constitute a core body of knowledge of both theory and practice on the subject of mental health in the workplace.

Rieber, Elaine R. (1984). The Question of Safety in Secondary School Science Laboratories. Canadian Council on Children and Youth, Ottawa Canada; May 1984.

This paper was prepared in response to a 1983 letter from a secondary school teacher in Quebec concerning the issue of mercury vapour in science laboratories and its potential harmful effect on the health of children. It was concluded that there are no provincial regulations with respect to safety standards for secondary school science labs, and no systematic monitoring of safety conditions exists at any level.

Riesenberg, Donald E; Arehart-Treichel, Joan (1986). "Sick Building" Syndrome Plagues Workers, Dwellers. Journal of the American Medical Association; Vol. 255, No. 22, p. 3063; June, 1986.

"Sick building" syndrome is increasingly becoming a valid diagnosis used to explain illness in occupants of energy-efficient, tightly-constructed office and residential buildings with artificial ventilation. Whether it be hypersensitivity pneumonitis, irritation from fiberglass or other chemicals, or lassitude, scientists working in this area predict that physicians are likely to see increasing numbers of patients who work and live in sick buildings. Increased awareness of climatic pathogens will be required because physicians are, on the whole, poorly informed about the relationship between workplace exposures and ill health.

Riley, E.C.; Murphy, G.; Riley, R.L. (1978). Airborne Spread of Measles in a Suburban Elementary School. American Journal of Epidemiology, Vol. 107, No. 5, pp. 421-432; 1978; Johns Hopkins University School of Hygiene and Public Health.

A measles epidemic in a modern suburban elementary school in upstate New York in spring, 1974 is analyzed in terms of a model which provides a basis for apportioning the chance of infection from classmates sharing the same home room, from airborne organisms recirculated by the ventilating system, and from exposure in school buses. The epidemic was notable because of its explosive nature and its occurrence in a school where 97% of the children had been vaccinated. Many had been vaccinated at less than one year of age. The index case was a girl in second grade who produced 28 secondary cases in 14 different classrooms. Organisms recirculated by the ventilating system were strongly implicated. After two subsequent generations, 60 children had been infected, and the epidemic subsided. From estimates of major physical and biologic factors, it was possible to calculate that the index case produced approx. 93 units of airborne infection per minute. The exceptional infectiousness of the index case, inadequate immunization of many of the children, and the high percentage of air recirculated throughout the school, are believed to account for the extent and sharpness of the outbreak.

Riley, R.L. (1982). Indoor Airborne Infection. *Environment International*, Vol 8, pp 317-320, 1982.

Airborne infection from person to person is an indoor phenomenon. The infectious organisms are atomized by coughing, sneezing, singing, and even talking. The smallest of droplets evaporate to droplet nuclei and disperse rapidly and randomly throughout the air of enclosed spaces. Droplet nuclei have negligible settling velocity and travel wherever the air goes. Measles and other childhood contagions, the common respiratory virus infections, pulmonary tuberculosis, and Legionnaires Disease are typically airborne indoors. In analyzing a measles outbreak, the probability that a susceptible person would breathe a randomly distributed quantum of airborne infection during one generation of an outbreak was expressed mathematically. Estimates of the rate of production of infectious droplet nuclei ranged between 93 and 8 per minute, and the concentration in the air produced by the index case was about 1 quantum per 5 m³ of air. Control of indoor airborne infection can, according to the author, be approached through immunization, therapeutic medication, and air disinfection with ultraviolet radiation.

Rimland, Bernard; and Larson, Gerald E. (1981). Nutritional Approaches to the Reduction of Criminality, Delinquency and Violence. *The Journal of Applied Nutrition*, Volume 33, Number 2, 1981, pp. 116-133.

The traditional approaches to the treatment of offenders, through counseling and psychosocial rehabilitation, have in repeated studies been found to be useless or counterproductive. These methods assume antisocial behaviour is primarily sociogenic in origin. The author presents another point of view, that much crime, particularly violent crime, is caused by biological malfunctioning of the brain. The strong association between learning disabilities and antisocial behaviour suggests that both learning problems and problems with the law may stem from a common cause - brain malfunction. Maternal smoking, poor nutrition, toxic metal and other chemical exposure, and food allergies are among the adverse factors considered as causes of brain malfunction in youth.

Rittfeldt, Lars; Sandberg, Maria; and Ahlberg, Mats S. (1984). Indoor Air Pollutants Due to Vinyl Floor Tiles. *Proceedings, 3rd International Conference on Indoor Air Quality and Climate*, held in Stockholm, Sweden, August 20-24, 1984. Volume 3: Sensory and Hyperreactivity Reactions to Sick Buildings, pp. 297-301.

Most pollutants emitted from vinyl floor tiles are found to originate from the surface film and the patterned film used in the manufacturing of the tiles. These films are plasticized with butyl benzyl phthalate, which makes the tiles easy to clean. Butyl benzyl phthalate was shown to be contaminated with benzyl chloride and benzal chloride, compounds which

are irritating to the eyes and respiratory mucosa and which also are carcinogenic. The concentrations in the films and in the vinyl floor tiles of these substances were measured and their emission rates from a newly produced tile were determined. The author suggests that the emission of benzyl chloride and benzal chloride from vinyl floor tiles plasticized with butyl benzyl phthalate may be a factor contributing to the irritations in so-called "temporary sick buildings".

Robbins, Albert F. (1987). Neurotoxic Effects of Pesticides Called Critical Risk to Health. American Academy of Environmental Medicine Newsletter, Volume 22, No. 1, p 6; Spring, 1987.

The Center for Science in the Public Interest recently asserted that most health symptoms caused by pesticides are effects on the nervous system. In a petition to the US Environmental Protection Agency, CSPI was joined by the American Public Health Association, the American Psychological Association and 8 other groups to require that pesticides be evaluated for their neuro-toxic and neuro-behavioural effects.

Robertson, Leon S. (1986). Injury: from Handbook of Prevention; Edelstein, Barry A. and Michelson, Larry, editors. Plenum Press, New York, 1986; pp. 343 - 360.

Injury is among the most important public health problems in terms of the numbers of people affected and severity in cases of permanent disability and death. About one third of the population of the US is injured severely enough in a year to report an injury in the National Health Survey. It is the fourth leading cause of deaths. Traditionally, injury prevention has been considered a problem of accident prevention, a supposition which contains several unfortunate implications that have tended to retard consideration of the full range of options available to prevent injuries or reduce their severity. The continued high injury rates associated with road vehicles, guns, farm machinery, sports equipment, cigarettes, and other consumer products are not for lack of knowledge of what can be done to sharply reduce them. For major gains to be made in injury control, the government must act to set standards for the distribution and use of potentially injurious products.

Robinson, Alan M.; Sheppard, Howard; and committee (1983). Child Abuse: Second Report on Family Violence:. Government of Ontario; December, 1983.

The primary responsibility of children's aids societies must be the protection of children. Interprofessional cooperation is essential at all stages of a case of child abuse. The central child abuse register must be made more effective. Abusers should be charged under the criminal code and child welfare act. Public and professional awareness of child abuse must be increased. The movement towards Native autonomy in the provision of child welfare services must be accelerated. Research into the problem

of child abuse must be encouraged and funded. There is a particular need for long-term studies of abuse victims. Research into the environmental factors, including diet, that could affect the incidence of child abuse must also be conducted.

Rogers, Sherry A. (1986). Indoor Air Quality and Environmentally Induced Illness: A Technique to Evoke Chemically Induced Symptoms in Patients. IAQ'86, p 71 - 77; American Society of Heating, Refrigerating and AirConditioning Engineers, Inc.; 1986.

Formaldehyde is but one of many chemicals capable of causing environmentally induced illness. The spectrum of symptoms it may induce includes attacks of headache, flushing, laryngitis, dizziness, extreme weakness or exhaustion, inability to think clearly, arrhythmia or muscle spasms. The nonspecificity of such symptoms can baffle physicians from many specialties. Presented in this report is a simple office method for demonstrating that formaldehyde is among the etiologic agents triggering such symptoms. The very symptoms that patients complain of can be provoked within minutes, and subsequently abolished, with an intradermal injection of the appropriate strength of formaldehyde. This aids in convincing the patient that his symptoms can result from formaldehyde, one of the many triggers of environmentally induced illness. The patient can then begin to relate symptoms to environment exposures and initiate measures to avoid such triggers and bring his disease under control.

Root, David E.; Anderson, Joan (1986). Reducing Toxic Body Burdens: Advancing in Innovative Technique. Occupational Health and Safety News Digest, Vol. 2, No. 4; April, 1986.

Assessing and reducing chemical toxicity is of increasing interest to industry. Some corporations and medical advisors have developed policies emphasizing early detection and preventive measures. Case histories are presented where a regimen was employed which reduced body stores of some highly bioaccumulative substances, including organohalides such as polychlorinated biphenyls. It included aerobic exercise, polyunsaturated oil supplement, sauna, nutritional supplements, and water and salts taken as needed. The effect of this regimen was to enhance the mobilization of toxic substances from tissues.

Roy, Alan W. (1987). Letter from Union of Ontario Indians About Pollution of Ojibway Land. Personal Correspondence from Alan W. Roy, Environmental Director, Union of Ontario Indians, Toronto, Ontario.

The author cites the attitudes taken by the federal government and certain corporations, towards the Ojibway people (Anishinabek) and the pollution of their land, as contributing to an unhealthy environment for Ontario native people. In the 1950s, a sulphuric acid factory was built adjacent to an existing native community, within the Serpent River Indian Reserve, half-way between Sault Ste. Marie and Sudbury, on the

North Shore of Georgian Bay. In 1963, the factory was abandoned for economic reasons, leaving a toxic waste dump containing 120,000 tons of sulphur, calcine and pyrite which eventually contaminated soils over 100 acres of the reserve. (See also 'Cutler Acid Plant Study', Stokes, 1981.)

The Union of Ontario Indians suggests that the original cooperation by the Department of Indian Affairs to offer Reserve Land to industry, for such an environmentally deleterious operation, was indicative of an unhealthy attitude, at that time, towards Ontario's aboriginal people. They charge that 30 years of delay in working out financing for an appropriate rehabilitation program (estimated at \$2 million) is proof that this attitude has persisted. The author speculates that if this situation existed in Southern Ontario, with a constituency of the dominant society middle class Ontarians, the site would have been rehabilitated much more quickly.

Sachs, H.M.; Hernandez, T.L.; Ring, J.W. (1982). Regional Geology and Radon Variability in Buildings: from Indoor Air Pollution; Spengler, John—editor. Environ. Int., Vol. 8, No. 1/6, pp. 97-103; 1982; Pergamon Press Ltd., Oxford, Toronto.

Radon concentrations in dwellings vary by more than two orders of magnitude. Predicting where and when concentrations are likely to be high requires studying the variability of the contributors to radon in buildings. Among common sources, geological factors (water supply and substrate) are the most variable, whereas building materials are much less variable. Ventilation variation among houses is generally responsible for radon variations comparable to those introduced by building materials, but it is more significant at lower ventilation rates. In some regions with relatively high proportions of houses with elevated radon concentrations, mappable geological factors are associated with most cases of high radon concentrations.

Saegert, Susan (date?). Crowding: Cognitive Overload and Behavioral Constraint. Environmental Design Research, Vol. 2; pp 254 - 260.

Hypotheses concerning cognitive and behavioral effects of crowding are developed by looking separately at its two physical components: number of people in a space and amount of space per person. The first would tend to increase cognitive complexity and uncertainty in the situation and the difficulty of organizing behavior. The second would make others in the situation more salient as stimuli and require greater coordination of behavior. Together these factors would tend to produce cognitive overload and behavioral constraint. Supporting results from an exploratory study conducted in a mid-Manhattan department store are reported.

Saegert, Susan (undated). High Density Environments: Their personal and Social Consequences. Human Responses to Crowding, Lawrence Erlbaum Associates, Norwood, NJ; Baum, A.; Epstein, Y., editors.

Many studies have been undertaken which make it clear that high-density experiences and living conditions can provoke stress and anti-social behaviours and attitudes. But other studies present contradictory findings. There is no such thing as density per se. The term refers to people living in and moving through certain amounts of space that varies in configuration, definition and types of interpersonal interaction as well as perception on the part of the people involved. Since high density systems seem to require more from people, economic, health and social disadvantages appear to take more of a toll in these environments, although the many opportunities provided may at times offset such problems. Some implications for physical design and planning of high density environments are given which include conducting investigations of standing patterns of social relations and activity systems prior to planning; constructing facilities and policies to either support or develop affinity groups and patterns of behaviour that have existed in other situations; definition of manageable-size groups and provision of flexible shields from interaction for groups and individuals; and provision for access to lower density environments for individuals during the course of their normal life activities. An excellent bibliography is included.

Saegert, Susan; Winkel, Gary (date?). The Home; A Critical Problem for Changing Sex-Roles. Unpublished.

The home is a significant physical and symbolic environment for both men and women. The activity patterns of women, especially mothers of young children, are strongly influenced by the home's location, as well as the women's sense of options and feelings about their identity. Urban men appear to be less emotionally invested in the home than their wives; suburban men are happier in suburban homes. This raises questions for women who want their homes to be meaningful centres of the family and who want to expand their roles outside the home. If women are to expand their roles outside the home, a wide range of home-related activities and values must be changed in some way. Higher density, mixed-use residential environments give women more options. The time schedule and reward structure of most work is one of the strongest pressures against the combination of physical and psychological investment in the home and outside. The geographic segregation of residential environments from public life reinforces the cultural choice of work or home, especially for women, who do not have the luxury of a wife.

Saegert, Susan (1975). Stress-Inducing and Reducing Qualities of Environments. Environmental Psychology, 2nd edition; Proshansky, H.M. Ittelson, W.H., and Rivlin, L.G. editors; Holt, Rinehart & Winston; 1975.

Environments can both induce and reduce stress. Stress-inducing environments include those which are physically threatening, noisy and polluted. These environmental stressors have not been extensively investigated, and never in the context of their full complexity. Stimulus or information overload is also a stress-provoking quality of environments, as is decisional overload which results from a person's relationship to the environment. This combination of physical environment and psychosocial orientation contributes to the dimension of experience in cities which is referred to as the pace of life. Also to be considered are the suitability of an environment for the particular people and the activities that engage them (the frustration experienced by commuters in a hurry to get home but tied up in traffic or by children wanting to play in high-rise buildings which prohibit such activity) and the psychologically and socially meaningful messages associated with particular physical environments (people experiencing increased status become healthier and those whose educational or class background exceed their current status are less healthy than others in the same circumstances). Implications for stress-reducing environments are discussed. An extensive bibliography is included.

Saegert, Susan (1980). Masculine Cities and Feminine Suburbs: Polarized Ideas, Contradictory Realities. *Signs: Journal of Women in Culture and Society*, Vol. 5, No. 3 suppl. pp s96 - s111; 1980.

A wide range of symbolic associations attach to women and suburbs versus men and cities. Urban life and men tend to be thought of as more aggressive, assertive, definers of important world events, intellectual, powerful, active, and sometimes dangerous. Women and suburbs share domesticity, repose, closeness to nature, lack of seriousness, mindlessness, and safety. The growth of suburban populations during a time of increased female labour force participation can be added to a list of contradictory trends in this era that place rather greater demands on women. Many now have two roles — work outside the home and responsibility for domestic life. The public services that might help women manage are not given priority by suburbs which operate under the myth of the housewife in an effort to preserve class and racial exclusiveness. Higher density, mixed-use residential environments give women more options. Collective visions of an integrated life of domestic work, productive work, and leisure must inform public policy and physical planning in order to bring forth an organization of time and space in which that integration is fostered.

Saegert, Susan (1981). Environment and Children's Mental Health: Residential Density and Low Income Children. *Handbook of Psychology and Health*, Volume 2; Erlbaum Associates; 1981; Baum, A.; Singer, J., editors.

A review of research on children and density is presented, along with a description of recent research conducted by the author which examines the influence of residential crowding and high rise living on children's health and psychological development. Higher levels of occupancy and interaction in high density apartments result in more frequent conflicts

between children. In large buildings, children have more difficulty in accurately conceptualizing the larger buildings and know proportionally fewer people, which leads to perceptions of less friendliness among tenants, less likelihood of aid being offered, and less guilt among children about committing antisocial acts.

Salvaggio, John E. (1986). Human Symptomatology and Epidemiology of Fungi in Air. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

This report covers occupational and environmental problems due to air exposure to fungal contamination. It includes a discussion of relevant clinical data on patients, specific measurements made with air sampling devices, the importance of particle size, and lung biopsy data where appropriate. An exhaustive bibliography is included.

Sandia National Labs (1982). Indoor Air Quality Handbook: For Designers, Builders, and Users of Energy Efficient Residences. US Govt.; NTIS DE83002315; Sept. 1982.

Keywords: indoor air pollution; residential buildings; air pollution control; energy efficiency; pollution sources; radon; ventilation; formaldehyde; carbon monoxide; nitrogen oxides; organic compounds; smokes; low pollution design

Sarason, Irwin G.; Spielberger, Charles D. (1983). Stress and Anxiety: Volume 6. Hemisphere Publishing Corporation; 1983.

A concern with life crises and with the stress and anxiety generated by such crises projects the concept of environmental security into a prominent place in the person-environment conceptual framework. This concept expresses the affective correlates of individuals' degrees of competence, control, and knowledge in relation to their physical worlds. To have such control, understanding, and competence are matters of critical importance to a person's self-identity and related feelings of self-esteem. Physical settings, however, because of their frequently growing complexity and accelerated change, increasingly make demands on people to the extent that the pursuit of environmental security in an urban context becomes an unending process for urban dwellers.

Saric, Marko; Fugas, Mirka; Hrustic, Omer (1981). Effects of Urban Air Pollution on School-Age Children. Archives of Environmental Health, Vol. 36, No. 3, pp. 101-108; 1981.

From November 1977 to March 1978, forced expiratory volumes (FEVs) of 78 second graders living in a high sulfur dioxide and smoke pollution area were compared with FEVs of 70 second graders living in a clean air area.

The incidence of acute respiratory diseases in these children and their families was also comparatively studied during the same period. Indoor and outdoor measurements of sulfur dioxide and smoke, as well as additional measurements of suspended particulate matter (SPM) and sulfate were conducted. The results of the study suggest that at the actual average annual exposure to sulfur dioxide, smoke, and SPM of 70-80 ug/m³, 60-80 ug/m³, and 130-200 ug/m³, respectively, with frequent exposures to three to five times higher daily sulfur dioxide and smoke concentrations and two times higher daily concentrations of SPM during the heating season, certain effects on the ventilatory functions and occurrence of acute respiratory diseases can be expected.

Sass, Robert (1986). Workplace Health and Safety: Report from Canada: Original Articles on Occupational Health Policy. International Journal of Health Services, Volume 16, Number 4, 1986; pp 565 - 582; Baywood Publishing Co., Inc.

This article represents a critical analysis of the major policy responses to workplace health and safety in Canada. It examines the deficiencies inherent in the legislative development of Joint Health and Safety Committees in most Canadian jurisdictions, the limitations regarding standard-setting of worker exposure to contaminants, and disincentive for employers to positively improve the workplace because of Workers Compensation legislation. Collective bargaining agreements in Canada have had only limited positive effects, while the ultimate legal sanction of criminal prosecution by the regulatory agencies has weakened enforcement and compliance of existing regulation. There has never been a successful criminal prosecution of an employer in Canada, even for multiple deaths.

Four reasons are suggested: the concealment of the dimension of the incidence of industrial disease based on Workers Compensation Board statistics; the application of an incorrect theory of causation of both industrial disease and injury by both managers and government administrators of occupational health and safety programs; the resistance of both senior and middle managers against increased worker participation in both work organization and job design questions; and the general moral underdevelopment rather than ignorance of managers in favouring economic considerations or values at the expense of worker health and safety. The author proposes the need for greater workplace democratization of production and industry as a necessary and sufficient reform of workplace health and safety.

Savitz, David A.; Calle, Eugenia E. (1987). Leukemia and Occupational Exposure to Electromagnetic Fields: Review of Epidemiologic Surveys. Journal of Occupational Medicine, Vol. 29, No. 1, pp 47 - 51; January, 1987.

Several recent surveys have presented data suggesting an increased risk of leukemia among men with occupational exposure to electromagnetic fields. Eleven pertinent data sets were compiled in order to assess the

consistency of this pattern and to identify those occupations most deserving closer examination. Results for total leukemia show a modest excess risk for men in exposed occupations, with an enhanced risk elevation for acute leukemia and especially acute myelogenous leukemia. These studies are inherently limited regarding the effect of electromagnetic fields due to the absence of exposure characterization. Nonetheless, telegraph, radio, and radar operators, power and telephone linemen, and electrical and electronic engineers showed the most consistent results and warrant further study to ascertain potential occupational health hazards.

Savitz, David A.; Calle, Eugenia E. (1987). Leukemia and Occupational Exposure to Electromagnetic Fields: Review of Epidemiological Surveys. *Journal of Occupational Medicine*, Volume 29. No. 1, January 1987; pp. 47-51; American Occupational Medicine Association;

This review compiled eleven sets of data from previous study to assess leukemia risk among men with occupational exposure to electromagnetic fields. Results for total leukemia show a modest excess risk for men in exposed occupations, with an enhanced risk elevation for acute leukemia and especially acute myelogenous leukemia. The studies lacked characterization of the electromagnetic exposure. Nevertheless, telegraph, radio and radar operators, power and telephone linemen, and electrical and electronic engineers showed the most consistent results. The authors caution that the available data are inadequate to conclude that electromagnetic field exposures are the reason for that elevation, and they suggest further study to ascertain potential occupational health hazards.

Scadding, Glenis K.; Brostoff, J. (1986). Low Dose Sublingual Therapy in Patients With Allergic Rhinitis Due To House Dust Mite. *Clinical Allergy*, Vol. 16, pp 483-491; 1986.

In a double-blind placebo-controlled trial, low dose sublingual therapy with house dust mite was effective in relieving symptoms in 72% of the group of patients with perennial rhinitis due to house dust mite. Following active treatment, there was a significant increase in morning peak nasal inspiratory flow rate in those who improved and resistance to nasal provocation with house dust mite also increased, in some cases up to 1000-fold. Oral therapy is safe and avoids the side effects of desensitizing injections which can be serious. The potential for oral desensitization is great and further studies on this form of treatment are needed.

Schaumburg, Herbert; Spencer, Peter S. (1978). Environmental Hydrocarbons Produce Degeneration in Cat Hypothalamus and Optic Tract. *Science*, Vol. 199, pp. 199-200; January 1978.

Keywords: environmental hydrocarbons; animal studies; health effects;

Schauss, Alexander (1980). Diet, Crime and Delinquency. Parker House, Berkeley CA 94704; 1980.

Junk food diets, sugar starvation, vitamin deficiencies, lead pollution; and food allergies can convert a normal brain into a criminal mind. Children absorb and retain a greater percentage of ingested lead than adults. Lead levels formerly considered safe have recently been shown to increase abnormal behaviour, cause learning difficulties and reduce intelligence.

Schauss, Alexander (1980). Lead, Behaviour and Criminality: from Diet, Crime and Delinquency; chapter 111; pp. 32-49. Parker House, Berkeley CA; 1980.

Lead levels formerly considered safe have recently been shown to increase abnormal behaviour, cause learning difficulties and reduce intelligence. Children can absorb enough lead to impair their performance on tests of reasoning, coordination, intelligence and reading. Some symptoms of lead toxicity are: hyperactivity, perceptual disorders, mental retardation, fatigue, irritability, temper tantrums, learning disabilities, speech disturbances, perceptual motor dysfunctions, and emotional or behavioural problems. There are two new techniques helpful in determining heavy metal concentrations, replacing blood tests. These are hematofluorometry and hair trace mineral analysis.

Schiefer, H.B. (1985). Health Effects from Mycotoxins (volatile or absorbed to particulates): A Review of the Relevant Data in Animal Experiments. University of Saskatchewan; December 1985.

Assuming that it is correct that potentially toxigenic fungi can be isolated from dust in indoor polluted homes, and given the clinical descriptions of health problems of inhabitants of such dwellings, it is possible to hypothesize that the health problems encountered are due to trichothecene mycotoxins. The subtle effects on the immune system will, however, be masked by the more obvious clinical and pathological findings of an opportunistic infection, thus escaping the attention of clinicians. There is a chance that an aero-allergenic reaction is taking place due to the proteins contained in the dust or associated with the fungal spores or other inorganic material or chemicals. The question should be pursued with care and priority.

Schlesinger, Richard B. (1985). Effects of Inhaled Acids on Respiratory Tract Defense Mechanisms. Environmental Health Perspectives, Vol. 63, pp 25 - 38; 1985.

The respiratory tract is endowed with an interlocking array of non-specific and specific defense mechanisms which protect it from the effects of inhaled microbes and toxicants, and reduce the risk of

absorption of materials into the bloodstream. Ambient acids may compromise these defenses, perhaps providing a link between exposure and development of chronic and acute pulmonary disease. This paper reviews the effects of inhaled acids upon the nonspecific clearance system of the lungs. An extensive bibliographic list is included.

Schnare, David W.; Ben, Max; Shields, Megan G. (1984). Body Burden Reductions of PCBs, PBBs and Chlorinated Pesticides in Human Subjects. *Ambio, A Journal of the Human Environment*, Vol. 13, No. 5-6; pp 378 - 380; 1984.

With human exposure to environmental contaminants inevitable despite the best application of environmental laws and protection technologies, interest has grown in the potential to reduce the levels of contamination carried in the human host. This study demonstrates the promise of a comprehensive treatment for reduction of body burdens of polychlorinated and polybrominated biphenyls and chlorinated pesticides. Adipose tissue concentrations were determined for seven individuals accidentally exposed to PBB. These patients underwent detoxification treatment to eliminate fat-stored foreign compounds. Of the 16 organohalides examined, 13 were present in lower concentrations at post-treatment sampling. 7 of the 13 reductions were statistically significant. To determine whether reductions reflected movement to other body compartments or actual burden reduction, a post-treatment follow-up sample was taken four months later. Analysis showed a reduction in all 16 chemicals. Future research stemming from this study should include further investigation of mobilization and excretion of xenobiotics in humans.

Schrecker, T.F. (1984). Political Economy of Environment Hazards: Protection of Life Series, Study Paper. Law Reform Commission of Canada; 1984.

'Political' should be taken to not only include the actions of elected officials and bureaucrats, but also the process of selecting the issues and defining the alternative courses of action which are considered by these and other actors. The law itself will usually reflect similar mobilizations of bias. This paper deals both with the process by which environmental hazard law and policy are made and with the conceptual frameworks which are used to define objectives and strategies for controlling environmental hazards. Also examined is the role of the large profit-oriented corporation as a policy-making institution. The system of property rights which allows business relatively unfettered control over investment flows, on the basis of maximizing profitability, seriously restricts the ability of governments to intervene in many kinds of decisions which are now the prerogative of the corporation.

Schreiter, Anne (1984). Human Ecology in the Schools. Human Ecology Foundation of Canada, Kitchener Branch.

This article outlines causes of and solutions for chemically sensitive children in the school system. Suggestions are made for reducing the chemical load of the children and teachers and solutions are presented that have been used in actual situations of chemical sensitivity.

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

Schreiter, Anne (1984). A Follow-up to Human Ecology in the Schools. The Human Ecology Foundation of Canada, Kitchener Branch.

Suggestions and solutions for chemically sensitive children in the school system. Suggestions are made for reducing the chemical load of the children and teachers, including that smoking be banned on school property.

Schroeder, Henry A. (1973). Cadmium, High Blood Pressure and Water. from Trace Elements and Man; chapter VIII, pp. 97-114; Devin-Adair Co.; 1973.

There is evidence that cadmium from refined foods, water pipes and contaminated air and water accumulates in human kidneys and causes hypertension or high blood pressure.

Scott, P.M. (1986). Mycotoxins Produced by Certain Fungi Associated with Thermal Insulation. Significance of Fungi in Indoor Air, Part II, Working Papers; Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Mycotoxins associated with thermal insulation are described in detail. An exhaustive bibliography is included.

Scott, Peter M. (1983). Other Mycotoxins. Proc. Int. Symp. Mycotoxins, pp. 87 - 110; 1983.

With the exclusion of the aflatoxins, fusarium toxins, and penicillium toxins, the remaining known mycotoxins are produced by fungi of the genera *Aspergillus*, *Alternaria*, *Trichothecium*, *Stachybotrys*, *Myrothecium*, *Claviceps*, *Deplodia*, *Chaetomium*, and over a dozen others. Natural occurrence has been demonstrated for some in foods or agricultural commodities as a result of fungal infection. The epidemics of human ergotism in Europe in medieval times, more recent incidences in India, and many diseases in animals, are attributable to these toxins. An excellent bibliography is included.

Seguin, Marilynne (1987). Dying With Dignity: A Canadian Society Concerned with the Quality of Dying. Dying With Dignity; 1987.

Dying With Dignity is a charitable, volunteer organization that believes people should have the right to control the ending of their own lives. This includes the right to voluntary euthanasia, suicide, and to refuse treatment. It is based on the concern of many people that modern technology has made it possible to keep people alive past the point at which life has ceased to be worth living.

Seifert, Bernd; Drews, Marianne; Aurand, Karl (1984). Indoor Heavy Metal Exposure of the Population Around a Secondary Lead Smelter: Buildings, Ventilation and Thermal Climate; Berglund, Birgitta; Lindvall, Thomas; Sundell, Jan — editors. Indoor Air, Vol. 2, pp. 177-181; Swedish Council for Building Research, Stockholm Sweden, 1984.

In the course of earlier studies, it had been observed that measuring the concentration of heavy metals in deposited dust is a much better means to characterize the exposure of population groups living close to emission sources than analysing samples of suspended particulate matter. Thus, a simple method has been developed in which house dust collected with a passive sampler is used for a rapid screening of the heavy metal content of dust deposited in houses located near a source. Analysing such dust samples for lead and cadmium permitted the authors to detect the most polluted spots in an area surrounding a secondary lead smelter at Oker, Federal Republic of Germany. The method as well as the results of field measurements carried out in 1982 are discussed.

Selikoff, Irving J. (1975). Investigations of Health Hazards in the Painting Trades: Final Report. City University of New York, Mount Sinai School of Medicine, Environmental Sciences Laboratory, New York NY; December 5 1975.

Keywords: occupational health and safety; painting; air pollution; health hazards;

Selner, John C.; Studdenmayer, Herman (1986). The Relationship of the Environment and Food to Allergic and Psychiatric Illness. Psychobiological Aspects of Allergic Disorders; chapter 6, pp 102 - 146; Praeger Publishers; 1986; Young, Stuart H., Rubin, James M., Daman, Harlan R., ed.

Patients with environmental illness are described, as are methods for diagnosis and treatment. Foods and chemicals can cause acute and chronic symptoms previously viewed as expressions of somatoform disorders by most physicians. Environmental Care Units, isolated facilities which screen out known chemical irritants and sensitizers, are described as useful treatment facilities and as a means of separating patients with ulterior motives from those with environmental illness.

Selye, Hans (1976). Stress: Its Relationship to Man and his Environment. Health Promotion Through Designed Environments, pp 107 - 126; Health and Welfare Canada, Health Programs Branch; October, 1976.

The unprecedented development of technology in this century has created many new sources of stress: pollution of the atmosphere by chemicals and noise, the speed of industrial life, fear of atomic war, etc. The manifestations of stress and techniques for reducing it are presented.

Seppalainen, Anna Maria; Harkonen, Hannu (1976). Neurophysiological Findings Among Workers Occupationally Exposed to Styrene. *Scand. j. work environ. & health*, No. 3, pp. 140-146; 1976.

Keywords: styrene; electroencephalography; encephalopathy; electroneuromyography; mandelic acid; occupational health and safety;

Severs, Richard K. (1980). Air Pollution and Health. from *Environment and Health*, chapter 5, pp. 123-162; Trieff, Norman M. editor; Ann Arbor Science Publishers Inc.; 1980.

Synergistic effects of pollutants have been demonstrated. Air pollution aggravates existing disease conditions or puts at higher risk those predisposed to ill health. Nonspecific measurements of air pollutants that are associated with disease states in humans should be treated as though they were surrogates for causal factors. Little relevance comes from studies that expose humans to only one pollutant at a time. Adverse effects of the following air pollutants on the cardiopulmonary system are chronicled: sulfur dioxide, nitrogen dioxide, photochemical oxidants, carbon monoxide, hydrocarbons, and carcinogens.

Seyal, A. Rashid; Awan, M. Hayat; Mumatz, Hussain (date?). Premature Ventricular Contractions: The Relationship of Synthetic vs Natural Fabrics Worn Next to the Skin. *Clinical Ecology*, Vol. IV, Number 4, pp 149 - 154.

Variations in the frequency of premature ventricular contractions (PVCs) with change in the type of garments were evaluated in 16 subjects who had no evidence of systemic or cardiovascular disease. These subjects, wearing either synthetic or cotton garments, underwent three programmable ambulatory electrocardiograph recordings to achieve a one hour trace within 24 hours. "Within subjects" the reduction in frequency of PVCs was 59 to 100%, whereas "within days" this reduction was 80 to 96%. With pooled data, the major variation in the PVC frequency occurred "within subjects" wearing synthetic garments. However, with cotton clothing this contribution was reduced by 50%. The results also show that the random contribution to PVCs attributed to unavoidable sources is much higher with cotton (50%) than with synthetic garments, the contribution of which was negligible.

Shah, Chandrakant P.; Farkas, Carrol Spindell (1985). The Health of Indians in Canadian Cities: a Challenge to the Health Care System. *Can Med Assoc Journal*, Vol 133, pp 859 - 863; November 1985.

It is well known that Canadian native people living on reserves have high morbidity and mortality rates, but less is known about the health of those who migrated to urban centres. Several studies have shown that these people have high rates of mental health problems, specific

diseases, injuries, infant death and hospital admission. In addition, there is evidence that cultural differences create barriers to their use of health care facilities. The low socioeconomic status, cultural differences, and discrimination that they find in cities are identified as the primary blocks to good health and adequate health care. More epidemiologic studies need to be done to identify health problems, needs and barriers to health care. Federal, provincial and civic governments along with the appropriate departments of faculties of medicine should begin working with native organizations to improve the health of native people living in Canada's cities.

Shain, Martin (1982). Alcohol, Drugs and Safety; An Updated Perspective on Problems and Their Management in the Workplace. *Accid. Anal. & Prev.*, Pergamon Press Ltd., Vol. 14, No. 3, pp 239-246.

The relationship between alcohol, drugs and accidents in the workplace is explored in the context of available evidence, the balance of which suggests that the contribution of substance abuse and misuse to personal injury and property damage losses has been underestimated. Recommendations are made requiring a more complete data base be developed in this area. Necessary steps are seen as: 1) conducting independent prevalence surveys of alcohol and drug use in specific work organizations; 2) relating this information to safety-related behaviour as monitored through observation and interviews and searches of official records, in particular Workmen's Compensation files. Also, it is suggested that the relationships between the management of alcohol and drug related problems and the management of safety be more closely coordinated. The emphasis, however, should be as much on heavy drinking as on alcoholism, and as much on inappropriate prescription use of drugs as on illicit drug abuse.

Shakman, Robert A. (1974). Nutritional Influences on the Toxicity of Environmental Pollutants: A Review. *Arch Environ Health*, Vol. 28, pp. 105-113; February 1974.

Keywords: nutrition; air pollution; toxicity; health effects;

Sharp, Dan S.; Eskenazi, Brenda; Harrison, Robert; Callas, Peter; Smith, Allan H. (1986). Delayed Health Hazards of Pesticide Exposure. *Ann. Rev. Public Health*, No. 7; pp. 441 - 471; 1986.

The delayed health hazards of pesticide use have been difficult to detect. The need continues for surveillance and assessment of delayed health effects from pesticide exposure. Evidence is presented for pesticides causing various cancers, deleterious reproductive outcomes, and subtle neurologic sequelae. Epidemiologic evidence provides the focus, but pertinent animal and clinical research is also presented. Studies of farmers and pesticide users are presented in detail as are studies linking phenoxy herbicides, chlorophenols, dioxins,

and amitrole with mesenchymally derived tumors in Sweden, American chemical manufacturing facilities, New Zealand, and with Vietnam veterans. Other evidence links arsenic and organochlorine pesticides to cancer in humans. Reproductive hazards have been found to result from exposure to DBCP, and there are unconfirmed suspicions that exposure to DDT and the phenoxy herbicides may be associated with congenital malformations. The nervous system has been recognized as a target organ for pesticide toxicity for several decades.

Shearer, Ruth (1983). On Safe Use of Pesticides. author; 1983.

The idea that people can be exposed to pesticides safely is based on a number of faulty assumptions. These include: pesticides registered with the government have passed stringent health testing; a no-observable-effect level can be determined for any pesticide; pesticides are safe when used according to the label; after a pesticide has been excreted from the body, all poisoning symptoms are gone.

Shearer, Ruth W. (1983). Health Effects of 2,4-D Herbicide. author; April 1983.

The author is familiar with the medical histories of more than 30 people who have been acutely poisoned by 2,4-D alone or in combination with other herbicides. Acute symptoms include nausea, vomiting, diarrhea, headache, temporary loss of vision, weakness, burning eyes, sore throat with burning in chest, and difficulty in thinking. Residual effects include impairment, bleeding tendency, concentration and memory problems, and hypersensitivity to non-physiologic chemicals which prevents participation in most modern job environments. Likely carcinogenicity is discussed in detail, including a discussion of recent animal studies.

Sheehy, Noel P.; Chapman, Antony J. (1985). Adults' and Children's Perceptions of Hazard in Familiar Environments. Children Within Environments, Chapter 4, pp 51 - 63; Garling, Tommy and Valsiner, Jaan editors; Plenum Press; 1985.

Children's accidents constitute an endemic social problem of epidemic proportions. This chapter examines the relationship between children's and adults' perceptions of hazard and accident liability. Two studies are described which question the presumption that adults comprehend the child's perspective and empathize with the child which have traditionally governed safety campaigns and educational measures. Future research should consider how hazard identification, tactical responding, and knowledge of personal limitations interact, and how independently and jointly they may precipitate an accident. Extensive references are included.

Shifrin, L. (1987). The Changing Face of Poverty. Toronto Star, November 30, 1987.

The author quotes Statistics Canada figures showing the proportion of Canadians in poverty to be 14.9 percent in 1986. He notes that the percentage of mother-led households among those below poverty-level incomes had increased from 9.9% to 12.7% since 1979. The poverty rate among single-parent families headed by women had risen from 55.4% to 56.1%, while the number of families had continued to grow.

Shy, Carl M.; Creason, John P.; Pearlman, Martin E.; McClain, Kathryn E.; Benson, Ferris B.; Young, Marion M. (1970). The Chattanooga School Children Study: Effects of Community Exposure to Nitrogen Dioxide: 1. Methods, Description of Pollutant Exposure, and Results of Ventilatory Function Testing. Journal of the Air Pollution Control Association, Volume 20, No. 8, pp. 539-545; August 1970.

Elementary schools in four areas of Chattanooga Tennessee USA were selected for a study of the effects of community exposure to nitrogen dioxide. One area, in close proximity to a TNT plant, had high NO₂ exposure, another had relatively high suspended particulate exposure, and two areas served as "clean" controls. The similarity of the economic levels of high NO₂ and control areas and moderately lower economic level of the high particulate area were documented. Ventilatory performance of second-grade school children in the high NO₂ exposure area was significantly lower than the performance of children in the control area. The data suggested that ventilatory performance was adversely affected when an NO₂ threshold was exceeded but that above this threshold no further impairment of performance could be detected.

Silberstein, Samuel (1979). Heating System-Generated Indoor Air Pollution. Energy and Buildings, Vol. 2, No. 4, pp. 271-278; Dec. 1979; Elsevier Sequoia S.A., Lausanne Switzerland.

Keywords: buildings; space heating; air pollution; energy conservation; indoor air pollution;

Silberstein, Samuel (1979). Energy Conservation and Indoor Air Pollution. Energy and Buildings, Vol. 2, No. 3, pp. 185-189; August 1979; Elsevier Sequoia S.A., Lausanne Switzerland.

Keywords: buildings; energy conservation; air pollution; ventilation; indoor air pollution; air exchange;

Silver, Francis (1976). More on the Plastic Problem. *Ecologist*, p. 9, 1976; Environmental Health Assoc. of Greater Washington, Box 3162, Falls Church VA 22043.

Many people experience burning sensations, loss of attention and drowsiness when sitting on plastic chairs. It is possible that incidences of misbehavior by children such as wrecking school buses, hyperactivity and learning difficulties may be all be influenced by exposure to tri-cresyl phosphate in the vinyl seating of the buses, school seats or cleaning agents.

Silver, Francis (1976). Carbon Monoxide. from *Clinical Ecology*, Dickey, Lawrence D., ed.; pp. 269-272; Charles C. Thomas; 1976.

Easily measurable impairment of some of the more delicate senses can be measured at 1% or 2% of hemoglobin bound by carbon monoxide. This blood level can result from five to ten parts per million in the breathed air. At 3% to 5% carboxyhemoglobin, ordinary intellectual functioning starts to suffer in a way that can be measured. Creativity and finer intellectual coordination may be harmed at much lower levels. It is difficult to measure and document such low levels of impairment to intellectual functioning, coordination and creativity, hence the many studies claiming to report "no effect" after heavy CO exposure.

Silver, Roxame, L.; Wortman, Camille B. (1980). Coping With Undesirable Live Events. *Human Helplessness*; Academic Press; Garber, J., Seligman, Martin E.P., editors; pp 71 - 95; 1980.

This is a study of how people cope with stressful life events which deals with the following questions: Are there any reactions that are universally experienced? Do persons who encounter different life crises show similarities in response? Do people progress through an orderly sequence of stages as they attempt to cope? Is it true that with time, people accept or recover from their crisis and move on? What is successful adjustment? Outsiders frequently underestimate the nature and duration of the distress and individuals are likely to hold unrealistic expectations to their own responses which may serve to intensify their distress. Health care professionals must legitimize the feelings and reactions that commonly occur among people who have encountered negative life events in order to assist them to cope. Outsiders must be sensitive to the fact that for most people, an aversive life event is never really forgotten. The transient return of unresolved feelings must not be viewed as an indication of instability or mental illness but as an acceptable way of living with the crisis. Further research is needed. An extensive bibliography is included.

Silverman, Frances (1979). Asthma and Respiratory Irritants (Ozone). *Environmental Health Perspectives*, Vol. 29, pp. 131-136; 1979.

Asthmatics appear to be more susceptible to the effects of air pollutants than nonasthmatics. The present studies were undertaken to examine the effects of exposing asthmatics to ozone concentrations that occur in the environment and indicate that acute exposures to ozone at realistic concentrations in the environment can produce adverse responses in some asthmatics.

Silverman, F.; Corey, P.; Mintz, S.; and Hosein, R.H. (1985). Factors That Influence Assessments of Health Effects of Air Pollution. *Proceedings, 3rd International Conference on Indoor Air Quality and Climate, held in Stockholm, Sweden, August 20-24, 1984. Volume 4: Chemical Characterization and Personal Exposure*, pp. 123-127.

Scientists at the Gage Research Institute, University of Toronto examined the health effects of air pollution in asthmatics and healthy non-asthmatics. The study took into account the fact that the indoor environment may be a modifier of human exposure to air pollution, by using small portable multi-pollutant samplers for nitrogen dioxide, sulphur dioxide, and particulate matter.

Simard, Monique (1986). The Reduction and Rearrangement of Worktime: A Priority and a Challenge: A New Work Agenda for Canada. *Canadian Mental Health Association, Toronto Canada; 1986.*

Policies for reducing and re-organizing worktime are aimed at reducing inequalities and responding to the needs of individuals and groups, especially those traditionally underprivileged. New approaches to worktime can allow us to improve the quality of life at work if introduced with drastic modifications to the organization of work patterns, greater individual and collective control over the work milieu and new technology, and more individual autonomy and creativity at work. The organization of worktime must also respond to the specific needs of women who now make up a significant part of the labour force and who seem to end up sharing unstable jobs most of the time. Women's specific expectations must be taken into account and a real sharing of housework between men and women as a daily reality must be ensured.

Simonsson, B.G. (1986). Allergies and Other Hypersensitivity Reactions to Indoor Pollutants: volume subtitled: Evaluations and Conclusions for Health Sciences and Technology. *Indoor Air*, Vol. 6, pp 57-59; 1986; Swedish Council for Building Research, Stockholm Sweden; Berglund, B.; Berglund, U.; Lindvall, T.; Sundell, J.; editors.

In patients with asthma and hyperreactivity, small concentrations of a pollutant like sulphur dioxide can cause substantial and clinically

important respiratory effects. In acute severe episodes of air pollution, predominantly people with pre-existing heart and lung disease show an increase in mortality. Nitrogen dioxide and formaldehyde can also cause respiratory symptoms especially but not exclusively in hyperreactive subjects like patients with asthma. The compounds can possibly induce airways disease in combination with other pollutants in high concentrations and during exposure for a long time.

Simpson, Michael (1983). Indoor Air Quality and Health Impacts of Energy Conservation: Some Congressional Options. The Library of Congress, Congressional Research Service, Washington DC 20540; updated 09/08/83.

Keywords: indoor air quality; health hazards; energy conservation; research; public policy;

Sivak, Michael (1981). Human Factors and the Highway-Accident Causation: Some Theoretical Considerations. *Accid. Anal. & Prev.*, Vol. 13, No. 2; pp. 61-64; 1981; Pergamon Press Ltd.

Skrzycki, Cindy (1987). Three of Five in U.S. Like Going To Work Survey Finds. *Toronto Star*, October 14, 1987.

A survey for The Conference Board found that three out of five Americans generally like their jobs and are interested in their work. A large majority said they are comfortable with their supervisors and their fellow workers. Only two out of five were reasonably satisfied with their pay cheques.

Small, Bruce M. (1982). Chemical Susceptibility and Urea-Formaldehyde Foam Insulation. National Research Council of Canada, Division of Building Research; February 1982.

Not everyone responds the same way when exposed to pollutants in the air. Initial observations of people exposed to formaldehyde and other gases emitted by Urea-formaldehyde Foam Insulation show a wide variation. But recent evidence indicates that there may be a larger group who are in general more susceptible to low levels of chemical exposures of various kinds. Preliminary reports indicate that some people who are exposed to UFFI gases, and who have no previous indication of chemical susceptibility, may become generally chemically susceptible as a result of the exposure. There is also evidence that for some people other allergic-like sensitivities may also have been triggered by the UFFI gas exposure, even in cases with no obvious history of allergy.

Small, Bruce M. (1982). Environmental Health Factors in Falling Accidents. National Research Council of Canada, Div. of Electrical Engineering, Medical Engineering Section, Ottawa Canada; October 1982.

This study surveys the literature and other sources of expertise to assess the possible role of environmental air pollutants in accidents involving falls in buildings. Evidence is presented which shows that certain pollutants can cause both gross and subtle perceptual and motor ability changes which clearly add to the risk of a falling accident occurring. In addition, it is shown that some people can be much more highly susceptible to common indoor air pollutants than others, and specific precautions to help such persons minimize risk of accidents are outlined. The study recommends that indoor air pollution should be recognized as both a health problem and a safety problem, and a number of followup investigations are suggested.

Small, Bruce M.; Lorimer, Judy; Russell, Peter (1983). Indoor Air Pollution and Housing Technology. Canada Mortgage and Housing Corp., Ottawa Canada; August 1983.

This reference contains a general review of the possible sources of indoor air pollution in residences, including a discussion of the literature describing the health effects of these contaminants. The report also reviews the principles of low-pollution design, with particular reference to an experimental building designed by the author. Available from Technology and Health Foundation, R.R.#1, Goodwood, Ontario L0C 1A0.

Small, Bruce M. (1984). Update on Indoor Air Quality Studies. American Academy of Environmental Medicine, 18th Advanced Seminar in Clinical Ecology Program, pp. 14-15; October 1984.

When populations with common risk factors under exposure to various pollutants are added up, greater than a quarter of the population is seen to be at risk, with the known chemically susceptible population representing only a small part of this group. Those at greatest risk from increased pollutant exposure include pregnant women and young infants, the elderly, other persons with existing respiratory and cardiac problems, the chemically susceptible portion of the allergic population, and smokers.

Small, Bruce M. (1985). Recommendations for Action on Pollution and Education in Toronto: A Report Prepared for the Pollution and Education Review Group of the Board of Education for the City of Toronto. Consultation Paper, May 1985, Board of Education for the City of Toronto, 155 College Street, Toronto, Ontario M5T 1P6.

The evidence presented in this report shows that staff and students are exposed to many pollutants which originate both within their schools and in the neighbouring communities. Many of these pollutants affect brain function, learning ability, behaviour, and hence education. They may also present a general health hazard. The sources of pollution range

from industrial emissions and automobile exhaust outside, to painting, cleaning chemicals, art and science materials, tobacco smoke, and even chalk dust, inside. There is evidence that in some schools the ventilation may have been insufficient to adequately exhaust the pollutants being generated inside. Some students and staff are experiencing acute adverse effects from such exposures, while others do not appear to be suffering, at least in the short term, from the same exposures. Recent evidence indicates that this situation reflects the wide range of vulnerability to pollutants within our population. Some people suffer even at very low levels of contamination, well within previously accepted guidelines for tolerable levels of pollutants.

Small, Bruce M. (1985). Low-Pollution Building Design and Construction. Air Pollution Control Association.

Principles of low-pollution design are reviewed which may be applied in the Canadian climate either for minimizing pollutant levels in conventional housing or for producing specialized housing for persons who may be particularly prone to health problems in the presence of common indoor pollutants. Some of the primary techniques which have been useful in energy conservation are also either helpful or essential in low-pollution construction. For example, a tightly-sealed building envelope is indispensable if random leakage inward through walls brings pollutants from the exterior structure. Outward leakage of humid indoor air through building walls may produce condensation and mould growth, a possible indoor air hazard. It is also important to seal the building envelope below grade, to avoid drawing in soil gases which can include radon, water vapour, mildew odours, and other components.

Other principles highlighted are the provision of clean enough interior construction to allow fluctuations in ventilation and/or filtration without suffering badly deteriorated air during off-times for filtration or periods of repair or emergency. The provision of adequate storage combined with spot ventilation (for economical exhaust of pollutants) are essential features of design at least for chemically susceptible persons and would be of advantage in lowering general pollution levels in other applications.

Smith, Cathy (1985). Letter to the Prime Minister of Canada Concerning Budget Cuts in Environmental and Toxicological Programs, February 13, 1985. Canadian Association for Children and Adults with Learning Disabilities, Maison Kildare House, 323 Chapel, Suite 101, Ottawa, K1N 7Z2.

This letter emphasizes the need for further research on the effects of toxins on human health, productivity and human quality. The author states that there is new mounting scientific evidence showing that the human brain, especially in unborn and young children, is particularly vulnerable to toxic insult, and that such effects appear to be permanent.

Smith, Cyril W. (1987). Electromagnetic Man & His Electromagnetic Environment in Health and Disease. Unpublished presentation to the 5th Annual International Symposium on Man and His Environment in Health and Disease, February 26, 1987, Dallas, Texas, USA;.

This researcher emphasized that a human system can act as an unstable amplifier in response to various stimuli, including a wide range of frequencies of electromagnetic energy. Clinical experiments with exposure of sensitive patients to weak electromagnetic fields indicated 'windows' of frequency response for both initiation and termination of physical symptoms. He emphasized the individual nature of this kind of sensitivity response.

Smith, Lendon (1979). Feed Your Kids Right. Dell Publishing Co., Inc.; 1079.

Illness, hyperactivity, learning problems and stress can be prevented by diet and removal of environmental pollutants.

Smith, Lesbia F.; Victor, Janet (1987). Studies in Assessment of Risk and Management of Environmental Contamination. Occupational Health in Ontario, Vol. 8, No. 3, pp 90 - 101; Summer, 1987; Ontario Ministry of Labour, Occupational Health Branch, Toronto Canada.

Presented in this paper are some of the issues that arise during the response to episodes of environmental contamination and response of the Ministry of Health to local agencies and the public's fear of passive exposure for environmental sources. The essential ingredients required to meet the expectations of scientists, government and community are: the designation of credible local authorities easily accessible to the public; maintaining a steady flow of information, even if it means admitting the temporary absence of data; obtaining expert advice on potential toxicity and having accepted experts available to the community; describing positive and scientifically sound plans to assess potential dangers to health; and keeping the media informed. Cases studies of the PCB spill in Kenora in 1985 and drinking water contamination from the Love Canal and in southwestern Ontario are presented in detail.

Solomon, R.L.; Natusch, D.F.S. (1977). Environmental Contamination by Lead and Other Heavy Metals. Vol. 3. Distribution and Characterization of Urban Dusts: Final Report. US Govt.; NTIS PB-287 153/1SL; July 1977.

This volume describes studies of the sources and distribution of lead and cadmium in the dust and soils of an urban area. The work was conducted as part of a larger effort to identify the sources, distribution, and fate of lead and other heavy metals in both the rural and urban environments. Urban soils and settled dusts are of special interest from the standpoint of human health because it is with these potential carriers of

heavy metal pollutants that humans are most likely to come into contact. It is shown that current urban concentrations of heavy metals in dusts and soils, both indoors and outdoors, could have a significant impact on human health. A field survey of soils and settled dusts was conducted in the small urban area of Champaign-Urbana, Illinois. Objectives of the studies were: to determine where high levels of lead and cadmium are found in urban areas; to determine the sources of lead, cadmium, and polycyclic organic matter (POM); and to determine those physical and chemical properties of lead, cadmium, and POM which determine their environmental impact. Samples collected from several types of locations in an urban environment during the field survey were analyzed for their lead and cadmium content to determine the general distribution of these elements within the area and to identify local patterns of their concentrations in and around selected buildings.

Somers, E. (1987). Making Decisions From Numbers. *Regulatory Toxicology and Pharmacology*, No. 7, pp 35- 42; 1987.

Regulatory agencies require numbers to provide health protection. The manner in which these numbers are derived from animal experiments and human epidemiology is considered together with the limitations and inadequacies of these numbers. Some recent examples of risk assessment in Canada are given including asbestos, drinking water, and indoor air quality. The value of these numbers in providing a measure of the hazard in a wider perspective is stressed, although they can never be the sole determinant of public policy.

Soskolene, Colin L.; Coates, Randall A.; Sears, Abby G. (1986). Characteristics of a Male Homosexual/Bisexual Study Population in Toronto, Canada. *Canadian Journal of Public Health*, Vol 77, pp. 12-16; January/February 1986.

A 1983 survey provided baseline information on the male homosexual and bisexual population of Metropolitan Toronto. The median numbers of different male sexual partners over preceding time periods were about 50% lower than figures reported from United States centres. Population characteristics reported may be important when explaining observed differences in the frequency of AIDS around the world. The authors discuss the difficulty in establishing baseline information for good health research, in a population which is socially stigmatized, and, as a result, not enumerable.

Spasoff, Robert (1987). Special Study of Ontario's Health Care System. Government of Ontario.

The Spasoff Report, unavailable to this research team at the time of writing of the present review, is reported to have concluded that children in families living on welfare get sick because of poor nutrition and hostile living environments. Dr. Spasoff notes that historically, the poor live in the most dangerous areas.

Spedding, D.J.; Hamilton, R.B. (1982). Adsorption of Mercury Vapor by Indoor Surfaces. Environmental Research, Vol. 29, No. 1, pp. 30-41; 1982; Academic Press, Inc.

Keywords: indoor air pollution; metal pollution; mercury adsorption;

Speizer, Frank E.; Ferris, Benjamin, Jr.; Bishop, Yvonne M.M.; Spengler, John (1980). Respiratory Disease Rates and Pulmonary Function in Children Associated with NO₂ Exposure. American Review of Respiratory Disease, Vol. 121, No. 1; pp. 3-10; Jan. 1980.

As part of a long-range, prospective study of the health effects of air pollution, approximately 8,000 children from 6 years to 10 years of age from 6 communities had questionnaires completed by their parents and had simple spirometry performed in school. Comparisons were made between children living in homes with gas stoves and those living in homes with electric stoves. Children from households with gas stoves had a greater history of respiratory illness before age 2 and small but significantly lower levels of FEV₁ and FVC corrected for height. These findings were not explained by differences in social class or by parental smoking habits. Measurements taken in the homes for 24 hour periods showed that NO₂ levels were 4 to 7 times higher in homes with gas stoves than in homes with electric stoves. However, these 24 hour measurements were generally well below the current federal (US) 24 hour outdoor standard of 100 ug/m³. Short-term peak exposures, which were in excess of 1,100 ug/m³, regularly occurred in kitchens. Further work will be required to determine the importance of these short-term peaks in explaining the effects noted.

Spencer, Christopher; Blades, Mark (1985). Children at Risk: Are We Underestimating Their General Environmental Competence Whilst Overestimating Their Performance?. Children Within Environments, Chapter 3, pp 39 - 49; Garling, Tommy and Valsiner, Jaan editors; Plenum Press; 1985.

Adult feelings of responsibility for the child are predicated upon beliefs that the child's competence is less than that of the able-bodied adult; the child is an apprentice in environmental skills. But parents often overestimate their children's performance in traffic. Recent realizations by environmental and developmental psychologists of the child's environmental skills is but one area of the current re-evaluation of a whole range of competencies in perceptual and cognitive domains that is going on in psychology. Ways are suggested in which adults can enhance children's performance in novel environments, with implications for road safety education and practice as it relates to children.

Spengler, John D.; Sexton, Ken (1983). Indoor Air Pollution: A Public Health Perspective. SCIENCE, Vol. 221, No. 4605, pp. 9-17; July 1 1983.

Although official efforts to control air pollution have traditionally focused on outdoor air, it is now apparent that elevated contaminant concentrations are common inside some private and public buildings. Concerns about potential public health problems due to indoor air pollution are based on evidence that urban residents typically spend more than 90 percent of their time indoors, concentrations of some contaminants are higher indoors than outdoors, and for some pollutants personal exposures are not characterized adequately by outdoor measurements. Among the more important indoor contaminants associated with health or irritation effects are passive tobacco smoke, radon decay products, carbon monoxide, nitrogen dioxide, formaldehyde, asbestos fibers, microorganisms, and aeroallergens. Efforts to assess health risks associated with indoor air pollution are limited by insufficient information about the number of people exposed, the pattern and severity of exposures, and the health consequences of exposures. An overall strategy should be developed to investigate indoor exposures, health effects, control options, and public policy alternatives.

Spitzer, Walter O. (1984). A Study of the Health Status of Residents of the Junction Triangle, Toronto: Final Report. Dept. of Public Health, City of Toronto; September 1984.

The purpose of the study was to determine whether the residents of the highly industrialized Toronto Junction Triangle experience adverse health effects more frequently than do residents of a demographically similar but minimally exposed (non-industrialized) area of the City of Toronto, and/or residents of areas adjacent to the Junction Triangle. A comprehensive questionnaire was administered to 110 children and 290 adults in each of: the Junction Triangle, the combined neighbourhoods adjacent to the Triangle, and the comparison tract. The author concludes that there is no difference in the health status of the adult residents of the Junction Triangle and those of the most comparable census tract of Toronto who were not exposed to the emanations from industry identified as being a problem in the Junction Triangle area. However, the study confirmed that the residents of the area experience markedly greater discomfort than people living in other areas of the city and the report recommends that industry guidelines should be established to "sharply reduce imposed factors that diminish the quality of life". It was also concluded that there is enough evidence of unfavourable health experience among children in the Junction Triangle area to warrant in depth, all-inclusive clinical evaluation of their health status.

Sprague, D.E.; Rea, W.J.; Smiley, R.E.; Johnson, A.R.; Lopez de Victoria, A.; Tucker, W.F.; Fenyves, E.J. (1982). Formaldehyde Sensitivity Following Exposure to Building Materials. presented at the 16th Advanced Seminar in Clinical Ecology, Banff, Canada.

Keywords: formaldehyde; building materials; sensitivity;

Squires, Bruce P. (1987). Medical Education To Achieve Health For All. CMAJ, Vol. 136, pp 474 - 475; March 1987.

The "Achieving Health For All" report lends even more credence to the growing concern among many medical educators that the undergraduate curricula of Canadian faculties of medicine do not provide educational programs that realistically reflect the health needs of the Canadian people. They present the concept that health is the absence of disease and that the goal of a medical education is to teach students to recognize and treat disease. Medical schools must devise undergraduate clinical programs that permit students to spend considerably more time in settings where they can acquire a more realistic view of the nature and extent of the major health problems that afflict people and develop the skills to cope with them. They must also allot considerably more time to the teaching of preventive medicine and health promotion and to imbuing their students with the recognition that physicians have a heavy direct and indirect responsibility in helping their patients and communities develop the behaviours that promote health.

St. Lawrence, Ivy (1986). The Twelfth Annual Report of The Ontario Advisory Council on Senior Citizens: For the twelve month period ending March 31, 1986. Advisory Council on Senior Citizens, Government of Ontario, Canada; 1986.

The Advisory Council on Senior Citizens promotes the view that seniors should be encouraged to accept responsibility for themselves to the fullest extent possible, including responsibility for a lifestyle affecting one's own health. However, health promotion strategies should be directed at the entire population, should begin well before retirement, and should be presented in a manner that is relevant to today's seniors. Health promotion should encompass the following: education demonstrating ways and means of illness prevention; nutrition; exercise and activities; wise use of medications and alcohol; and encouragement to become socially active as volunteers and participate in one's community. To a large extent, the current medical model in health promotion works against this objective. Physicians and other health professionals are trained to cure illness. This presents great problems for many of them in working with seniors, for whose illnesses cure is not always possible but who often require health management counselling.

Stebbing, James H., Jr.; Fogleman, Diane G. (1979). Identifying a Susceptible Subgroup: Effects of the Pittsburgh Air Pollution Episode Upon Schoolchildren. Am J Epidemiol, No. 110, No. 1; pp. 27-40; 1979.

Pulmonary function test results on 224 school children collected during and after the Pittsburgh air pollution episode of November 1975 were re-analyzed to determine whether a small subgroup of susceptible children could be defined. Individual regressions of three-quarter second forced

expiratory volumes and forced vital capacities (FVC) on time over the six day study period were calculated, and the distributions of individual slopes for the four exposed and two control schools were compared. Excesses of strong upward trends in the exposed areas would suggest effects of suspended particulate air pollution by indicating significant improvement following the episode. A highly statistically significant excess of strong upward trends in the FVC among exposed students was observed, and was consistent by sex and by school within sex. Approx. 10 to 15% of the students appear susceptible to an average impairment of about 20% of the FVC. The findings are limited by the small number of subjects with strong post-episode upward trends in the FVC, and by lack of validation by replication of the study design, but do suggest that episode levels of suspended particulates induce lung damage, and that this may occur in a small susceptible subgroup. Children with low baseline pulmonary function values, histories of asthma, or with acute respiratory symptoms immediately following the episode were not found to be especially susceptible to these effects of suspended particulates.

Steensberg, Jens (1984). Indoor Climate Problems in Institutions for Children—Practical, Administrative and Policy Perspectives: Buildings, Ventilation and Thermal Climate; Berglund, Birgitta; Lindvall, Thomas; Sundell, Jan — editors. Indoor Air, Vol. 1, pp. 179-180; Swedish Council for Building Research, Stockholm Sweden, 1984.

On the basis of practical indoor climate problems in institutions for children, the study illustrates the administrative and policy perspectives of the decision making process, and some conclusions are attempted on possible ways of improving environmental health decision making. Examples of practical indoor climate problems in institutions for children are given. There are tentative answers to the following questions: How important are value attitudes versus scientific knowledge? Are existing regulations sufficient? How do local and central administrators handle indoor climate problems? Do the responsible politicians influence the course of events? What is the role of the press? Is the pressure from interest groups decisive?

Stephens, R. (1981). Human Exposure to Lead from Motor Vehicle Emissions. Intern. J. Environmental Studies, Vol. 17, pp.73-83; 1981; Gordon and Breach Science Publishers Inc.

A strictly quantitative assessment of the contribution made by car exhaust gas lead to the body lead burdens of young children will probably never be attained. However, there is now available a substantial scientific literature relating airborne lead to lead in humans, especially in young children who ingest deposited and impacted aerosol lead on both food and non-food items. It is clear from blood lead surveys that pre-school children absorb more lead than other sections of the community because young children have a relatively higher metabolic rate and ventilation at rest. The distributions of lead burdens reported

for new-born and pre-school children have been set against levels of lead absorption recently associated with pregnancy problems in women and impaired cognitive/behavioural functioning in children. A study by Needleman et al. in Boston Massachusetts of 158 seven and a half year old children shows that high lead children (greater than 6 ppm) performed significantly less well on a range of I.Q. tests and behavioural ratings than low lead children.

Sterling, E.M.; Sterling, T.D.; Kobayashi, D. Hartel; McIntyre, E.D. (1983). Health and Comfort in Modern Office Buildings: Results of a Work Environment Survey. Theodor D. Sterling Limited, Vancouver B.C.; 1983.

A Work Environment Survey was undertaken to identify the causes of various health and comfort problems experienced by 1106 members of OPEIU Local 153 working in 12 office buildings. Problems were strongly associated with poor ventilation; poor lighting; fluctuations in temperature and humidity; inability to adjust and control ventilation, lighting, temperature, and humidity to individual office comfort needs; and use of video display terminals. Stress at work had a significant influence on health and comfort of employees. Not only were health related symptoms affected by physical factors, but they were also aggravated by inability to participate in the way time at work was allocated, by job insecurity and by poor relations with supervisors. In general there was a complex interaction between the states of environmental conditions and of stressors. Presence of smokers did not show any significant relationship to health and comfort complaints among either smokers or non smokers. But all indices of stress were heightened when smoking was restricted or prohibited.

The authors call for detailed architectural, epidemiological, and air quality studies that could provide regulatory agencies the evidence required to formulate standards which will ensure acceptable working conditions in modern office buildings, especially in structures specifically designed to conserve energy.

Sterling, Elia; Sterling, Theodore; McIntyre, David (1983). New Health Hazards in Sealed Buildings. AIA Journal, pp. 64-67; April 1983; American Institute of Architecture.

Keywords: indoor air pollution; health hazards; housing design;

Sterling, T.D.; Sterling, E.; Dimich-Ward, H. (1983). Building Illness in the White-collar Workplace. Int. J. Health Serv., Vol. 13, No. 2, pp. 277-287; 1983.

The authors describe an increasing incidence of "building illness" among white-collar workers due to the high pollutant content of air in modern energy-efficient office buildings. These buildings are hermetically sealed, mechanically ventilated, and contain many materials that give

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

off a variety of toxic fumes and aerosols. Severe outbreaks of illness have also been traced to ventilation problems in sealed hospital buildings.

Sterling, Theodore D.; Collett, Chris W.; Sterling, Elia M. (1987). Environmental Tobacco Smoke and Indoor Air Quality in Modern Office Work Environments. *Journal of Occupational Medicine*, Vol 29, No. 1; pp 57 - 62; January 87.

Recent attempts to clean the air in modern sealed office buildings appear to have focused on one component of indoor air quality, environmental tobacco smoke. Prohibiting smoking entirely or designating specific smoking areas has been suggested to improve comfort of office workers and reduce acute symptoms of building illness. The effectiveness of such methods, as well as the overall relations of tobacco smoke to indoor air quality is evaluated, based on reviews of a large number of studies of indoor air quality in modern office buildings under normal use and occupancy. Tobacco smoke does not appear to contribute significantly in a build-up of contaminants in offices. Also, in two large series of studies of buildings with health and comfort complaints in the US and Canada, tobacco smoke does not appear to be associated with cases of building illness.

Designation of special smoking areas might remove multiple sources of irritation to smokers and non-smokers alike. On the other hand, the segregation of smokers to specially designated smoking areas may have little effect and may well have undesirable impacts on ventilation performance. Concentrating smokers in designated smoking areas may place an excessive local burden on existing ventilation systems with the result that smoking office workers are exposed to high levels of irritants for short-time periods with any benefits accruing to their non-smoking coworkers. To be of any real use, specially designated smoking areas may require installation and operation of higher volume ventilation systems with more effective air cleaning devices and possible direct venting outside.

Stewart, Donald W.; Toews, John A. (1982). The Chronic Mental Patient: A Selective Annotated Bibliography of the Psychiatric Literature, 1970-1980. Canadian Mental Health Association; August 1982.

This work constitutes part of a national study undertaken to assess the situation of the chronically mentally ill in Canada. It is a selective annotated bibliography of over 400 references from the professional literature on the service system for people with chronic mental disorders.

Stewart, Richard D. (1976). Paint-remover Hazard. *Journal of the American Medical Association*, Vol. 235, No. 4, pp. 398- 401; January 26, 1976.

Keywords: paint remover; indoor air pollution; health hazards;

Stewart, Richard D.; Hake, Carl L. (1976). Paint-Remover Hazard. JAMA, Vol. 235, No. 4, Jan. 26, 1976.

The in-home use of paint removers containing methylene chloride results in the absorption of this solvent, which is metabolized to carbon monoxide. Exposure for two to three hours can result in the elevation of carboxyhemoglobin (COHb) to levels that stress the cardiovascular system. The metabolic formation of COHb continues following the paint-remover exposure, doubling the duration of the cardiovascular stress produced by a comparable COHb level after exposure to CO. Patients with diseased cardiovascular systems may not be able to tolerate this unexpected stress.

Stokes, P.M.; Forester A.; Hale, L.; Hutchinson, T.C.; and Lee, B. (1985). The Cutler Acid Plant Study: A Multidisciplinary Approach to Rehabilitation. Trace Substances in Environmental Health - XV: A Symposium; pp. 340-349; 1981.

The abandoned 100-acre site of operation of a multi-million dollar sulphuric acid factory, on part of the Serpent River Indian Reserve, is still devoid of vegetation over large areas, with a total volume of 60,000 cubic yards of foreign material, mainly in the form of waste piles of pyrite sulphur and calcine and old sulphur storage areas. The surface waters have pH as low as 2.5, with elevated levels of heavy metals, and drain into Aird Bay on Lake Huron. This study identified five rehabilitation options, with associated costs and types of land use for each. The Cutler site is one of several pollution concerns on the Reserve. The Serpent River itself is contaminated with radioactive material from Elliot Lake, and the Aird Bay fishery has been tainted by pulp and paper mill effluent from Espanola.

Stokinger, H.E.; Scheel, L.D. (1973). Hypersusceptibility and Genetic Problems in Occupational Medicine—A Consensus Report. Journal of Occupational Medicine, Vol. 15, No. 7, pp. 564-573, 858, 860; July 1973.

Keywords: hypersusceptibility; occupational medicine;

Stotsky, Karen (1987). Occupational Health and Safety. Integration and Participation, chapter 6, pp 85 - 101; Canadian Advisory Council on the Status of Women; 1987.

As a result of their segregation into a few occupational categories, large numbers of women in the paid workforce are exposed to particular types of health and safety hazards. These include clerical workers who are exposed to VDT radiation hazards, poor lighting, excessive noise, toxic substances, and poor ventilation as well as poorly designed furniture and monotonous jobs. Retail and service workers face hazards associated with bending, lifting and carrying. Hairdressers working

around tonics, dyes, pungent chemicals, and aerosol sprays on a daily basis are susceptible to respiratory problems and skin conditions. Teachers and child care workers are exposed to a variety of communicable diseases, as are nurses. Health care workers are also exposed to radiation, anaesthetic gases, and other toxic substances. Regardless of their place of work, stress is a common problem for most women workers. Factors causing stress include heavy workload, low pay, little job control, lack of recognition, monotonous work, unrealistic deadlines, and the added burden of home responsibilities. A useful bibliography is included.

Stratton, Peter (1985). The Role of the Family in Childhood Risk: The Origins of Competence. *Children Within Environments*, Chapter 9, pp 129 - 144; Garling, Tommy and Valsiner, Jaan editors; Plenum Press; 1985.

Children and other groups such as the aged may be put at risk because of egocentric thinking by those with more power to structure their environments. An evaluation of the dangerousness of particular environments requires a sophisticated understanding of how the environmental characteristics relate to those concerned. A failure to recognize the different perceptual capabilities, decision processes, motivations, etc. of a particular group can lead to the provision of environments which present no great risks to a healthy adult, but which can be extremely hazardous to others such as children. The prospects of reducing risks to children then depend largely on achieving a full and detailed appreciation of how their characteristics relate to those of the contexts in which they must function, and on transmitting this information to those who care for them.

Strickland, Bonnie R. (1982). Implications of Food and Chemical Susceptibilities for Clinical Psychology. *International Journal of Biosocial Research*, Vol.3, No. 1, pp. 39-43; 1982.

Many problems result when the clinician attempts to understand the role of food and chemical substances in behavior problems. The enormous range of individual differences, public misconceptions, lack of research, and the problem of applying old approaches to new situations, all affect how the clinician relates to patients. The public and professionals must become educated as to primary prevention. Clinicians must broaden their repertoire of assessment and treatment techniques.

Subcommittee on Formaldehyde and Air Contamination in Public Buildings (1983). Report of the Subcommittee on Formaldehyde and Air Contamination in Public Buildings. Inter-Ministry Committee of Safety and Occupational Health, British Columbia BC; July, 1983.

Health problems have been linked to poor air quality in a few public buildings in the province, including schools. Twenty school districts have had reports of alleged symptomatic health effects among students,

particularly in portable classrooms. Most of those districts have carried out air testing for formaldehyde and implemented remedial measures. Research is needed into a variety of aspects of the indoor air pollutant problem, including the relationship between short-term exposure to high concentrations and long periods of low level exposure; the effects of humidity on the emission of formaldehyde; and the production and evaluation of controls and standards for mechanical ventilation and air-cleaning systems.

Suedfeld, Peter (1983). Stressful Levels of Environmental Stimulation: Misplaced Emphasis. Stress and Anxiety, Chapter 6, pp 109 - 127; Sarason, Irwin G.; Spielberger, Charles D. ed; Hemisphere Publishing Corporation; 1983.

There are stressful levels of environmental stimulation or environmental load. Social isolation and restricted environmental stimulation can conversely lead to stimulus underload. Comprehensive bibliography is included.

Suess, Michael J. (1984). Exposure and Health Effects of Indoor Air Pollution. Clinical Ecology, Vol 11, No. 3, pp. 130-136; Summer 1984.

Increased concern about indoor air pollution has led to increased effort to determine its sources and how to prevent it. Indoor pollutants may be generated outdoors as well as indoors through a variety of mechanisms. Vehicles and transportation terminals present special problems as do public buildings and medical facilities. More indoor pollution data is needed before adequate assessments of indoor pollution can be made.

Sugita, Kido; Ogihara, Arata; Murabayashi, Hiroshi; Ichikawa, Seichi; Shishido, Masao (1981). The Relationship of Air Pollution and Respiratory Function and Bacterioflora in Pharyngeal Mucus on Junior High School Students. Yokohama Med. Bull., Vol. 32, No. 3-6, pp. 215-225; 1981.

The studies were performed to investigate the effect of multiple air pollutants on the human body. Examination of respiratory function were made by means of the flow-volume curve method and the bacterioflora in pharyngeal mucus were examined. Students from three junior high schools located in a polluted area, a less-polluted area, and a non-polluted area, were tested and the results were compared. The results obtained were as follows: comparing maximal expiratory flow rate in summer, fall and winter, it was highest in summer; measures of central airway respiratory function were the lowest at the school in the most polluted area; streptococci occurrence was high in winter and gram negative bacilli occurrence was high in summer; gram negative bacilli were the lowest at the school in the non-polluted area.

Sullivan, John L. (undated). Office Air Contaminants and Their Sources. University of Western Ontario, Occupational Health & Safety Resource Centre.

Several studies of buildings have been conducted as a result of complaints about air quality. The increased importance of building ventilation due to recent sealing of buildings for energy conservation is highlighted. Typical sources of indoor air pollutants are described. These include sulfur oxides, ozone, pollens, lead, chlorine, nitrogen dioxide, carbon monoxide and dioxide, particles, radon, formaldehyde, asbestos, synthetic fibres, polycyclic hydrocarbons, ammonia, mercury, aerosols and allergens.

Sutton, Robert (1987). Tenant's Inquest Told of Problems Faced by Disabled. Toronto Star, 1987.

The inquest into the death of a disabled woman was told that there were many things wrong with the Ontario Housing Corp highrise where she had lived. The legless widow died as a result of infection from injuries due to falls. Only two of four main entrances were usable by those in wheelchairs, while one ramp was dangerously steep, too narrow, and lacked handrails, according to a study of the building. There were many more access-oriented problems in the building which was originally not designed with the disabled in mind.

Symonds, Alexandra (1986). The Dynamics of Depression in Functioning Women — Sexism in the Family. *Journal of the American Academy of Psychoanalysis*, 14(3); pp 395 - 406; 1986; John Wiley & Sons, Inc.

The incidence of depression in women is at least two times as high as in men, yet there is very little discussion in the psychoanalytic or psychiatric literature which addresses itself to this gender difference in any meaningful way. There is a group of functioning women who are doing fairly well at home and at work, but who have a chronic, underlying depression which has lasted for years. Outward manifestation of this is persistently low self-esteem and a deep sense of insecurity, regardless of external facts to the contrary. Many of this doctor's patients had in common a background which included a brother who was favoured because he was a boy. While this is not the only cause of the increased incidence of depression in women, it intensifies the entire dynamics with which most girls must cope while growing up in a culture of prejudice. Rejection because of gender, being demeaned, trained to serve others, and to repress their own needs create the emotional climate which leads to low self-esteem, insecurity, and depression. This produces lifelong effects in many women which can seriously hinder growth.

Szymusiak, Susan M.; Ryan, Joseph P. (1982). Prevention of Slip and Fall Injuries: Professional Safety. *Professional Safety*, pp. 11-15; June 1982.

Keywords: falling accidents; accident prevention;

Tager, Ira B.; Weiss, Scott T.; Rosner, Bernard; Speizer, Frank E. (1979). Effect of Parental Cigarette Smoking on the Pulmonary Function of Children. American Journal of Epidemiology, Vol. 110, No. 1, pp. 15-26; 1979; Johns Hopkins University School of Hygiene and Public Health.

The authors have investigated the effects of parental smoking patterns on the pulmonary function of children in East Boston, Massachusetts. A crude inverse dose-response relationship was observed between the level of forced expiratory flow 25-75 per cent of forced vital capacity per cent predicted (FEF 25-75 %) of children who never smoked and the number of smoking parents in the household. Compared to children with two non-smoking parents, the level of FEF 25-75 % predicted was 0.156 and 0.355 standard deviation units lower for children with one and two currently smoking parents, respectively. An additional decline in level of FEF 25-75 % predicted was observed for children who themselves had smoked. Smoking children with two smoking parents had an average FEF 25-75 % predicted level which was 0.355 standard deviation units lower than non-smoking children with two smoking parents. These data not only confirm that cigarette smoking by young children and teenagers has direct measurable effects on their pulmonary function, but also show that cigarette smoking by parents has a measurable effect on the pulmonary function of their children which is independent of any direct use of cigarettes by the children.

Tall, F.D. (1987). Air Quality Complaints in University Academic Buildings. Personal interview with Prof. Franklin D. Tall, University of Toronto, December 9, 1987.

Franklin D. Tall is Professor of Mathematics and Chair of the Health and Safety Committee for the Main Academic Building for the Faculty of Arts and Science at the University of Toronto. He notes that classrooms and academic offices are more suitable for chemically susceptible individuals when fresh air is introduced directly into the room, rather than through a central recirculation system which mixes fresh air with large volumes of polluted air. He finds that organizations are extremely slow to respond to air quality complaints, and that provincial inspectors don't know how to deal with indoor air quality problems in non-industrial situations.

Task Force on the Child as Citizen (1978). Admittance Restricted: The Child as Citizen in Canada. Canadian Council on Children and Youth; 1978.

This report discusses the needs of children in relations to Canadian law and social practice and argues for the recognition of children as persons in their own right. Decisions made for and on behalf of children must recognize the individuality of each child's interest and the need for equality of opportunity within the context of the family and the broader society. The existence of true citizenship status for the young will be determined by the degree to which governments and institutions

are mindful of the needs of the young as they pursue their mandates. When the systems lose sight of the purpose for which they were created and the needs of all the citizens they were designed to serve, then it is the weakest members of society who suffer the gravest consequences.

If we are to judge from recent trends, children are, on the one hand, viewed as exotic pets and on the other, as adults in miniature, a consumer market to be ruthlessly exploited. Children are essentially the property of their parents who are free to do with them virtually what they will. The circumstances of children are usually determined by those of their families. Children who are left solely dependent on the economic circumstances of their families have nothing approaching equal access to the full range of goods and amenities society has to offer. Under present circumstances, a health system dominated by an individualized curative approach is unlikely to use its highly specialized personnel to reach out to families to begin active disease prevention and health promotion.

Taylor, Ralph B.; Brower, Sidney (1985). Home and Near-Home Territories. Home Environments, Human Behavior and Environment; Vol. 8; pp. 183 - 212; Plenum Press; New York; 1985; Altman, Irwin and Werner, Carol M., editors.

This is a discussion of the exterior spaces adjoining the home: porches, steps, front yards, back yards, driveways, sidewalks, and alleys. They are important for two reasons: what happens there strongly influences the quality of life in the home; and they represent spaces where the private and personal interpenetrate with the public and shared. The control exercised over these outdoor locations, the responsibility for management of activities, and upkeep, contribute to the immediate society by helping to define and stabilize the standing pattern of behavior on the street block. Different socio-economic groups deal with assaults on this territory in different ways. For instance, residents on a high-crime, low-income street have little material or political means to use as a leverage against possible threats and co-resident distrust is likely, thereby precluding effective group social action against a threat. When people experience less control over this territory than is desired, the discrepancy is viewed as stressful.

Thatcher, R.W.; Lester, M.L.; McAlaster, R.; Horst, R. (1982). Effects of Low Levels of Cadmium and Lead on Cognitive Functioning in Children. Archives of Environmental Health, Vol. 37, No. 3, pp. 159-166; May/June 1982.

Hair cadmium and lead content were related to intelligence tests, motor impairment assessments, and school achievement scores from 149 children aged 5 to 16 enrolled in rural Maryland public school systems. Hair cadmium and lead were significantly correlated with both intelligence scores and school achievement scores, but not motor impairment scores. Significant relations with IQ were obtained after regressing out demographic factors and were observed, even in children within a normal

IQ range. Evidence of different effects of cadmium and lead on cognitive development was obtained. Hierarchical regression analyses suggest that cadmium has a significantly stronger effect on verbal IQ than does lead and that lead has a stronger effect on performance IQ than does cadmium.

Thatcher, R.W.; Lester, M.L. (1985). Nutrition, Environment Toxins and Computerized EEG: A Mini-Max Approach to Learning Disabilities. *Journal of Learning Disabilities*; Vol. 18, No. 5, pp. 287 - 297; May, 1985.

A computerized EEG process called neurometrics has been used to help identify environmental and nutritional factors that may have either a negative or positive effect on child development. A two-pronged strategy involves investigation of environmental factors, the maximization or minimization of which are potentially within the control of parent, child, and educational or health care professional working with a family. The environmental factors which may be of importance for an individual child are determined. Evidence of environmental toxin contribution is routinely assessed when children with learning and behavioural problems are evaluated. If assessment is positive, efforts can be taken to reduce or eliminate the toxins from the body, to minimize contact with the environmental sources of the toxins, and to strengthen overall health by means of a good diet.

Thielebeule, U.; Pelech, L.; Grosser, P.-J.; Horn, K. (1980). Height and Bone Age of School Children in Areas of Different Concentrations of Air Pollution: A Repeat Study Conducted in Bitterfeld and Berlin, East Germany. *Zeitschrift Fuer Die Gesamte Hygiene Und Ihre Grenzgebiete*; Vol. 26, No. 10; pp. 771-774; 1980.

An examination of the height and bone age of school children living in areas of different concentrations of air pollution in comparison with their calendar age was repeated ten years later. There was a substantial drop in dust emission in the highly polluted area of Bitterfeld in the period between the first and repeat examination thanks to considerable investments in the field of energy economy. This was accompanied by a more favourable development of the height and bone age of children living in Bitterfeld than in the preceding period and they are now closer to those of children living in Berlin. It is assumed that these improvements are connected with improvements in the environmental conditions.

Tobin, Richard S. (1986). Health Effects of Airborne Bacteria. Health and Welfare Canada Working Group on Fungi and Indoor Air; March, 1986.

Some adverse human health effects related to bacteria in air are documented. It covers person-to-person transmission of bacterial disease by the air route and problems caused by the excessive concentrations of various types of bacteria in indoor air, particularly after amplification by devices such as humidifiers. It does not include endotoxins nor

Legionella. The most serious consequences on human health are usually observed in especially susceptible populations, but allergic responses ranging from trivial to life-threatening can result in the general population. There is currently little information relating to the presence or absence of airborne bacteria from thermal insulation materials.

Tobin, Richard S.; Baranowski, Eugene; Gilman, Andrew P.; Kuiper-Goodman, Tine; Miller, J. David; Giddings, Michele (1987). Significance of Fungi in Indoor Air: Report of a Working Group. Canadian Journal of Public Health, Vol. 78, Number 2; Canadian Public Health Association; 1987.

Insulation and sealing of homes has reduced the exchange of air between indoors and outdoors and has led in many homes to an accumulation of previously vented pollutants. This report is in response to a request from Consumer and Corporate Affairs Canada for an assessment on the extent to which microorganisms could be a cause of illness in homes.

It was concluded that fungi are common contaminants of indoor air and dust, causing illness such as respiratory and skin allergies, infection of tissues, as well as systemic toxicity. Symptoms from documented exposures have ranged from very minor to severe, including death. Conditions in the home are crucial determinants of the airflora. Fungi thrive in moist conditions, the presence of suitable substrate, and moderate temperatures. Little is known about the types of effects of mycotoxins and fungal volatiles in indoor air. There is a wide range of sensitivities among humans. Recommendations included that research efforts on indoor air quality should consider fungi as an integral and important parameter to study; effort should be made in design, construction, modification, and maintenance of buildings to ensure that exposure to indoor microflora is kept as low as possible; assessment of the health effects of fungi and their components should include data obtained from immuno-toxicity assays and studies conducted in inhalation chambers; efforts should be directing improving human allergy testing; and testing and approval of materials for homes should consider the potential to lead to biological problems. An extensive bibliography is included.

Todd, Dave (1987). Central America Still Uses Banned Pesticides. Toronto Star, July 8, 1987.

Canadian consumers may not be able to fully protect themselves from harmful chemical substances in their food supply, even though the chemicals may be banned in this country. Central America has become a lethal dumping yard for pesticides that North American and European chemical companies are banned from selling or producing. Many foods which are imported to Canada from Central America contain residues of chemicals which are proven to cause cancer and/or central nervous system damage such as DDT, paraquat, Phosvel, DBCP and others.

Toews, J.; Barnes, Gordon (1982). Chronic Mental Disorders in Canada: A Needs Assessment Project prepared for the Canadian Mental Health Association. Health and Welfare Canada; December 1982.

Mental institutions have, in the past few years, been downgraded as the cornerstone of the mental health care system in favour of community and general hospital treatment. However, increasing concern has been raised about the fate of the de-institutionalized patient in the community, including reports of inadequate housing, life skills, activity and work preparation programs, as well as staff overload and frustration at lack of resources. There is a stigma which operates at both the societal level promoting ghettoization, and at the professional level. At the service level, the attitude is often paternalistic. This project included a needs assessment survey of 222 mental health professionals, provincial forums, a national forum, and a selective annotated bibliography. Many recommendations were made concerning the specific aspects of services. These include an intensified advocacy stance on the part of the Canadian Mental Health Association which would include public education, governmental lobbying, and volunteer programs. Studies were suggested into mental health needs in northern and remote areas and the perspectives of clients, friends and relatives.

Toronto Star (1985). '20th Century Disease' Kids Get Separate Special Class. Toronto Star, March 27, 1985.

This newspaper article describes action taken by principal Harry McCosh of Kitchener Collegiate Institute in Kitchener, Ontario, to provide a special low-pollution classroom in this secondary school, for use by students who are hypersensitive to low levels of indoor air pollutants.

Toronto Star (1987). Japan Mistreats its Kids Who've Lived Abroad, Critics Say. Toronto Star, April 18, 1987.

Japanese children who have lived abroad are reported to be treated like aliens when they return home, an experience which can be very stressful. Education Ministry studies indicate that more than 80% of children who have lived abroad experience problems when they return. Parents blame Japan's rigid education system. Educators blame deterioration in the children's ability to speak Japanese. Others feel that differing perceptions of childhood are to blame, and that strong pressure on Japanese children eventually causes frustration and aggression.

Toronto Star (1987). Countdown Acid Rain: Summary and Analysis of the Second Progress Reports by Ontario's Four Major Sources of Sulphur Dioxide. Ontario Ministry of the Environment; February, 1987.

Under Ontario's Countdown Acid Rain program, the four major sulphur dioxide sources are submitting technical progress reports every six

Healthy Environments for Canadians: PART III: BIBLIOGRAPHY

months. Each of the companies has complied with the legal requirements and technical progress continues to be made across a broad front. Detailed financial information is not available for most of the needed technical changes, in view of the fact that much of the developmental work has not been completed.

Toronto Star (1987). Acid Rain Threatens Half-Billion People, UN Study Says. Toronto Star, July 7, 1987.

Acid rain is now a health threat to more than half a billion people, according to a new United Nations report. The report, prepared jointly by the UN Environment Program and the World Health Organization, says that the pollution is endangering some 600 million city-dwellers worldwide. Another billion people are exposed to high pollution levels that result from coal, wood, oil combustion and automobile traffic dust.

Toronto Star (1987). Doctor Urges Basements Be Tested For Radon Gas Linked To Lung Cancer. Toronto Star, September 17, 1987.

Homeowners should be testing for radon that may be sneaking through the cracks and crevices of their basements, says Dr. Michael Moss, a medical professor at Dalhousie University in Halifax. Moss says that radon gas, produced naturally in many parts of the country from uranium in the ground, causes at least 400 lung cancer deaths a year in Canada.

Toronto Star (1987). Ottawa Bans Smoking on Short-Haul Flights: Passengers Who Won't Butt Out Could Face Fines of \$5,000 Under New Rules. Toronto Star, September 16, 1987.

Smoking will be banned on most Canadian airline flights within North America under new federal legislation to take effect on December 9. Fines for smoking will be \$5,000 and airlines will be fined \$25,000 for allowing smoking. Representatives of Canada's two largest airlines say smoke-free air travel has been increasingly popular among passengers.

Toronto Star (1987). Modern Offices Hazard to Health, Civil Servants Say. Toronto Star, July 1, 1987.

Modern office buildings are full of materials, equipment, and furniture that are hazardous to workers' health, according to a statement by the Economists', Sociologists' and Statisticians' Association, a union which represents 2,400 federal public servants. The impact is compounded by the trend toward sealed buildings with internal ventilation systems.

Toronto Star (1987). Poor Kids Shorter or Fatter, Study Says. Toronto Star, June 25, 1987.

A surprising number of children under 5 from low-income families are shorter or fatter than most other youngsters, which could be blamed on chronic under-nutrition, according to U.S. health researchers from the National Centers for Disease Control.

Toronto Star (1987). 24% Feel They're Victims of Discrimination: Gallup. Toronto Star, November 26, 1987.

This popular press report, released just prior to the final draft of the present study, indicates that about one in four Canadians feel they have been discriminated against in one way or another. The figure of 24 per cent, obtained in a recent Gallup poll, is an increase from 19 per cent in a similar poll in 1981. Gallup attributed the increase to a broader awareness of human rights and to legislation resulting from the Charter of Rights and Freedoms. The equality of rights section of the Charter bans any discrimination on the grounds of race, national or ethnic origin, sex, age, or mental or physical disability.

Tosteson, Tor D.; Spengler, John D.; Weker, Robert A. (1982). Aluminum, Iron, and Lead Concentrations of Personal, Indoor, and Outdoor Respirable Particles: from Indoor Air Pollution; Spengler, John—editor; pp. 265-268. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

Samples of respirable particulate matter collected during a personal monitoring study in Topeka, KA, were analyzed for iron, aluminum and lead content. The sampling protocol and instrumentation and described in detail. Lead indoor concentrations (median = 79ng/m³) were found to be less than both personal (median = 112ng/m³) and outdoor lead concentrations (median = 106ng/m³). The indoor, outdoor, and personal levels of iron and aluminum were not significantly different. In addition, it was determined that outdoor respirable particulate mass does not correlate well with the personal or indoor metal concentrations, and that the amount of time spent in motor vehicles is a relatively good indicator of lead exposures. The relationships between indoor, outdoor, and personal lead are discussed in greater detail, with references to supporting evidence from other studies.

Townson, Monica (1986). A New Work Agenda For Women: A New Work Agenda for Canada. Canadian Mental Health Association, Toronto Canada; 1986.

Over the past 15 years, there have been remarkable changes in the economic role of women. The impact on family life is likely to be profound. But policy-makers have been slow to recognize the nature and extent of the changes that have taken place. Policies and programs based on an outdated perception of "the family" and of the relationship between family life and work may be hampering a smooth transition to a new social structure. Unless a different approach is adopted, there could be long-lasting negative consequences for families, children, and society in

general. A new framework is needed within which policies can be developed to address the new reality of family life and work. A new work agenda for women must focus on the particular interests and concerns of women. New patterns of worktime arrangements must make sure that women have access to a fair share of meaningful, dignifying paid work. We must value alternative forms of work such as family responsibilities and voluntarism but also recognize that for too long women have been engaged in these activities without recognition or pay in addition to their paid employment.

Trainor, John; Church, Kathryn (1984). A Framework for Support for People With Severe Mental Disabilities. Canadian Mental Health Association, Toronto Canada; December 1984.

In our collective relationship to people with severe mental disabilities we find ourselves balanced between institutional care, a familiar strategy which we know to be debilitating in the long term, and community integration, a relatively new approach which has been unsuccessful to date because of a lack of full commitment to its promise. The following questions need to be asked: How do people with severe mental disabilities in your community receive the support they need? Do you view the provision of support to people with severe mental disabilities as solely the responsibility of trained professionals? What creative ideas do you have for building new forms of support when existing forms are inappropriate?

Traynor, G.W.; Apte, M.G.; Girman, J.R.; Hollowell, C.D. (1981). Indoor Air Pollution from Domestic Combustion Appliances. US Govt.; June 1981.

Keywords: gas appliances; carbon monoxide; formaldehyde; indoor air pollution; nitrogen dioxide; energy conservation; kerosene; homes; space heaters; stoves; ventilation;

Traynor, G.W.; Apte, M.G.; Dillworth, J.F.; Hollowell, C.D.; Sterling, E.M. (1982). The Effects of Ventilation on Residential Air Pollution Due To Emissions from a Gas-Fired Stove: from Indoor Air Pollution; Spengler, John—editor; pp. 447-452. Environment International, Vol. 8, No. 1/6; 1982; Pergamon Press, Oxford, Toronto; 1982.

The use of indoor combustion appliances can cause an increase in the levels of many different pollutants. The work presented here shows the usefulness of a model for extrapolating environmental chamber results on pollutant emissions from combustion appliances to determine indoor pollutant concentrations in actual residences. In addition, the effects of infiltration, whole-house ventilation, and spot ventilation on pollutant levels are investigated. The results show that a range hood is the most effective means of removing pollutants from a gas-fired range; removal rates varied from 60% to 87%.

Turiel, I.; Hollowell, C.D.; Miksch, R.R.; Rudy, J.V.; Young, R.A.; Coye, M.J. (1983). The Effects of Reduced Ventilation on Indoor Air Quality in an Office Building. *Atmospheric Environment*, Vol. 17, No. 1, pp. 51-64; 1983.

Keywords: human microorganism; office buildings; ventilation; humidity;

Turiel, Isaac (1985). *Indoor Air Quality and Human Health*. Stanford University Press, Stanford CA; 1985.

Air pollution inside buildings may be very much worse than it is outside, according to a recent government report. But efforts to control air pollution have traditionally focused on outdoor air, despite the evidence that urban residents typically spend about 90 percent of their time indoors. This is a readable summary of what is known about the several threats to be found in indoor air, whether in private homes, office buildings, or public places. It assesses various health risks associated with indoor air pollution, and suggests what can be done to lessen the effects of the major contaminants — combustion products from cooking, heating, and cigarette smoke; chemical evaporants from household products and building materials; microbes and allergens; and radon. Two special problems are also discussed: the unhealthy side effects of promoting energy-efficient houses and airtight office buildings, and the complicated regulatory and legal sanctions that impinge on the quality of indoor air. Appendices list relevant U.S. government agencies and describe commercially available air-purification devices. An extensive bibliography is included.

Turner, G; Jackins, H.; (1981). *An Interview with Gill Turner on Young People's Liberation*. Young and Powerful, No. 2; 1981; Rational Island Publishers, Box 2081 Main Office Station, Seattle, Washington 98111, USA.

The author, 17 years of age at the time of the interview, stresses the importance of young people maintaining pride in themselves, and not colluding with many false ideas about youth that are pushed at them by older people. She describes feelings that many young people get, of being superfluous, of not fitting in, and that it doesn't matter what they do. She notes that more obvious forms of oppression of young people, such as physical violence and battering, are still common, but not as universal as they once were. Less obvious, but still critical forms of oppression of young people by adults include: not expecting the best from young people; expecting less than full humanness from them; ignoring their potential; and treating them as less than human. Such attitudes erode a young person's sense of importance, of confidence, and of power.

She adds: "The most terrible thing about oppression (of young people) is not being treated with respect, not being viewed as important, not having real information. This usually results eventually in the young person giving up and allowing her/himself to just drag along with the

patterned world." She emphasizes that because young people are themselves made to feel powerless, while young, they are more inclined to go along with oppression of any other people when they are older. It is part of social conditioning that the society puts upon every person as they grow up, to accept being oppressed and to co-operate in the oppression of others.

Twigge-Molecey, C. (1987). The Impact Assessment Study. Occupational Health in Ontario, Vol. 8, No. 2, pp 53 - 68, Spring, 1987; Ontario Ministry of Labour, Occupational Health Branch, Toronto Canada.

Implementation of the proposed right-to-know laws will put tremendous pressure on companies. While the objectives of the legislation are highly desirable, the implementation period will be difficult, costly, and likely fraught with confusion.

US National Research Council (1969). Effects of Chronic Exposure to Low Levels of Carbon Monoxide on Human Health, Behavior and Performance: Final Report. US Govt.; 1969.

This is an assessment of the state of knowledge concerning the effects on human health of carbon monoxide, one of the major atmospheric pollutants. Presented are well established observations on effects of low levels of CO encountered on urban streets and in traffic tunnels. Research is recommended to broaden present knowledge of physiology and biochemistry of CO toxicity.

US Radiation Policy Council (1980). Report of the Task Force on Radon in Structures. US Govt.; NTIS PB81-166258; August 15, 1980.

Keywords: radon; natural radioactivity; buildings; health risks; NHW

Ulsamer, A.G.; Gupta, K.C.; Cohn, M.S.; Preuss, P.W. (1982). Formaldehyde in Indoor Air: Toxicity and Risk. Air Pollution Control Association; June 1982.

The Consumer Product Safety Commission has received over 3,000 complaints involving consumer exposure to formaldehyde vapor released from building materials. Indoor levels of formaldehyde may range from less than 0.01 to approximately 3 ppm. The effects of formaldehyde on human health derive in part from its properties as a strong irritant and sensitizer. Exposure to formaldehyde can produce a variety of symptoms depending on the mode, duration, and concentration of exposure. They range from statistically significant irritant responses of the eye, nose and throat to prolonged eye, nose and throat irritation, coughing, wheezing, diarrhea, nausea, vomiting, headaches, dizziness, lethargy, irritability, disturbed sleep, olfactory fatigue, and skin irritation. It has been demonstrated to be mutagenic in bacteria, fungi, insects, and mouse lymphoma cells. Of additional concern is the carcinogenic potential of formaldehyde in humans.

United Nations (1973). Report of the United Nations Conference on the Human Environment. United Nations; New York; 1973.

The declaration of the conference proclaims that both the natural and "man-made" environments are essential to people's well-being and to the enjoyment of basic human rights. The protection and improvement of the human environment is the duty of all governments. We must shape our actions with a more prudent care for their environmental consequences. Individuals, communities, enterprises, and institutions must accept equitable responsibility for the task. The report called for protection of non-renewable resources, the halting of the discharge of toxic substances and the release of heat in such quantities as to exceed the capacity of the environment to render them harmless, economic and social development for ensuring favourable living and working environments, integrated and coordinated approaches to development planning so as to ensure that development is compatible with the need to protect and improve environments, education in environmental matters for both young and old, scientific research and development in the context of environmental problems, and international cooperation to spare the earth the effects of nuclear weapons and all other means of mass destruction.

University of Michigan (1965). The Effect of Windowless Classrooms on Elementary School Children. Architectural Research Laboratory, Department of Architecture, the University of Michigan; 1965.

A study was made of windowless classrooms in elementary schools. Teachers like the absence of windows in teaching spaces because they prevent the students from being distracted by outside happenings and extra wall space can be put to good instructional use. Several professional educators, however, have questioned whether the elimination of outside distractions is always something to be desired — an exterior happening may frequently provide a fruitful stimulus to educational activity. Others contend that the curious child in a well-conducted classroom already has an information overload. Windows designed solely as eyes to the outside environment should be quite different in shape and size and location than the traditional window designs. Ideally, they should be ports or apertures in the building shell that will permit the building occupants to have a view of the outside in any desired direction at any desired time.

Unknown (1982). High Aspergillus Count in Hospitals Due to Construction and Renovation. Infectious Diseases, 14; April 1982.

Keywords: indoor air pollution; bacteria; construction; renovation;

Unknown (1982). Lung-cancer Increase Linked to Breathing Diesel-engine Fumes. Medical World News, pp. 22-24; April 26 1982.

Heavy exposure to exhausts from diesel engines may raise American's lung-cancer risk by as much as 40%, according to Dr. Marc B. Schenker of Harvard University. A review was made of health records of 2,519 railroad workers. Men with the highest exposure to diesel fumes had a lung cancer rate nearly 40% higher than unexposed workers.

Unknown (1987). DOE Report Assesses Environmental Impact of Waste Oil Industry: Hazardous Waste News. International Journal of Air Pollution Control and Hazardous Waste Management, Vol. 37, No. 7, pp 780 - 781, 839 - 840; July, 1987.

Many practices which have previously not been questioned as to their environmental or health impact are beginning to be examined. One such practice involves the disposal and re-use of waste oil. It is commonly applied to land, for example, as a road oil or dust suppressant, or is indiscriminately dumped. This creates the risk of contaminating air, water, and soil with substances that pose substantial hazards to humans, animals, and plant life. The hazardous nature of used oil is primarily related to the many contaminants which are known to have carcinogenic, mutagenic, teratogenic, and other chronic and acutely toxic properties.

Valciukas, Jose A.; Lilis, Ruth; Fischbein, Alf; Selikoff, L.J.; Eisinger, Josef; Blumberg, William E. (1978). Central Nervous Dysfunction Due to Lead Exposure. Science, Vol. 201, pp. 465-467; August 1978.

Central nervous system dysfunction was investigated in workers at a secondary lead smelter by means of performance tests. Correlations between test scores and zinc protoporphyrin levels, a biological indicator of lead toxicity, are statistically significant. This correlation should prove to be useful in current efforts to evaluate effects of lead exposure.

Valciukas, Jose A.; Lilis, Ruth (1980). Psychometric Techniques in Environmental Research. Environmental Research, Vol. 21, pp. 275-297; Academic Press Inc.; 1980.

Behavioral changes may be the earliest and only manifestation of neurotoxicity. Moreover, it is well known that extensive brain damage can occur with little or nondetectable clinical neurological deficit. Psychometric techniques now in use in toxicological and epidemiological research have been proposed to assess in an objective manner early manifestations of functional neurological changes that may be due to environmental neurotoxic agents.

Van Assendelft, A.; Forsen, K.-O.; Keskinen, Helena; Alanko, Kari (1979). Humidifier-associated Extrinsic Allergic Alveolitis. *Scand. j. work environ. & health*, No. 5, pp. 35-41; 1979.

Keywords: allergic alveolitis; cool-mist humidifiers; *Thermoactinomyces vulgaris*; *Aspergillus fumigatus*; *Amoeba proteus*;

Van Rensburg, J.P.; Van Der Walt, W.H.; Van Der Linde, A.; Kielblock, A.J.; Strydom, N.B. (1982). Lead Absorption in Distance Runners Exposed to Motor Vehicle Exhaust Fumes. *S.A. Journal for Research in Sport, Physical Education and Recreation*, Vol. 5, No. 1, pp. 21-44; 1982.

In order to establish the extent to which lead in exhaust fumes of motor vehicles is absorbed by athletes training along motorways, blood and urine samples of 94 distance runners from various regions in Johannesburg were analysed for lead and its derivatives. These results were compared with those obtained from a sedentary control group (n=122) and a group occupationally exposed to high levels of atmospheric lead (n=84). The above analyses were extended to erythrocyte count, haematocrit, haemoglobin and mean cell volume with regard to the runners and the sedentary control group. The mean whole blood level concentration of the distance runners was significantly elevated above that pertaining to the control group. In the absence of any irregularities in the haemopoietic pattern, it is suggested that these elevations do not constitute a short-term health hazard. In contrast, the industrial reference group exhibited significantly higher levels for all parameters. The overall impression is, therefore, that the airborne lead levels to which runners are exposed are not high enough to elicit any adverse biological effects. However, the fact that significantly higher whole blood lead levels were found in the runners, points to an "above average" exposure. The long-term or chronic effects of these "above average" levels are not known.

Vayda, Eugene (1986). The Canadian Health Care System: An Overview. *Journal of Public Health Policy*, pp 205 - 210; Summer 1986.

An overview of the history and operation of the Canadian health care system is presented, with comparisons to the performance of the American system. Although the system has worked fairly well, there are concerns: that the system is rigid and inflexible, that costs may be poised to increase rapidly, and that a technology focus coupled with an oversupply of physicians and services may have produced a system which will be unable to adapt to the problems of an aging population, a post-industrial society, and changing human service needs. These concerns are healthy in that they may produce needed changes.

Visual Arts Ontario (1981). The State of the Art: Health Hazards, Conservation and New Materials For the Visual Artist. Visual Arts Ontario; November 1981.

The objective of the conference is to make known the hazardous effects certain commonly used art materials can have and the precautions that should be exercised. Artists are among the highest sufferers of cancer, among other diseases, due to use of solvents, heavy metals, mineral dusts, gases, and other hazardous chemicals such as acids and alkalis. Adequate ventilation, proper use of protective equipment and basic safety equipment must become the concerns of art educators. Elementary, secondary and post secondary institutions must incorporate into their curriculum, health and safety programmes regarding the use of art materials.

Vogt, R.L.; Witherell, L.; Larue, D.; Klaucke, D.N. (1982). Acute Fluoride Poisoning Associated with an On-site Fluoridator in a Vermont USA Elementary School. American Journal of Public Health, Vol. 72, No. 10, pp. 1168-1169; 1982.

On August 30 1980, an outbreak of minor illnesses consisting of nausea and vomiting affected 22 individuals attending a farmers' market at a school. Illness was associated with the consumption of beverages made from school water. Analysis of the water showed high levels of fluoride (1,041 mg/l). The most likely source of the contamination was the school fluoridator, which had accidentally been left on continuous operation.

Voronova, B.Z.; Elkovskaya, E.A. (1980). Effect of the Ventilation Regime on the Functional State of Young Schoolchildren. Gigiena I Sanitariya, No. 6, pp. 31-35; 1980.

In model experiments, an appreciable improvement in air quality occurred when the rate of air supply was increased to 40 or more cubic meters per hour per child, whereas the functional state of the children improved only when the air supply was increased to at least 60 cubic meters per hour per child. The functioning of different bodily systems did not improve simultaneously. The first to improve were the respiratory and cardiovascular functions, followed by that of the central nervous system.

Wadden, Richard A.; Scheff, Peter A. (1983). Indoor Air Pollution: Characterization, Prediction, and Control. John Wiley & Sons, Inc., Box 63, Somerset NJ 08873; 1983.

This text is organized into four parts: characterization, prediction, control and application. The health implications, external and internal contributions, and the measurement of indoor air pollution are described. The current status of prediction techniques is outlined, including areas such as one-compartment models, infiltration estimation, and empirical models. The most common control methods are summarized:

filters, electronic air cleaners, gas filters and traps. Finally, the application of modeling techniques to several typical indoor settings is examined in detail.

In a course run by the authors in Ottawa in late April, 1985, and for which this publication served as a textbook, Messrs. Wadden and Scheff suggested that the samples of "Occupational Standards" which are discussed around Table 1.3 on page 5 of this publication are approximately ten times too high for an application to buildings to be used by a general cross-section of the population, while Table 7.1 on page 136 (ASHRAE) would lead to acceptable standards. The range of chemicals, covered by "Occupational Standards", however, is very extensive. The authors therefore suggest that where no established ASHRAE standards exist, use should be made of the occupational standards, but divided by a factor of ten.

The Toronto Board of Education's Planner-Demographer has suggested in turn that the ASHRAE standard might be divided by ten to produce a "school-safe" standard, but this may be difficult to achieve in practice. Despite practical difficulties, however, it may be necessary to restrict the levels of known sensitizing chemicals to a small fraction of the occupational or ASHRAE standards (e.g. 1/100 or 1/1000).

Wadden, Richard A.; Scheff, Peter A. (1985). Estimation of Activated Carbon Requirements for Controlling Air Quality in an Art Workshop: Chapter 12 (Supplemental) to "Indoor Air Pollution: Characterization, Prediction, and Control" (John Wiley & Sons, Inc., Box 63, Somerset NJ 08873; 1983). Environmental and Occupational Health Sciences, School of Public Health, University of Illinois, Chicago, IL, USA. (treat as copyrighted material not to be distributed except by authors at this address)..

A variety of solvents and other potentially hazardous compounds are typically used in art workshops, including acetone, styrene, toluene, benzene, methyl cellulose acetate, xylene, and others. Methyl cellulose acetate (MCA) was chosen as a representative paint and printing solvent. The authors demonstrate how activated carbon requirements for reducing indoor solvent odours may be estimated for several different control efficiencies.

Wales, Rosemary (1984). Residential Indoor Air Quality; Defining the Problems, Searching For Solutions. Habitat, Vol 27, NO. 1, 1984; Canada Mortgage and Housing Corp., Ottawa.

In recent years, as fuel prices have soared, making our houses increasingly airtight in order to conserve heat has led to an accompanying increase in indoor air pollution. Five of the most troublesome pollutants which can find their way into domestic air are; tobacco smoke, carbon monoxide, formaldehyde, radon, and particles of biological origin. The burning of fossil fuels and wood and the use of products derived from gas

and oil can cause health problems. Chemical susceptibility varies from individual to individual and in the same individual according to age and physical condition. Chemical sensitivity problems related to housing are commonly the result of continuous, cumulative exposure, and often affect more than one bodily system at the same time.

Since current indoor air quality standards generally only apply to the industrial environment, there is a need to determine the safe limits of exposure to indoor air contaminants in living environments. There is also a need to determine the proportion of the population which is most adversely affected, and the groups which are most at risk. The role of government with respect to indoor air quality remains largely to be determined.

Walkinshaw, D.S.; Tsuchiya, Y.; Hoffmann, I. (1987). Exploratory Field Studies of Total Volatile Organic Compound Concentrations in Relation to Sources and Ventilation Rates. ASHRAE; 1987.

Current interest in Total Volatile Organic Compounds (TVOCs) in the indoor environment has been stimulated by Molhave's (1986) postulate of a threshold for human sensory irritation to TVOCs at a concentration similar to those in some European buildings with complaints, and by several U.S. and European studies reporting indoor VOC exposures for many compounds that far exceeded outdoor exposures. The present study examined TVOCs and ventilation rates in eight Canadian settings: an office, an office/library, an office/laboratory, two schools, two hospitals, and a residence. VOCs were collected with a three-part sorbent sampler and then thermally desorbed, with TVOCs quantified by flame ionization detection and characterized by gas chromatography/mass spectrometry. Ventilation rates were estimated from carbon dioxide concentrations and building populations. On the basis of these techniques which were recently developed and are not yet standardized, TVOCs in some locations of four of the eight buildings were found at times to meet or exceed the levels of 2-5 mg/m³ associated by Molhave with mucous membrane irritation and impaired ability to concentrate. However, the isoparaffinic hydrocarbons generated by liquid process photocopiers and identified as the primary TVOC constituent in the air of these four buildings, were well below their proposed occupational threshold limit value. Ventilation rates in all buildings but the schools greatly exceeded the minimum outdoor air proposed for the revised ASHRAE ventilation standard, although rates for the residence on occasion fell below the additional proposed minimum of 0.35 air changes per hour.

Waller, Julian A. (1978). Falls Among the Elderly—Human and Environmental Factors. *Accid. Anal. & Prev.*, Vol. 10, pp. 21-33; 1978; Pergamon Press.

Keywords: falling accidents; elderly;

Wartew, G.A. (1983). The Health Hazards of Formaldehyde. *Journal of Applied Toxicology*, Vol. 3, No. 3, 1983.

Formaldehyde is common in the environment. As an aqueous solution, it causes burns on contact with the skin and eyes, and the vapour is irritating to the eyes and respiratory system. It is a sensitizing agent and may affect susceptible individuals at levels below the current British TLV of 2 ppm. It has been tested for teratogenicity in several animal species by a variety of routes, but was not positive in any of the studies. Inhalation of the vapour has been shown to cause nasal cancer in rats and mice. However, epidemiological studies have failed to confirm whether or not the compound is carcinogenic in humans.

Waterman, Fern K. (1984). Equilibrium of CO Between the Pregnant Mother and the Foetus. *Occupational Health in Ontario*, Vol. 5, No. 1, pp.10-22; January 1984.

The uptake of carbon monoxide by women and its relevance with respect to the foetus is discussed. No threshold can be established below which there is a margin of safety for the foetus.

Webber, Gerald M.B.; Clark, Alan J. (1979). Building Related Home Accidents: A Preliminary Study. *Journal of Consumer Studies and Home Economics*, No. 3, pp. 277-287; 1979.

Keywords: accidents; residential safety;

Weber-Tschopp, Annetta; Fischer, A.; Grandjean, E. (1976). Physiological and Irritating Effects of Indoor Air Pollution Due to Cigarette Smoke. *Ergonomics*, Vol. 19, No. 3, p. 377; May 1976; Taylor & Francis Ltd., London WC2B 5NF England.

Keywords: smoke; density measurement;

Weicker, Lowell P. Jr. (1986). Health Research and National Priorities. *Journal of Medical Education*, Vol. 61, pp 100 - 103; February, 1986.

The nation is in a critical period when national priorities have diverted resources away from the business of a better life and into the never-ending search for better destruction. A \$6.5 billion appropriation for health research and \$50 billion spent over 45 years is compared to a research budget for the Department of Defense in fiscal year 1986 of more than \$40 billion. There needs to be an ordering of public priorities so that sufficient funds ensure a continuing research effort in the cause and prevention of diseases.

Weinstein, Malcolm S. (198?). Health in the City: Environmental and Behavioral Influences. Pergamon Press; 198?.

Today's city has the potential to meet our physical needs better than ever before. Modern technology has been developed to assure clean water and effective sanitation services. But the prevalence of chronic diseases, accidents, and mental disorders has risen dramatically and it has a long way to go to promote the kinds of social environments and healthy lifestyles required to cut down on chronic diseases. Physical problems were given a great deal of early attention while the seeds of social and health problems resulting from overcrowding and poor housing were ignored. Today's health in the city reflects this early priority of physical over social needs. Health inspections need to include standards for social as well as physical environments. The most dehumanizing aspect of today's city is that, as a result of cars and mass transit, we have lost our freedom to move in a human scale. Cities must be designed to meet people's basic spatial and personal needs.

Weiss, Bernard; Spyker, Joan M. (1974). Behavioral Implications of Prenatal and Early Postnatal Exposure to Chemical Pollutants. *Pediatrics*, Vol. 53, No. 5, Part II, pp. 851-856; May 1974.

Some health effects caused by long term exposure to small levels of chemical pollutants may be represented as a slow process similar to natural aging. It is extremely difficult to evaluate whether such differences are significant biological changes and whether they represent a potential for significant functional differences. The brain possesses an enormous reserve capacity. Still, a process that reduces this reserve capacity may, at some time later in life, because of additional losses, make the brain incapable of coping with any additional loads. Subtle neurological deficits and behavioral impairments are common developments of early exposure to lead. These handicaps are often not recognized until the child enters school when he or she exhibits a short attention span, hyperirritability and aggressiveness, sensory and motor impairments. Conventional neurological examinations often fail to detect such instances of minimal brain dysfunction.

Weiss, Bernard (1983). Behavioral Toxicology and Environmental Health Science: Opportunity and Challenge for Psychology. *American Psychologist*, pp. 1174-1187; American Psychological Association, Inc.; November 1983.

Behavioral toxicology is now established as a component of the environmental health sciences. Its rise paralleled recognition that the adverse health impact of environmental chemicals should be gauged by how people feel and function, not solely by death or overt damage. Its compass extends across the total spectrum of environmental chemicals, including heavy metals, solvents, fuels, pesticides, air pollutants, and even food additives. Psychology can help resolve many critical issues in environmental health science and toxicology. Government agencies have to deter-

mine how to predict hazard to humans from animal testing, how to calculate risk on the basis of functional measures. Behavioral toxicologists must design adequate behavioral tests so that, for instance, behavior will be included in food additive testing protocols.

Werner, Carol M.; Altman, Irwin; Oxley, Diana (1985). Temporal Aspects of Homes: A Transactional Perspective. Home Environments, Human Behavior and Environment; Vol. 8; pp 1 - 32; Plenum Press; New York; 1985; Altman, Irwin and Werner, Carol M., editors.

Homes can be viewed as transactional unities. Instead of researching and theorizing about the separate physical, psychological and interpersonal qualities of homes, this approach calls for an examination of homes as integrated unities of physical, psychological and temporal features. Concepts of social relationships need to be studied simultaneously; the home is defined by and gains meaning from the psychological and interpersonal events that occur in it.

Whiston Spirn, Anne (1984). The Granite Garden: Urban Nature and Human Design. Basic Books Inc./Harper Colophon Books; 1984.

In order to create better, more habitable urban environments, it is necessary to understand the natural settings of cities — their air, water, geology, plant and animal life. Various factors of designing an urban ecosystem which works for its inhabitants are described in detail using examples from cities around the world.

White, Frederick A. (1983). Physiological and Psychological Effects of Sound. Our Acoustic Environment, chapter 17, pp 460 - 488; John Wiley & Sons; 1983.

High sound levels can induce responses in the human body that are not specifically related to the auditory system. These physiological responses represent essentially a failure of the human organism to remain a passive system. In most instances, a physiological change is evident only during, or for a short time following, the noise exposure. Some effects studied include the effect of noise on blood circulation, the resistance of the skin to electrical potentials, skeletal-muscle tension, hearing, breathing, and the effect of noise on sleep. There is also evidence that noise effects the gastrointestinal tract, changes the size of the pupils of the eyes, and changes the rate of saliva and gastric secretions.

Psychological and sociological response to noise is difficult to quantify. Objective measurements can be made on specific work tasks, but many reactions to noise seem to be related more to personality factors than to specific noise conditions. A person's psychological response will also be conditioned by whether he or she feels that the noise is a highly personal infringement of a basic right to acoustic privacy. Other in-

fluencing factors are whether the noise is essential, the relationship of the noise to personal activity, the noise's predictability or unpredictability, and frequency of occurrence. There is an aftereffect, especially to unpredictable noise.

An ideal acoustic environment would provide the most pleasant and efficient atmosphere for every individual to work, play or sleep. This is impossible when a large number of individuals are involved. A compromise is required between a rigid no-noise code and a societal attitude that is highly permissive. Advance planning and modern engineering methods can sometimes create strategies for noise reduction. For many years to come, the most complex problem will be the abatement of noise in the residential environment. Community planning of a long-range nature will be required to isolate residential from commercial and industrial areas. But in the final analysis, an individual's acoustic privacy will also depend upon good neighborliness and a genuine consideration for the acoustic sensibilities of others.

Wigle, Donald T.; Mao, Yang; and Grace, Michael (1980). Relative Importance of Smoking as a Risk Factor for Selected Cancers. *Canadian Journal of Public Health*, Vol. 71, July/August 1980, pp. 269-275.

To estimate the proportion of smoking-related cancers that might be prevented if persons did not smoke, a case-control study was performed using questionnaire data gathered from 3924 persons in Alberta who had developed cancer during the period 1971 to 1973. The population attributable risk percent by cancer site among persons who had ever smoked any type of tobacco was: lip: 44% for males; tongue, mouth and pharynx: 84% for males, 37% for females; esophagus: 67% for males; larynx: 84% for males; trachea, bronchus and lung: 88% for males, 52% for females; and bladder: 58% for males, 39% for females.

Wigle, D.T. (1982). Tobacco Smoke and the Non-smoker. *Chronic Diseases in Canada*, Vol. 3, No. 1, pp. 3-5; June 1982.

Keywords: tobacco smoke; passive smoking; indoor air pollution; health effects;

Wigle, D.T. (1982). Tobacco Smoke and the Non-Smoker. *Chronic Diseases in Canada*, Volume 3, No. 1, June 1982, pp. 5-8.

Some 37% of Canadian adults still smoke regularly. Any plan to reduce the extent of smoking must overcome at least two major obstacles. First, smoking is a powerful addiction with failure rates after cessation as high or higher than those for heroin addicts or alcoholics. Secondly, smoking is still socially acceptable (at the time of writing) or at least allowed in most areas. Thus, smokers do not receive enough negative feedback from non-smokers to have a major impact on the problem.

Wilde, Vicki L.; Glunt, Eric K. (1985). Intergenerational Housing: A report prepared for Phipps Houses pursuant to a study undertaken for the American Jewish Committee. City University of New York; January, 1985.

In order to investigate options for meeting housing needs of both students and the elderly, the American Jewish Committee sponsored a study on intergenerational housing. The elderly, because they tend to live on low and fixed incomes, and students, because they choose to devote their time to academic and culturally but not economically productive work, share certain housing and support needs. A proposal was made to move graduate students into under-occupied or abandoned buildings in neighbourhoods with a high percentage of elderly people. The aim for the elderly would be to find housing options that are affordable, age-integrated, supportive of special needs, secure, and help them to remain in their current homes or neighbourhoods. A comprehensive bibliography is included.

Williams, D.L.; Muller, H.K.; Lugg, D.J. (1986). Cell-mediated Immunity in Healthy Adults in Antarctica and the SubAntarctic. *J. Clin. Lab. Immunol.*, No. 20, pp. 43 - 49; 1986.

Cell-mediated responses were studied in 2 Antarctic and sub-Antarctic groups at quarterly intervals over a wintering year, using the cutaneous CMI Multitest. The sub-Antarctic population had levels of responsiveness and hypoergy comparable to other healthy populations in temperate regions. The Antarctic group showed decreased scores and total number of positive responses to antigens, and a significantly elevated incidence of anergy and hypoergy. It is concluded that environment and stress factors in Antarctic expeditions are responsible for the decreased immunological responsiveness.

Williams, John S., Jr.; Leyman, Edward; Karp, Stephen A.; Wilson, Paul T. (1973). *Environmental Pollution and Mental Health*. Information Resources Press, Washington DC; 1973.

This book brings together the existing literature on the subject of environmental pollution and mental health in a manner which provides a general overall view and suggests research priorities on the subject. Among the findings are that the ingestion of some pollutants such as lead, results in direct physical damage to brain tissue. Also discussed are mercury, carbon monoxide, organophosphorous compounds, chlorinated hydrocarbons, and noise pollution. Research priorities include the need for more complete information on the details of physiological effects; the need to explore the idea of mediators between environmental conditions and adverse mental health effects; details about the different kinds, intensities, and periods of exposure; the idiosyncratic nature of individual responses; the interactive effects of exposure to multiple pollutants; and refinement of the measurement of the mental health response.

Wilson, Richard (1978). Risks Caused by Low Levels of Pollution. Yale Journal of Biology and Medicine, No. 51, pp. 37-51; 1978.

Keywords: pollution; health hazards; risk factors;

Winkel, Gary H.; O'Hanlon, Timothy; Mussen, Irwin (1974). Black Families in White Neighborhoods: Experiences and Attitudes. City University of New York; May, 1974.

The increased physical diversification of the suburbs of large American cities during the past 25 years has not been matched by a corresponding diversification in the population residing there. Minority groups have experienced the most serious obstacles in gaining access to suburban living. This study describes the experiences of a group of sixty-two Black families who have purchased homes in racially-integrated suburban neighbourhoods. These people were affluent and well-educated with children who had moved from rental housing. The ease of locating a new home was made most difficult by inadequate information, various forms of racial discrimination from real estate brokers, sellers, neighbours of those selling, and occasionally from a bank. Once they had moved, families reported predominantly positive experiences with their new neighbors. Most of the families were not generally in favour of maintaining social class homogeneity in their new setting.

Winkel, Gary H.; Holahan, Charles J. (1985). The Environmental Psychology of the Hospital: Is the Cure Worse Than The Illness?. The Built Environment, pp 11 - 33; Haworth Press Inc.; 1985.

This paper presents a framework and summarizes evidence bearing on the role that the physical environment plays in the prevention and reduction of psychological and social problems encountered by patients in acute care and psychiatric institutions. Factors that are considered important to preventive strategies include issues such as the spatial layout and design of hospital environments, privacy problems, personal control and independence, information interventions, hospital social relationships, and levels of environmental stimulation. Two case studies are utilized to illustrate these issues within the context of both acute care and psychiatric facilities in a large municipal hospital. Greatest emphasis is placed on the use of the physical environment in the promotion of primary and secondary prevention within tertiary care settings.

Winkelstein, Warren, Jr.; Levin, Lynn I.; Johnson, Kathryn (1982). Health Effects of Particulate Pollution: Reappraising the Evidence. American Journal of Epidemiology, Vol. 115, No. 3, pp. 471-475; 1982; The Johns Hopkins University of Hygiene and Public Health.

Keywords: particulate pollution; lung diseases; particle size;

Witek, Theodore J.; Schachter, E. Neil; Colice, Gene; Beck, Gerald J.; Leaderer, Brian P.; Cain, William S. (1984). Characterization of Irritative Effects from Low-Dose SO₂ Exposure: Sensory and Hyperreactivity Reactions to Sick Buildings; Berglund, Birgitta; Lindvall, Thomas; Sundell, Jan — editors. Indoor Air, Vol. 3, pp. 211-215; Swedish Council for Building Research, Stockholm Sweden, 1984.

Sulfur dioxide (SO₂) is a known irritant that is commonly found in indoor air. The authors measured subjective respiratory complaints in asthmatics and healthy subjects during exposure to low levels of SO₂ in an environmental chamber. Asthmatics complained more of lower airway irritation such as wheezing and chest tightness while healthy persons noted more upper airway irritation such as taste and unusual odour. Exercise increased complaints in asthmatics, but not in healthy persons. In addition, it was demonstrated that there is a relationship between lung function changes and non-specific airway reactivity as assessed by methacholine challenge within the group of asthmatics, suggesting that in some asthmatics airway hyperresponsiveness contributes to the sensitivity to SO₂.

Wolfe, Maxine (1976). Environmental Stimulation and Design; For the Different, Who Are Not So Different. Prepared for publication in Bednar, M.; Barriers in the Built Environment; Stroudsburg, PA: Dowden, Hutchinson & Ross, 1976.

In designing for the developmentally disabled, we also must design for those parts of people or their daily lives that are not disabled. The environments we design, both socially and physically, can help to create disabilities. Research is described which focused on children in a large psychiatric facility. The facility was large in size, having been designed for almost 200 children, but was housing only a fraction of that number. The need to observe and physically contain a small number of children in a large space lead to the program's nature being changed from open and unstructured to highly structured and physically contained. It is suggested that the number of people in a space cannot be thought of separately from the size of the space and the location of areas within that space. The people who are affected by the environment which is being designed can effectively be included in the design process.

It was also found that in most residential treatment settings for the emotionally disturbed, one goal is to foster socially interactive behaviours among the children and between children and staff by means of programming which emphasizes group activities. Yet, one of the clearest findings in behavioural research, both inside and outside of institutions, is that increasing the number of people in a space decreases the amount of social interaction.

Wolfe, Maxine (1986). Institutional Settings and Children's Lives; An Historical, Developmental and Environmental Perspective on Educational Facilities. City University of New York Graduate School; 1986.

A series of interdependent factors — physical, social, political, and economic, must be taken into account if we are understand people/environmental relationships, for example the relationship between learning environments and the behaviours, attitudes, feelings, and development, both cognitive and social, of the children in them. For twelve years, the author and colleagues observed the use of space and educational practices in many schools which were of different vintages and philosophies. It was found that in most of these settings, what occurred on a daily basis did not reflect the goals that teachers, administrators or designers said they were trying to achieve. She points out the institutional qualities which affect children such as structure and routine which take precedence over children as people; and control, authority and the accompanying surveillance which have the underlying assumption that in their absence children will be out of control and which create conformity and lack of privacy. Children are being taught to be passive rather than active creators of their own lives and experiences. Solutions include involving children in design and planning processes — creating alternative ways of creating environments so that the processes themselves will foster the healthy development of children.

Wolkenberg, R.C.; Gold, Calman; Tichauer, Erwin R. (1975). Delayed Effects of Acute Alcoholic Intoxication on Performance With Reference to Work Safety. Journal of Safety Research, Vol. 7, No. 3, pp 104-118; September 1975.

An investigation was undertaken of the aftereffects of alcohol in a simulated industrial work situation. Nine male subjects were administered a series of tests, over a two-week period, that took place before, during, and after an evening of social drinking. Tests involved a variety of coordination requirements commonly required in industry and a questionnaire that measured subjective mood. Delayed effects were observed up to 18 hours after ingestion. These included lengthened reaction time, poor motor performance, and decreased motor sensory skill, as well as inability to manipulate and position without tactile and/or visual facilitation. Some of these and other effects noted could create safety and health problems in a work situation.

Wood, Ronald W. (1981). Neurobehavioral Toxicity of Carbon Disulfide. Neurobehavioral Toxicology and Teratology, Vol. 3, pp. 397-405; 1981; Ankho International Inc.

Keywords: carbon disulfide; behavior; toxicity; central nervous system; peripheral nervous system;

World Commission on Environment and Development (1987). Our Common Future. Oxford University Press, New York NY; 1987.

The World Commission on Environment and Development, headed by Gro Harlem Brundtland, Prime Minister of Norway, was set up as an independent body in 1983 by the United Nations to re-examine the critical environmental and development problems on the planet and to formulate realistic proposals to solve them. This book concludes that the time has come for a marriage of economy and ecology, so that governments and their citizens can take responsibility not just for environmental damage, but for the policies that cause the damage.

World Health Organization (1979). Health Aspects of Wellbeing in Working Places: Report on a WHO Working Group/EURO Reports and Studies. World Health Organization, Regional Office for Europe, Copenhagen Denmark; 1980.

Keywords: occupational health; indoor air pollution; workplace hazards;

World Health Organization (1979). Health Aspects Related to Indoor Air Quality: Report on a WHO Working Group/EURO Reports and Studies. World Health Organization, Regional Office for Europe, Copenhagen Denmark; 1979.

The health aspects of indoor air quality were reviewed by a Working Group convened by WHO in collaboration with the Government of the Netherlands. It was recognized that indoor air quality depended on a number of factors operating simultaneously; these were examined separately and in relation to each other. Pollutants generated outdoors, such as sulphur oxides, nitrogen oxides and carbon monoxide particulates, photochemical oxidants and biologically active particulates, and their relationship to the indoor environment were discussed. A number of pollutants released from indoor sources were identified, such as formaldehyde from particleboard and foamed insulation, radon from the soil or building materials, and fibres of asbestos. Indoor generation of nitrogen oxides, carbon monoxide, carbon dioxide, water vapour and particulates, through human physiological processes or use of unvented gas burning appliances, was also considered.

World Health Organization (1982). Legionnaires' Disease: Report on a WHO Working Group/EURO Reports and Studies #72. World Health Organization, Regional Office for Europe, Scherfigsvej 8, DK-2100 Copenhagen Ø Denmark; 1983.

A meeting of the Working Group was convened by the WHO Regional Office for Europe in collaboration with the Austrian Government. Legionnaires' disease, a multisystem disease with pneumonia as the principal clinical feature, acquired its name following the mysterious outbreak among people attending an American legion convention in Philadelphia in July 1976. In January 1977, a previously unrecognized bacterium was shown to be the

agent responsible for the Philadelphia outbreak. The bacterium was named *Legionella pneumophila* and is now known to be a member of a large family of Legionellaceae. Dramatic common-source outbreaks have occurred in hotels, hospitals, and other establishments in association with contaminated water systems.

World Health Organization (1983). Indoor Air Pollutants: Exposure and Health Effects: Report on a WHO Working Group/EURO Reports and Studies #78. World Health Organization, Regional Office for Europe, Scherfigsvej 8, DK-2100 Copenhagen Q Denmark; 1983.

The assessment and monitoring aspects of exposure to indoor air pollutants were reviewed by a Working Group convened by the WHO Regional Office for Europe in collaboration with the Government of the Federal Republic of Germany. The purpose of the meeting was to review prior work and work underway, and to evaluate the extent to which such work is likely to provide early estimates of actual population exposures and the extent of the health impacts associated with them. Nine main aspects discussed were: population exposure, adverse health effects, concentration values, existing standards, international steering committee on indoor air quality, exposure assessment and priorities, sick building syndrome, sulphur dioxide concentrations and indoor/outdoor relationships, and health effects assessment methodologies and priorities.

World Health Organization (1983). Health and the Environment: Report on a WHO Meeting. World Health Organization, Regional Office for Europe; 1983.

A working group met to discuss how to put strategy into operation aimed at attaining health for all by the year 2000, with prevention, including environmental factors, being an important element. Recommendations include improving and organizing the data base on environmental health, especially in terms of linkages between morbidity/mortality data and environmental exposure; developing national environmental health policies aimed at prevention; developing more comprehensive and standardized risk assessment processes which take into account economic, social, and other factors; conducting research in the biological sciences with emphasis on individual sensitivities and the identification of defence mechanisms against toxic substances that contribute to ill health; and strengthening the components of environmental health in medical training. Public agencies should be encouraged to inform the public about comparative risks and the risks of individual hazards.

World Health Organization (1983). Health Impact of Different Energy Sources: A Challenge for the End of the Century. World Health Organization, Regional Office for Europe; July, 1983.

Current patterns of energy use were reviewed and trends estimated in the use of fossil and nuclear fuels and alternative sources of energy.

The adverse effects on health of different fuel cycles as well as conservation were analyzed. The number and severity of problems of the various sources increases from water, gas and the sun, to oil, nuclear energy, and coal. There are many gaps in knowledge. Sound and cost-effective studies should be made of the effects of energy-related toxicants on people, and on animals in experiments, to reduce current uncertainties about the nature and magnitude of the effects on health. Expanded research should be done to determine exactly how people and the environment are exposed to emissions. The capacity of large atmospheric and aquatic ecosystems to receive pollution should be determined. Research should be done on the genetic and teratogenic effects of energy-related pollutants and the conditions in which people are exposed to them.

World Health Organization (1984). The Effects of the Indoor Housing Climate on the Health of the Elderly: Report on a WHO Working Group. World Health Organization, Regional Office for Europe, Scherfigsvej 8, DK-2100 Copenhagen Ø Denmark; 1984.

A meeting of the Working Group on Indoor Housing Climate Impact on the Health of the Elderly was convened by the WHO Regional Office for Europe in collaboration with the Austrian Government. There has been increasing concern about the effects of the indoor climate of housing on people's health, especially those who may be considered to be at high risk, such as the elderly, the handicapped, and young children. These high-risk groups tend to spend more time at home than do other groups of people and apparently tend, for one reason or another, to be more sensitive or more reactive to extremes in indoor climatic conditions. There is concern not only among health specialists about the possible relationship between reduced ambient air temperatures and rates of ventilation in housing and respiratory diseases. U.S. research cited in the report indicated that indoor airborne viruses and bacteria were the most important causes of disabling illness in that country.

World Health Organization (1984). Summary Report: Working Group on Indoor Air Quality Research: Report on a WHO Working Group. World Health Organization, Regional Office for Europe, Scherfigsvej 8, DK-2100 Copenhagen Ø Denmark; 23 October 1984.

A meeting of the Working Group on Indoor Air Quality Research was convened by the WHO Regional Office for Europe immediately following the Third International Conference on Indoor Air Quality and Climate in Stockholm, Sweden in August, 1984. The Group noted an increased interest in, but little corresponding progress in the determination or estimation of, the overall impact on public health of different pollutants identified in the indoor environment. The Group concluded that changes in building design and engineering practices can have serious consequences for air quality, and that when such changes are introduced, followup studies on the occupants perceptions, reactions, comfort and health should be carried out. Among other things, the

Group also cited indications that the proportion of allergic or hypersensitive individuals in the population is increasing. They recommended that research be initiated into the possible contribution of poor indoor air quality to the causation of allergy and hyper-reactivity, and to the frequency and severity of the responses. Special environmental requirements for the protection of the affected groups should be assessed.

World Health Organization (1986). Intersectoral Action for Health: The Role of Intersectoral Cooperation in National Strategies for Health for All. World Health Organization; 1986.

Intersectoral cooperation has been accepted as one of the guiding principles of an international health strategy. Equity in health cannot be achieved by the health sector alone. It requires contributions from many sectors, in particular agriculture, animal husbandry, food, industry, education, housing, public works, and communications. The health sector has a vital interest in promoting equity-oriented educational policies that give priority to resources for primary education, and pay special attention to the health-related problems of women. Collaboration between health and education can best take place in the school setting. Health education has to be specifically targeted to the different vulnerable population groups. The link between housing and health has been relatively neglected in health strategies but is a major contributory cause of ill health. The health sector also needs to participate in surveillance and safeguards in the area of industrially-generated health hazards. Community participation has been identified as an important means of overcoming sectoral barriers. The priority for health in the national allocation of resources has to reflect the priority accorded to health as a goal.

Wotton, Ernest (1981). Lighting For Education. Ontario Ministry of Education; 1981AD -.

This booklet considers some of the qualities and quantities that must be dealt with to produce good lighting for education. Such things as distracting hum from fluorescent light ballasts is dealt with. Fluorescent lighting's possible effect on hyperactive children is discounted due to lack of conclusive data. It is suggested that students working under fluorescent lighting with very good colour rendering become less visually fatigued than those working under some other forms of fluorescent lighting.

Wotton, Ernest (1982). An Investigation of the Effects of Windows and Lighting in Offices. Health Facilities Design, Department of National Health and Welfare, Ottawa, Canada; May 1982.

This paper presents the results of a study of the effects of windows and lighting on the performance and well-being of office workers. Data was

collected by means of structured interviews with 235 managerial and clerical staff in 6 office buildings in southern Ontario. The buildings had a range of window glazing from 11% to 68%. The respondents had desk locations both near and far from a window. Light meter readings were also taken. No strong relationship was found between work productivity and access to daylight, between work productivity and percentage of glazing, or between job satisfaction and office lighting. Subjective perceptions of bright and dim daylight/electric light do seem to be related to the occurrence of headaches and eyestrain, and to mental well-being. While adequate lighting and access to a window are identified by office workers as important to them, the size of the window and the clearness of the view seem to be relatively unimportant. Within the range studied here, if office lighting is particularly low or high, it can have negative effects on the physical health of workers. However, mental well-being is not so dependent on objectively optimum lighting conditions than on how that lighting is perceived subjectively by workers. An extensive bibliography is included.

Wright, Michael; Kilian, D. Jack (1979). Panel Discussion: Role of the Knowledge of High Risk Groups in Occupational Health Policies and Practices. *Environmental Health Perspectives*, Vol. 29, pp. 143-153; 1979.

When setting out occupational health policies and practices, industry, while being aware of the risks to susceptible workers, must also recognize that there may be dangers to the health of so-called non-susceptible workers as well. Reducing environmental exposure may be the most appropriate solution to the risk problem and cigarette smoking is also a hypersusceptibility factor that must be considered.

Yang, William H.; Purchase, Emerson C.R. (1985). Adverse Reactions to Sulfites. *Can Med Assoc Journal*, Vol 133, pp 865 - 867, 880; November 1985.

Sulfites are widely used as preservatives in the food and pharmaceutical industries. In the United States more than 250 cases of sulfite-related adverse reactions, including anaphylactic shock, asthmatic attacks, urticaria and angioedema, nausea, abdominal pain and diarrhea, seizures and death, have been reported, including 6 deaths allegedly associated with restaurant food containing sulfites. In Canada 10 sulfite-related adverse reactions have been documented, and 1 death suspected to be sulfite-related has occurred. The exact mechanism of sulfite-induced reactions is unknown. Practising physicians should be aware of the clinical manifestations of sulfite-related adverse reactions as well as which foods and pharmaceuticals contain sulfites. Cases should be reported to health officials and proper advice given to the victims to prevent further exposure. The food industry, including beer and wine manufacturers, and the pharmaceutical industry should consider using alternative preservatives. In the interim, they should list any sulfites in their products.

Yassi, Annalee (1982). Occupational Disease and Workers' Compensation in Ontario. Ontario Ministry of Labour; 1982.

While occupational disease is not a newly uncovered problem, its potential magnitude has only recently been widely recognized. The increasing awareness has led to pressure for regulating industrial health hazards and to efforts for cleaning up the workplace. This report addresses the scope of occupational diseases under the headings of physical agents, biological hazards, chemical agents, ergonomic factors, and psychosocial stress. Measuring occupational disease is a difficult task. The various approaches to determining the work-relatedness of disease and decision-making frameworks for compensating victims in Ontario, Canada, the U.S., and Europe are examined. Cancer, respiratory disease, and work-related death are all examined in extensive detail. Recommendations are made concerning compensation policy and procedure in Ontario as well as on external matters.

Industrial disease is under-compensated. Many work-related diseases are not likely to be diagnosed as such because physicians lack awareness of occupational hazards and often neglect to take a comprehensive occupational history. Cases of occupational disease are under-reported because of the absence of records needed to alert workers and their physicians to the possible role of workplace factors in disease. Claims for potentially work-related diseases are almost systematically rejected in the absence of strong supportive research. There is therefore a need for greater resource allocation to research into industrial disease, possibly via a single research agency entrusted with the responsibility of conducting needed studies in industrial disease both for purposes of compensation and prevention. The establishment of a system of independent occupational health clinics would also facilitate the recognition and reporting of occupational disease.

Yocom, J.E. (1983). Industrial Sources of Metals. *NeuroToxicology*, Vol. 4, No. 3, pp. 91-102; 1983.

Any industrial process utilizing or producing inorganic materials releases metals (usually in the form of oxides or salts) into the environment. Metals and compounds containing metals may also be released from industrial processes based on organic materials where metals are used in the process. Significant quantities of metals are released from incinerators burning industrial and municipal waste. Metals are also released from combustion sources such as fuel burning for heat and power (including cars). Selected data is presented from the published literature on atmospheric emission of metals from several heavy industry categories and from incinerators.

Yoshida, Ken (1984). Review of Research Projects. unpublished.

This is a review of Dr. Yoshida's research projects undertaken on pesticides in environment, aerosol science, air pollution and industrial hygiene at the Saskatchewan Research Council and University of Saskatchewan, College of Medicine, since 1967. Projects include Board of Education sponsored studies of air pollution and health at elementary schools.

Yoshida, Ryo; Motomiya, Ken; Saito, Hiroyasu; Funabashi, Shigeru (1976). Clinical and Epidemiological Studies on Childhood Asthma in Air Polluted Areas in Japan. from Clinical Implications of Air Pollution Research, Finkel and Duel, editors; Chapter 13, pp. 165-176; Publishing Sciences Group, Inc.; 1976.

In recent years, the prevalence rate of bronchial asthma in school-children has been on the increase in Japan, and the rates in air polluted areas are higher than the rates in unpolluted areas. It has also been shown that the clinical findings in the air polluted areas are not fundamentally different from those in the unpolluted areas, and that bronchial asthma in the air polluted areas originates in allergic reactions. For symptomatic treatment of children with bronchial asthma in the air polluted areas, it is necessary to quarantine them from air-polluting substances by use of a dust-free room, but it is also fundamental to institute hyposensitization therapy by a positive study on antigens.

Young, G. Stewart; Hagopian; John H.; Hoyle, E. Robinson (1981). Potential Health Effects of Residential Energy Conservation Measures: Final Report, February 1980-February 1981. US Govt.; NTIS PB82-133315; July 1981.

Keywords: residential buildings; ventilation; energy conservation; indoor air pollution; NHW

Yulsman, Tom (1985). The New Threat From PCBs: Special Report: We've Brought The Poisons of a Waste Dump into the Office. Science Digest, pp. 66-68, 84; February 1985.

In 1981 an electrical fire in the basement of an 18 story New York state office building created an indoor environmental disaster. Oil containing PCBs spilled into the blaze and was changed by the heat into a dangerous brew containing, among other substances, dioxin. Contaminated soot was dispersed throughout the building by the ventilation system. Such fires have become fairly common, exposing large numbers of people to reproductive system damage, immune system damage, cancer-causing agents.

Zamm, Alfred V.; Gannon, Robert (1980). *Why Your House May Endanger Your Health*. Simon & Schuster, New York NY, 1980.

There are many environmental problems within buildings: chemical pollution, poisonous vapours, electrical particles, electromagnetic waves, dust, fumes from waxes and disinfectants. Hyperactivity, respiratory and digestive upsets, skin rashes, even symptoms of mental retardation can result from the presence of noxious chemicals indoors.

Zube, Ervin H. (1984). *Environmental Evaluation: Environment and Behavior Series*. Cambridge University Press; 1984.

Evaluation studies are intended to provide information to improve the quality of decision making with reference to environmental management as well as to change and modification. Whenever growth occurs, modification of the environment will occur, sometimes under the jurisdiction of federal or local policies relating to housing, recreation, transportation, etc. Evaluation studies provide a means for assessing the suitability of those policies through studying the success or failure of the subdivisions, apartments, parks, playground, and subway stations that result from their implementation. Three stages are important for assessment: the inventory or evaluation of existing environments, consideration of alternatives, and after the plan is implemented, the design or management program which evaluates the modified environment.

Zuckerman, Diana M.; Zuckerman, Barry S. (1985). *Television's Impact on Children*. *Pediatrics*, Vol. 75, No. 2, pp 233 - 240; February 1985.

Television has a major impact on children's knowledge, attitudes and behaviour. Research has demonstrated the association between television viewing and four areas: children's aggressive behaviour; racial and sex-role stereotypes; decreased interest in reading and school activities; and poorer health habits and attitudes. Methodological limitations make it difficult to draw firm conclusions about a causal relationship between television viewing and children's behaviour. Representative studies in these four areas are reviewed, important methodological concerns are pointed out, and conclusions from the research findings are drawn. The implications of the data for pediatricians and other health professionals are discussed. The modest statistical associations between television viewing and children's cognitive or behavioural problems may be masking the more extreme responses of small groups of especially vulnerable children. The identification of a high-risk subsample of children has been virtually ignored in the research that has been conducted thus far.

Zweideinger, Ruth A. (1977). *Organic Emissions from Automobile Interiors: Final Task Report, July 1975 to January 1976*. US Govt.; December 1977.

Keywords: volatile organic compounds; automobile interiors; vinyl chloride; gas chromatography; mass spectroscopy; monomers;

- Abel, C.B. A67,A68,B1
 Adam, B.D. A46,A47,A48,A79,B2
 Adams-Taylor, S. B138
 Adelman, R.C. A29,B2
 Adkinson, N.F. A55,B3
 Ahlberg, M.S. B178
 Aitken, R. B120
 Akimenko, V.V. A82,B3
 Alanko, K. B224
 Alarie, Y. B88
 Alfano, D.P. B159
 Allen, G.A. B155
 Allen, J.R. B73
 Allen, M. A47,B4
 Alphey, R.S. A4,B4
 Altman, I. A7,B4,B230
 Ames, B. A57,B4
 Ammann, H.M. A40,B5
 Amster, M. B89
 Andersen, I. A53,A55,A81,B3,B5,B6
 Anderson, D.E. A19,B7
 Anderson, J. B180
 Anderson, J.M. A39,B7
 Anger, W.K. A83,B7
 Angus, D.E. A71,B8
 Aoyama, H. B143
 Appleyard, D. A21,A67,B8,B9
 Apte, M.G. B73,B219
 Arehart-Treichel, J. B177
 Arlien-Soborg, P. A13,B9
 Armstrong, R. B4
 Aronow, W.S. A54,B10
 Asher, J.K. A20,B10
 Astle, A. B89
 Aurand, K. B190
 Awan, M.H. B191
 Axelson, O. A6,A13,B10,B11,B83
 Bach, B. A83,B11,B140
 Bakacs, T. A21,B12
 Baker, J.A. B76
 Baker, L.H. B36
 Baldassare, M. A19,A67,B12,B68
 Baldwin, D.G. B113
 Ball, I. A27,B12
 Ball, R. A12,B13
 Balser, D. A32,A33,A34,A35,B13
 Balster, R.L. B172
 Bamberg, D. B167
 Baranowski, E. B215
 Barchas, J. B40
 Barchas, P.R. B40
 Bardana, E.J.J. A13,B14
 Barker, R.H. A55,B14
 Barnes, G. B216
 Barnes, J.M. B14
 Baroncelli, P. A61,B15
 Barr, D. B25
 Barrett, P. B145
 Bartram, J. B127
 Bates, D.V. A51,B15
 Bates, L.A. B23
 Battisti, S. B15
 Batts Young, B. A13,B15,B103
 Bauman, K.A. A49,B16
 Beaconsfield, P. B16
 Beall, J.R. A55,B16
 Beard, R.R. B118
 Beck, G.J. B234
 Becker, H.A. B16
 Becker, R.O. A61,A62,B16
 Befekadu, F. B17
 Beirness, D.J. B130
 Bell, I.R. A64,B17,B18,B168
 Bellanti, J.A. A52,B18
 Ben, M. B188
 Benard, J.M. A16,B18
 Benson, F.B. B194
 Berglund, B. A12,A82,B19
 Berglund, U. B19
 Berk, J.V. A5,B19,P28,B91,B92
 Bertell, R. A62,B20
 Berwick, D.M. B20
 Besch, E.L. A54,B21
 Biersteker, K. B109
 Billings, C.E. B21
 Bishop, Y.M.M. B65,B202
 Bjorklid, P. A22,B21
 Biades, M. B202
 Blumberg, W.E. B223
 Blume, J. B83
 Blume, K.A. A8,B22
 Blumenthal, J.A. B22
 Blumenthal, M.D. A4,A29,B22
 Boegel, M.L. B92
 Boegman, R.J. A56,B23
 Boer, J. B23
 Bokina, A.I. A82,B23
 Bonham, G.S. A26,B24
 Bonica, J. B40
 Borland, B.L. A10,B24
 Borlaug, N. B16
 Borrows, P. A12,B24
 Bosnjakovic, B.F.M. B102
 Boulding, E. A27,A28,B25
 Bourne, P.G. A37,B25
 Bowen, R.P. B38
 Bowers, E.J. B88
 Bowles, A.M. B25
 Boyce, D.E. B88
 Boyes, W.K. B178
 Boylen, G. B89
 Bozzelli, J.W. A9,B25
 Bradley, E. A49,A84,B26,B27
 Bradley, J. A22,A81,B27
 Bradstreet, J.W. B89
 Brahim, S.A. B147
 Brasser, L.J. B41
 Breen, T.J. B172
 Breysse, P.A. A5,A6,B27
 Bridbord, K. A55,B27
 Brilliant, L.B. B64
 Broadbent, D.E. A61,B28
 Broadbent, M.H.P. B28
 Brodsky, C.M. B35
 Brooks, A.G.F. B135
 Brooks, C.R. B105
 Brostoff, J. B186
 Brower, S. B213
 Brown, K.S. B89
 Brubaker, P.E. B27
 Brundrett, G.W. A6,B28
 Budd, R.A. A61,B28
 Budnitz, R.J. A6,B28,B91
 Burg, M.M. B22
 Burge, P.S. B151
 Burnett, R.T. B70
 Butler, J.R. B53,B117,B150,B170
 Cain, W.S. A53,B29,B234
 Calabrese, D.V. A77,A84,B29
 Calabrese, E.J. A29,A41,A56,A84,B30
 Calles, P. B192
 Calle, E.E. B185,B186
 Camacho, T. B80
 Camp, J.E. A16,B30
 Campbell, D. A44,B31
 Campbell, J. A65,B31
 Campbell, J.S. A6,B31
 Campbell, K.I. A14,B32
 Canada Mortgage and Housing Corporation A29,A30,B32

- Canadian Advisory Council on the Status of Women A69, B33
Canadian Association for Children and Adults with Learning Disabilities B33
Canadian Centre for Occupational Health and Safety A15, B33
Canadian Home and School & Parent-Teacher Federation A26, B34
Canadian Mental Health Association A33, A34, A43, A69, B34, B35
Cannon, M. A16, B35
Carp, F.M. A31, B35
Carson, B.L. B36
Cassel, J. B36
Cassidy, M.W.A. A4, B38
Catalano A68, B37
Cavender, F.L. B160
Chai, R. B89
Chalupiak, S. A18, B37
Chamberlain, D. B50
Chapman, A.J. B193
Charpin, J. B153
Checcucci, A. B15
Chenier, N.M. A15, B38
Chester, E.H. A14, B38
Chinn, S. B135
Chipman, C. B38
Chisolm, J.J. A25, B38
Chown, G.A. A6, B38
Christensen, K.E. A18, B39
Christensen, L. A56, B39
Church, K. A44, A84, B39, B156, B219
Ciaranello, R. A64, B40
Clabo, D.A.J. B74
Clark, A.J. B228
Clark, D. A48, A49, A80, B40
Clark, E.J. A32, B40
Clark, J.D. B135
Clarke, N.A. B158
Clary, J.J. B31
Clayton, P. B55
Clelland, R.C. B88
Coalition for Gay Rights in Ontario A47, B41
Coates, R.A. B201
Coerr, S. B161
Cohen, M.A. B41
Conn, M.S. B221
Colice, G. B234
Collett, C.W. B207
Colley, J.R.T. A51, B41
Colligan, M.J. B42
Collishaw, N.E. A14, B42
Colome, S.D. A5, B42
Combs, P. B15
Combs, E.R. A30, B42, B106
Committee on Indoor Pollutants, N.A.S. A55, B43
Comptroller General of the U.S. B43
Comstock, G.W. B86
Conright, K.L. B115
Consumer Federation of America A5, B43
Contenta, S. A38, B44
Cooney, P.A. B137
Cooper, C.L. A17, B44, B50
Cooper, W.C. A84, B45
Corey, P. B93, B196
Cote, R.W. B108
Couchman, R. A35, A36, B45
Coulter, E.J. B138
Coye, M.J. B220
Craighead, I.B. B135
Crawford, L. B89
Crawford, T. A70, B45
Crawshaw, G.H. B46
Creason, J.P. B194
Crocker, L.H. B157
Cullen, J. A76, B46
Culot, M.V.J. B46
Czerski, P. B28, B102
D'Arcy, C. A69, B46
Dadd, D.L. A8, B47
Daglish, S. A26, A44, A57, B48
Dahl, A.R. B48
Dainty, E.D. A14, B49
Damstra, T. B49
Daniell, H. A20, B49
Danner, R.M. B158
Darlington, L.G. A76, B49
Dart, D.M. B21
Davey, P.G. B50
Davidson, M.J. A32, B50
Davie, J.W. B22
Davies, D.J. B136
Davies, F.L. A24, A27, B50
Davis, K. A76, B51
Dawkins Brown, O. B171
Day, D.M. A43, B51
Day, J. A77, B31, B51
DeLeon, L.R. B117, B170
Dean, A. A72, B52
Dean, J.H. A83, B52
Demers, L.M. B129
Dershewitz, R.A. A4, B52
Diamond, P. B53
Dickey, L.D. A44, B53
Didriksen, N. A10, A64, B53, B54
Dikshit, M.B. B100
Dillon, C. A78, B54
Dillworth, J.F. B219
Dimich-Ward, H. B206
Dimmick, R.L. A19, B54
Dinman, B.D. B118
Dockery, D.W. B54
Dodge, R. A5, B55
Dohrenwend, B. A72, B55
Doi, T. B105
Donelson, A.C. B130
Dossing, M. B152
Dougherty, R. B70
Douglas, C.J. A47, B55
Dowdle, E.B. B124
Drader, D. A29, B56
Dravnieks, A. A82, B56
Drews, M. B190
Ducharme, R.T. B100
Dudney, C.S. A53, B57
Duel, W.C. B67
Duffee, R.A. B89
Duhl, L.J. A21, A73, A81, B57, B58
Dunlop, M. A42, B58
Durlak, E. A4, A41, A45, B58
Edelstein, B.A. B59
Edling, C. B11
Edwards, D.D. A61, B59
Ehrlich, R. A52, B60
Eliff, A.W. B146
Eisdorfer, C. B61
Eisenbud, M. A67, B60
Eisinger, J. B223
Eksler, N.D. B23
Ellinson, L. A83, B61
Elkovskaya, E.A. B225
Elliott, G.R. B61
Ellis, H.V. B36
Elwood, J.M. A61, B62
Engen, T. A55, B62
English, K. A36, B62
Epstein, S.S. A62, B63
Erman, T. B26
Eskenazi, R. B192
Ettema, J.H. A40, B63

- Evans, J.G. B164
 Evans, J.R. A77,B63
 Faeder, E.J. B160
 Fanos, I. B26
 Farah, J.P. B18
 Farber, M. A38,B64
 Farkas, C.S. B191
 Faust, H.S. A8,B64
 Fawcett, I. B151
 Fein, G.G. B64
 Feingold, M. B89
 Fenters, J.D. B80
 Fenyves, E.J. B171,B203
 Ferguson, R. A75,B65
 Ferrario, C. B40
 Ferris, B.G. A5,B65,B121
 Ferris, B.J. B65,B202
 Feuerstein, M. A65,B66
 Figner, C.A. B66
 Ficula, T. B71
 Figley, D.A. B128
 Finberg, L. A25,B66
 Findlay, J.C. B60
 Fine, A. B138
 Finkel, A.S. A52,B67
 Finkel, J.M. A55,B67
 Finn, P. A46,B67
 Finn, R. A76,B67
 Fischbein, A. B223
 Fischer, A. B228
 Fischer, C. A72,B12,B68
 Fischer, C.S. A67,A72,B68
 Fishbeck, W.A. A15,B69
 Fisher, A.A. A8,B69
 Flachsbar, P. B155
 Fleishman, J. A16,B69
 Fleming, G.M. B38
 Florey, C.V. B135
 Fogleman, D.G. B204
 Foote, R.S. B69
 Forbes, W.F. B89
 Forester, A. B208
 Forsen, K.O. B224
 Franck, K.A. B69
 Frankenhäuser, M. B121
 Franklin, C.A. A22,A58,B70
 Frel, D. B127
 French, J.G. B27
 Friedman, M. A5,B70
 Fugas, M. B184
 Fukushima, K. B99
 Funabashi, S. B242
 Furlong, W.J. B158
 Gannon, R. B243
 Gardell, B. B121
 Gardner, D.E. B60
 Garling, T. A50,B70
 Gaudert, P.C. A6,B70,B106
 Gay, B.J. B27
 Gaynard, P. B153
 Gelfand, D.M. A66,B71
 Geller, E.S. B71
 George, E.L. B32
 Gerdes, K. A1,B72
 Gershon, S. A58,B72
 Gibson, B. A13,B72
 Giddings, M. B215
 Giddings, M.J. A14,A15,B73
 Gilka, L. B73
 Gilman, A.P. B215
 Girman, J.R. A5,B73,B219
 Glunt, E.K. B106,B232
 Godderz, W. B146
 Godish, T. A41,B73
 Golberg, L. B31
 Gold, C. B235
 Goldberg, S. A66,B74
 Goldman, R.F. B21
 Goldsmith, C.H. B158
 Goldsmith, J.R. B79,B88
 Goldstein, B.D. B135
 Gonzalez, M.G. B115
 Good, B.W. A5,B74
 Gordis, L. A24,B74
 Gordon, M. A29,B75
 Gosnell, D. A10,A11,A74,B75
 Gottlieb, N. B26
 Gottlieb, N.H. A73,B76
 Goven, A. B53
 Grace, M. B231
 Grandjean, B. B9
 Grandjean, E. B228
 Grandolfo, M. B15
 Gravesen, S. A9,A54,B76
 Grayson, D.A. B67
 Green, G.H. A63,B77
 Green, K.B. A60,B77
 Green, M. B129
 Greenwald, P. B46
 Greer, N.R. B78
 Gregory, J. A32,B78
 Griffith, E.E.H. A38,B79
 Griffith, E.J. B79
 Grimaud, C. B153
 Grimsrud, D.T. B79
 Groll-Knapp, E. B80
 Grosser, P.J. B214
 Grundy, R.D. B63
 Guebaly, N. B129
 Guidotti, T.L. A74,B79
 Gunnoe, C. B145
 Gupta, K.C. A53,B79,B221
 Gustavson, J. B11
 Gyntelberg, F. B76
 Haan, M. A70,B80
 Hadfield, A. B110
 Hadley, W.M. B48
 Hagopian, B242
 Haider, M. A82,B80
 Hake, C.L. B208
 Hale, F.A. B16
 Hale, I. B208
 Hall, C. A29,A30,B81
 Hall, M. A44,A45,B81
 Hall, R.H. B94
 Halton, D.M. A16,B81,B82
 Hambraeus, A. A54,B82
 Hamburg, B. B55
 Hamilton, R.B. B202
 Hancock, T. A77,A85,B82,B83
 Hane, M. A13,B10,B83
 Hanvey, L. B83
 Hanzai-Kashi, A. B42,B106
 Harding, D.H. B83
 Harkonen, H. B123,B191
 Harper, D. B129
 Harries, M.G. B151
 Harris, M. B131
 Harrison, J. A5,B84
 Harrison, R. B192
 Hart, D.R. A35,B84
 Hartley, R. B136
 Harvey, W.R. B74
 Hasebe, A. B107
 Hashimoto, T. B108
 Hastings, A. B106
 Hathaway, W.E. A61,B84
 Health and Welfare Canada A61,B85
 Heath, L. A47,B86
 Hein, K. B105

Helsing, K.J. A5,B86
Henderson, A.S. A72,B87
Henzi, H. A9,B87
Hernandez, T.L. B181
Hernberg, S. B123
Herndon, B.L. B36
Hertzman, C. A69,B87
Hess, C.T. A6,B88
Hexter, A.C. A52,B88
Hickey, R.J. B88
Hijazi, N. A82,B89
Hildingson, O. A6,B89
Himelein, M.J. B147
Hirdes, J.P. A29,B89
Hirschman, S.Z. B89
Hisashige, A. B143
Hoff, H. B90
Hoffmann, L. B227
Hogstedt, C. B10,B83
Holahan, C.J. B233
Holland, J. A29,B90
Holland, W.W. B90
Hollands, J. A16,B90
Hollenberg, C.H. B90
Holler, H. B80
Hollowell, C.D. A55,B19,B28,B73,B91,B92,B137,B219,
B220
Holma, B. B92
Holt, J. A27,B92
Hooker, C.A. A81,B93
Hopkirk, G. A18,B93
Horn, K. B214
Horst, R. B213
Horton, R.J.M. B21
Hosein, H.R. A5,B93
Hosein, R.H. B196
Houle, M. B66
House, J. B105
House, R.V. B52
Hoyle, E.R. B242
Hrustic, O. B184
Huggins, H.A. B94
Hughes, J.P. B162
Hume Hall, R. B94
Humiston, K. A21,A64,B94
Hutchinson, P. A43,B95
Hutchinson, T.C. B208
Hutchison, P. A43,B95
Hutchison, P.A. A39,B96
Ibsen, K.K. A9,B96
Ichikawa, S. B210
Illich, I. B96
Imbus, H.R. A5,B97
Ingraham, L. B97
Inman, E.E. B158
Innes de Neufville, J. A32,A84,B97,B98
Insel, P.M. A43,B98
Irwin, J. A35,A46,B99
Ishikawa, S. A58,B99
Israelson, D. A58,B99
Iversen, M. B124
Iyer, E.M. A63,B100
Izumi, K. B100
JML B100
Jackins, H. B220
Jacko, M.G. A20,B100
Jackson, G.M.A. B26
Jacobson, J.L. B64
Jacobson, S.W. B64
Jaeger, R.J. A5,B101
Jaffe, L.S. A52,B101
Jaffe, P. A33,B101
James, R.H. B67
Jammel, H. A62,B102
Jarvis, B.B. A54,B103
Johansson, I. B19
John, H. B242
John, H.H. B135
Johnson, A.R. B171,B203
Johnson, B. A42,B103
Johnson, K. B233
Johnson, K.A. A43,B103
Johnson-Lussenburg, C.M. A54,B104
Johnston, A. A81,A84,B104
Jonassen, N. A54,B104
Jones, M.R.L. B28
Jordan, S. B172
Kagamimori, S. B108
Kagawa, J. A40,B104,B105
Kagen, H.P. B133
Kahn, R. A72,B105
Kailin, E.W. A58,B105,B106
Kalita, D.K. B160
Kalman, C.M. B55
Kalman, T.P. B55
Kalinins, R. A6,B106
Kalymun, M. A30,B106
Kaminoff, R.D. A66,B107
Kaplan, G.A. B80
Karp, S.A. B232
Kasi, S. B105
Kasuga, H. A26,B107
Kasuya, M. B108
Kato, T. A22,B108
Kawano, S. B108
Kay, D.W.K. B87
Kaye, H. A82,B108
Keating, J.P. A64,B108
Keller, M.D. B108
Kelley, M.D. A5
Kennedy, C.B. B151
Kennedy, D.A. A21,B109
Kerigan, A.T. B158
Kerr, H.D. B109
Kerrebijn, K.F. A52,B109
Kershner, B. B110
Kershner, J. A8,B110
Kershner, J.R. A63,B110
Keskinen, H. B224
Keveova, E. B159
Khaleeluddin, K. B160
Kielblock, A.J. B224
Kilian, D.J. B240
Kim, Y.S. B110
King, D.S. A55,A56,B111
King, E. A52,B111
Kinloch, D. B31
Kinney, J.M. A31,B111
Kinsman, G. A47,B112
Kirkbride, J. B42,B129
Kirsh, S. A43,A69,A70,B112
Kittler, F.J. B113
Kjaer, G. B92
Klaucke, D.N. B225
Kleiman, C. A11,B113
Kling, H. B11
Knisely, J.S. B172
Kobayashi, D.H. B206
Komaroff, A.L. B20
Kon, S.H. A56,B113
Konopinski, V.J. A15,B113
Korosec-Serfaty, P. A7,B113
Korsgaard, J. B124
Kose, S. A4,B114
Kozuka, H. B108
Kral, V. B159
Krebs, H. B16
Kreiner, C. A35,B114
Kreiss, K. A19,B115
Krieger, M.H. B115

- Krietsch, K. B39
 Kruss, P. A58,B115
 Kruss, P. A22,B116
 Kuiper-Goodman, T. B215
 Kulle, T.J. B109
 Kume, Y. B143
 Kurumatani, N. B143
 Kusiak, R. B132
 Labelle, H. A78,B116
 Lechtman, D.S. B117
 Lanese, R.R. B108
 Larsen, L. B76
 Larson, G.E. B178
 Larue, D. B225
 Laseter, J.L. A82,B117,B170
 Lessiter, D.V. A14,B117
 Last, J.M. A84,B118
 Laties, V.G. A76,B118
 Lauer, L.D. B32
 Laufer, R.S. B118
 Laurie, R.D. A14,B118
 Laverne, A.A. A52,B119
 Layton, J. A67,B119
 LeBoutillier, J.C. B159
 Leach, S.J. B4
 Leaderer, B.P. B29,B234
 Lebowitz, M.D. A51,A53,B3,B120
 Lecuyer, G. A84,A85,B120
 Lee, A.J. A30,B120
 Lee, B. B208
 Lester, M.L. B213,B214
 Letourneau, E.G. A54,B121,B132
 Letz, R. B121
 Levi, L. A65,A68,B121
 Levin, A.S. B47
 Levin, L.I. B233
 Levine, S. B40
 Leviton, A. B145
 Lewis, A.L. B74
 Lewis, B.E. A42,B122
 Lewis, M. A33,B122
 Leyman, E. B232
 Lidwell, O.M. A9,B123
 Lillis, R. B223
 Lin, C.I. B91
 Lin, N. B52
 Lindstrom, K. A13,B123
 Lindvall, T. B3,B19
 Lipton, M. B40
 Liss, G.M. A13,B123
 Lofgren, I. B123
 Logsdon, R.F. B21
 Lopez de Victoria, A. B203
 Lord, J. B95
 Lorimer, J. B198
 Lowrey, A.H. B173,B174,B175
 Lubun, M. B66
 Lugg, D.J. B232
 Lund, S.M. B124
 Lundqvist, G.R. A19,B5,B124
 Lykke, A.W.J. B124
 MacLeod, K.E. A54,B124
 MacLennan, J.G. A8,B125
 Macpherson, A.S. A21,B125,B126
 Maeda, K. B147
 Mahaffey, K.R. A40,B126
 Maher, C. B145
 Makower, J. A15,A16,B126,B127
 Male, J.C. B28
 Mallov, J.S. A13,B127
 Mandell, M. B127
 Manga, P. B8
 Mansfield, J.R. B49
 Mao, Y. B231
 Marchant, R. A6,A74,B127,B128
 Morgulis, S.T. A42,B128
 Marina, K. A62,B128
 Marino, A.A. B16
 Markham, J.W. B129
 Marks, J.G. A16,B129
 Marshall, J. B44
 Martinez-Catanchi, F.L. B38
 Massari, J.P. B153
 Matas, M. B129
 Matsuki, H. B107
 Maulfair, C.G. A56,B130
 May, C.D. B130
 Mayhew, D.R. A25,B130
 Mayron, L.W. B154
 McAlaster, R. B213
 McCann, J.L. B36
 McCarthy, S. B42
 McCauley, G. A24,B130
 McClain, K.E. B194
 McCluskey-Fawcett, K.A. A24,B131
 McCullough, R.S. B132
 McDonald, E.W. B38
 McDonald, J.C. B131
 McDowall, M.E. A61,B132
 McGee, C.T. A61,B132
 McGrath, T. B132
 McGregor, R.G. A6,B132
 McIlhenny, M.L. B109
 McIntyre, D. B206
 McIntyre, E.D. B206
 McKenna, T.A. B18
 McKillop Farlow, D. A68,B133
 McKinnon, D.L. B18
 McKnight, J.L. A44,B133
 McLaughlin, D.E. B133
 McLean, A. B105
 McLeod, L. A33,B134
 McNall, P.E.J. B21
 McNeer, A.E. B111
 McNeil, T. B67
 McPhee, J. B4
 McQuiston, F.C. B21
 McVittie, D.J. A14,B134
 McWhinney, I.R. A78,B134
 Meek, N. B131
 Medical Services Division staff, W.C.B. A13,B135
 Medrich, E.A. B25
 Meggs, W. A66,B135
 Melia, R.J.W. A5,B135
 Mendelsohn, R.S. A33,B136
 Mental Health and the Workplace Committee A17
 Merkur'yeva, R.V. B23
 Mes, J. A57,B136
 Meyer, B. A82,B136
 Meyer, M.B. B86
 Michelson, L. B59
 Miksch, R.R. A15,A17,B92,B137,B220
 Milby, T.H. B117
 Millar, I.B. A20,B137
 Millar, W.J. A70,B137
 Miller, C.A. A70,B138
 Miller, C.S. A17,B138
 Miller, H.C. B67
 Miller, J.B. A56,B138
 Miller, J.D. B215
 Miller, S.M. A67,A81,B138
 Milling Pedersen, L. B9
 Mintz, S. B196
 Mitchell, C.L. B164
 Mitchell, E.W. B49
 Mitchell, R.I. B108
 Miyata, M. B99
 Moch-Sibony, A. A60,B139
 Mohr, R.D. A49,A80,B139
 Molhave, L. A54,A55,B5,B11,B139,B140

- Monahan, J.T. B140
Montanaro, A. B14
Morgan, M.S. B30
Morton, L.L. B110
Moschandreas, D. A6,A53,B141
Moss, R.W. B167
Motomiya, K. B242
Mourmans, A.R.M. B109
Mueller Associates B141
Muittari, A. A55,B141
Muller, H.K. B232
Mumetz, H. B191
Murabayashi, H. B210
Murphy, G. B177
Murphy, M.R. B111
Murray, M.J. B52
Murray, T.K. B142
Mussen, I. B233
Mustard, F.J. A85,B142
NIOSH B46
Nagda, N.L. A82,B142
Nagira, T. A20,B143
Nakaza, M. B105
Namba, T. B99
Naoki, H. B114
Nathanson, C.A. B143
National Com. for the Protection of Human Subjects
A28,B144
National Program to Reduce Tobacco Use, C.P.I.C. A53,
B144
Natusch, D.F.S. B200
Nazaroff, W.W. B28,B92
McNall, P.E. A81,B145
Needleman, H.L. A25,B145
Nelson, S.R. B70
Nero, A.V. A54,B28,B146
Nethercott, J.R. A53,B146
Neuberger, M. B80
Neus, H. B146
New York Academy of Medicine B147
Newball, H.H. B147
Nicander-Bredberg, H. B19
Nieder, J. A42,B147
Nielsen, A. B96
Nielzel, M.T. A66,B147
Nitta, H. A5,B147
Norris-Baker, C. A30,B148
Norton, S.A. B88
Novick, M. A71,B148
Nussbaum, K. A15,B149
O'Banion, D.R. A45,A56,A76,B149,B150
O'Brien, I.M. A14,B151
O'Connor, D. A66
O'Hanlon, T. B233
O'Hara, B. A71,B151
O'Quinn, S.E. A55,B151
Ofshe, R.J. B68
Ogihara, A. B210
Okuwaki, K. B99
Olafsson, O. B151
Olsen, J.H. A19,B152
Olson, H.G. B46
Ontario Federation of Labour A15,B152
Ontario Hydro A62
Ontario Mental Health Foundation B152
Ontario Ministry of Energy B153
Ontario Ministry of Health A61,B153
Orehek, J. A41,B153
Osaka, F. B107
Osborne, W.L. B95
Ostapovich, I.K. A84,B154
Ott, J. B154
Ott, J.N. A10
Ott, J.R. A61
Ott, W.R. B174
Ott, W. A16,B155
Otten, H. B146
Oxley, D. B230
Ozkaynak, H. A5,A52,B155
Page, S. B51
Paolini, R. B70,B132
Pape, B. A44,B156
Paroski, P.A. A46,A49,B156
Parris Stephens, M.A. B111
Pasternack, B.S. B77
Pauls, J. A4,B157
Paulson, G.W. B157
Pearlin, L. B55
Pearlman, M.E. B194
Pearse, I.H. B157
Pedersen, N.B. A15,B157
Pedersen, O.F. B11,B140
Peek, L.A. B150
Pelech, L. B214
Pelmeier, P.L. B129
Pengelly, L.D. B158
Pepelko, W.E. A14,B158
Pepper, J.H. B19
Pepys, J. A13,B159
Peresie, H. B145
Peterkin, A. B129
Petit, T.L. A25,B159
Pfeiffer, J. A25,B159
Pfeiffer, C.C. B160
Phalen, R.F. A52,B160
Philpott, K. B160
Philpott, W.H. A43,A76,B160
Pim, L.R. A57,B161
Planek, B. A4
Planek, T.W. B761
Plumlee, L. A41,A84,B161
Poole, R. A52,B161
Porter, A.L. B16
Poyner, B. A4,B161
Pranti, F.A. B132
Pressman, N.E.P. A21,B162
Preuss, P.W. B79,B221
Priesnitz, W. A10,A28,B162
Proctor, D.F. B5
Proctor, N.H. A13,B162
Proshansky, H.M. A65,A66,B107,B163
Prudham, D. A29,B164
Pryor, G.T. A83,B164
Pryor, P. A9,B164
Purchase, E.C.R. B240
Raab, K.H. A7,B165
Rainsbury, R. B16
Raizenne, M.E. B70
Raijoff, J. A26,B165
Ramsey, N.W. B49
Rand, G. A16,A84,B165,B166
Randolph, T.G. A5,A6,A8,B166,B167
Rapoport, A. A7,B167
Rapp, D.J. A8,A26,B167
Ray, S.M. A22,B168
Raymer, W.J. A57,B168
Rea, W.J. A44,A55,A56,A62,A81,A83,B117,B168,B169,
B170,B171,B203
Rector, H.E. B141,B142
Reed, R. B145
Rees, D.C. A9,B172
Reible, D.D. A53,B172
Reid, L.M. A40,B172
Reinhardt, C.F. A17,B172
Reischi, P. B160
Remafedi, G.J. A46,A47,B173
Reno, S.J. B164
Repace, J.L. A16,A53,A54,A81,B173,B174,B175,B176
Repetti, R.L. B176
Resin, T.S. A17,B176

- Richter, J. B159
 Richwood, D. B87
 Rieber, E.R. A12,B177
 Rieker, P.P. B40
 Riesenber, D.E. A74,B177
 Riley, E.C. A9,B177
 Riley, M. B55,B66
 Riley, R.L. B177,B178
 Rimland, B. A76,B178
 Ring, J.W. B181
 Rittfeldt, L. A55,B178
 Rivlin, L. B26
 Roark, S.F. B22
 Robbins, A.F. B179
 Robertson, L.S. A50,B179
 Robinson, A.M. A28,B179
 Robinson, J.D. B106
 Rogers, S.A. B180
 Root, D.E. A82,B180
 Rose, R.M. B55
 Rosenblum, I. B26
 Rosenfeld, A.H. B28
 Rosner, B. B212
 Roy, A.W. A38,B180
 Ruddel, H. B146
 Rudolph, J.P. B24
 Rudy, J.V. B220
 Russell, J.F. B25
 Russell, P. B198
 Ryan, J.P. B211
 Ryan, P.B. B155
 Sachs, H.M. A6,B181
 Sackner, M.A. B70
 Saegert, S. A34,A65,A68,B181,B182,B183
 Saito, H. B242
 Salvaggio, J.E. A14,B184
 Samadi, N. B171
 Sandberg, M. B178
 Sandia National Labs A81,B184
 Sarason, I.G. A66,B184
 Saric, M. A20,B184
 Sass, R. A15,B185
 Savage, H. B95
 Savitz, D.A. A61,B185,B186
 Scadding, G.K. B186
 Schachter, E.N. B234
 Schaumburg, H. A55,B186
 Schauss, A. A25,A52,B187
 Scheel, L.D. B208
 Scheere, A.R. B115
 Scheff, P.A. B225,B226
 Schiager, K.J. B46
 Schiefer, H.B. A77,B187
 Schlesinger, R.B. B187
 Schmidt, H.E. B137
 Schnakenberg, G.H.J. B49
 Schnare, D.W. A75,B188
 Schnarr, A. B95
 Schorr, L.B. B138
 Schrecker, T.F. A81,B188
 Schreiter, A. A11,B188,B189
 Schroeder, H.A. A58,B189
 Schulte, J.H. B118
 Schwartz, H.J. B38
 Schwartz, P.M. B64
 Scott, P.M. A54,A57,B189
 Scott, R. B87
 Sears, A.G. B201
 Seedorff, L. B6
 Seguin, M. A42,B189
 Seifert, B. B190
 Selikoff, I.J. A13,B190,B223
 Selner, J.C. A44,A75,B190
 Selye, H. A64,B190
 Semenenko, A.D. B23
 Seppalainen, A.M. B191
 Serio, A. B15
 Severs, R.K. A41,B191
 Sexton, K. B202
 Seyal, A.R. A55,B191
 Shah, C.P. A38,A39,B191
 Shain, M. A15,B192
 Shair, F.H. B172
 Shakman, R.A. A40,B192
 Sharp, D.S. A58,B192
 Shaw, F.H. B72
 Shearer, R.W. A58,B50,B193
 Sheehy, N.P. A50,B193
 Sheppard, H. B179
 Sherman, M.H. B79
 Shields, M.G. B188
 Shifrin, L. A36,B194
 Shirliff, C.J. B25,B38
 Shishido, M. B210
 Shore, R.E. B77
 Shy, C.M. B194
 Siddique, C.M. B46
 Silberstein, S. A6,B194
 Silver, F. B195
 Silver, R.L. B195
 Silverman, F. A40,B196
 Simard, M. A18,B196
 Siminovitch, L. B90
 Simonsson, B.G. A51,B196
 Simpson, H.M. B130
 Simpson, M. B197
 Sivak, M. B197
 Sizto, R. B15
 Skov, A. B6
 Skov, P. B76
 Skrzycki, C. A17,B197
 Small, B.M. A4,A5,A6,A7,A8,A11,A17,A29,A40,A76,A81,
 A83,B197,B198,B199
 Smiley, R.E. B168,B203
 Smith, A.H. B192
 Smith, C. B199
 Smith, C.W. A62,B200
 Smith, L. A66,A68,B200
 Smith, R. B13
 Solomon, R.L. A20,B200
 Somers, E. B201
 Sorensen, L. B26
 Soskolene, C.L. A47,B201
 Spasoff, R. A70,B201
 Spedding, D.J. B202
 Speizer, F.E. A5,B65,B202,B212
 Spencer, C. A27,B202
 Spencer, P.S. B186
 Spengler, J.D. A53,B42,B54,B65,B110,B121,B155,B202,
 B218
 Spielberger, C.D. B184
 Spitzer, W.O. A83,B203
 Sprague, D.E. A6,B203
 Spyker, J.M. B229
 Squires, B.P. A78,B204
 St. Lawrence, I. A30,B204
 Stagner, B. B39
 Stansfield, M. B110
 Stapleton, P.J. B62
 Steadwell, L. B25
 Stebbings, J.H.J. A41,B204
 Steensberg, J. B205
 Stein, M. B40
 Stephens, R. A24,A25,B205
 Sterling, E. B206
 Sterling, E.M. A16,A17,B206,B207,B219
 Sterling, T. B206
 Sterling, T.D. A16,B206,B207
 Stewart, B.W. B124
 Stewart, D.W. A44,B207

- Stewart, R.D. A41,B207,B208
Stidi, H. B80
Stokes, P.M. A39,B208
Stokholm, J. B92
Stokinger, H.E. B208
Stotsky, K. A32,B208
Stratton, P. A50,B209
Strickland, B.R. A76,B209
Strydom, N.B. B224
Stuudenmayer, H. B190
Subcommittee on Formaldehyde A9,B209
Suedfeld, F. A64,B210
Suess, M.J. A19,B210
Sugita, K. A52,B210
Suits, C.W. B168
Sullivan, J.L. B211
Sult, S. B66
Sun, W.F. B136
Sundell, L. B83
Suryanarayana, S. B100
Sutton, R. A42,B211
Svensson, P.G. B151
Swank, M.G. B69
Swidersky, P. B109
Swift, D.L. B5
Symonds, A. A33,B211
Szymusiak, S.M. A4,B211
Tager, I.B. A26,B212
Tall, F.D. A11,B212
Taniguchi, H. B132
Task Force on the Child as Citizen B212
Taylor, R.B. A66,B213
Thatcher, R.W. A25,A82,B213,B214
Thielebeule, U. A52,B214
Tichauer, E.R. B235
Tilson, H.A. B164
Tobin, R.S. A5,A54,B214,B215
Tockman, M.S. B86
Todd, D. A57,B215
Toews, J. A44,B216
Toews, J.A. B207
Toplack, S.A. B158
Toronto Star A11,A22,A27,A70,A79,B216,B217,B218
Tosteson, T.D. A52,B218
Townsend, J.C. B69
Townson, M. A34,B218
Toyama, T. B104,B105
Trainor, J. A44,B219
Trautlein, J.J. B129
Traynor, G.W. B73,B91,B92,B219
Trief, N.M. B168
Tsuchiya, Y. B227
Tsunoda, T. B105
Tsuru, K. B105
Tucker, W.F. B203
Turiel, L. A5,B91,B220
Turk, A. B21
Turner, G. A27,B220
Turner, R. B24
Turner, W.A. B155
Turton, D. B136
Twigge-Molecey, C. A81,A84,B221
US National Research Council A20,B221
US Radiation Policy Council B221
Ueno, M. B143
Ulsamer, A.G. A53,B16,B79,B221
United Nations A22,B222
University of Michigan A61,B222
Unknown A14,B222,B223
Uno, H. B114
Uyeno, E.T. B164
Valbjorn, O. B96
Valciukas, J.A. A76,B223
Valciote, L.M. B115,B116
Valsiner, J. B70
Van Assendelft, A. A6,B224
Van Der Linde, A. B224
Van Der Wall, W.H. B224
Van Rensburg, J.P. A20,B224
Vanderslice, S.F. B21
Vanderveen, J.E. B126
Vasudev, P. B132
Vaux, A. B140
Vayda, E. A74,B224
Vecchia, P. B15
Veneskoski, T. B141
Victor, J. B200
Vigoda, D.S. B89
Vilcins, G. B74
Visual Arts Ontario A14,B225
Vogl, R.L. B225
Voronova, B.Z. A84,B225
Wadden, R.A. A14,A82,B225,B226
Wales, R. A6,B226
Walker, P. B26
Walkinshaw, D.S. B227
Wallace, L.A. B174
Waller Scanlon, L. B127
Waller, J.A. A4,B227
Walsh, P.J. B57
Wanner, A. B70
Ward, E.C. B52
Ware, J.H. B121
Wartew, G.A. A53,B228
Washington, L.S.J. B32
Waterman, F.K. A24,B228
Webber, G.M.B. A4,B228
Weber-Tschopp, A. B228
Webster, X. B135
Weicker, L.P.J. B228
Weiffenback, C.V. B88
Weinstein, M.S. A21,B229
Weiss, B. A25,A76,B229
Weiss, S.T. B212
Weker, R.A. B218
Welch, P. B106
Werner, C.M. A7,B4,B230
Wessendarp, T. B118
Whiston Spinn, A. A22,B230
White, B. B39
White, F.A. A60,B230
White, R.P. B70
Whitfield, J. B56
Wigle, D.T. A19,A55,B42,B137,B231
Wilde, V.L. A30,B106,B232
Williams, D.L. B232
Williams, J.S.J. B232
Williamson, C. B62
Williamson, J.W. B52
Wilson, J. B87
Wilson, P.T. B232
Wilson, R. B233
Wilson, R.W. B24
Wilson, S. B101
Winkel, G. B182
Winkel, G.H. A39,A75,B233
Winkelstein, W.J. B233
Witek, T.J. A40,B234
Witherell, L. B225
Wnesley, B. B127
Wolfe, D.A. B101
Wolfe, M. A27,A43,A44,A84,B26,B27,B118,B234,B235
Wolkenberg, R.C. A15,B235
Wolochow, H. B54
Wood, R.W. A15,B235
Woods, J.E. B21
World Commission on Environment and Development A81,
A85,B236
World Health Organization A29,A44,A55,A59,A78,A85,
B236,B237,B238,B239

Wortman, C.B. B195
Wotton, E. A10,B239
Wright, G.R. B132
Wright, M. A17,B240
Yamamoto, M. B143
Yanagisawa, Y. B110
Yang, W.H. A57,B240
Yassi, A. A75,B241
Ydreborg, B. B83
Yocom, J.E. A52,B241
Yonts, P. B172
Yoshida, K. B127,B128,B242
Yoshida, R. B242
Youdim, S. B171
Young, G.S. A5,B242
Young, M.M. B194
Young, R.A. B19,B220
Yulsman, T. A50,B242
Zak, L. B101
Zamm, A.V. B243
Zarbatany, L. B71
Zielhuis, R.L. B63
Zilstorff, K. B9
Zube, E.H. B243
Zuckerman, B.S. B243
Zuckerman, D.M. A26,B243
Zweideinger, R.A. B243
Zwillich, C.W. B129

- abatement B46,B106,B127,B146
 abatement of noise B231
 abatement procedures B125,B165
 abatement programs B70
 abatement, dust B46
 abdominal pain A57,B240
 abdominal reflexes A9
 aberration, behavioral B103
 abilities, spatial A35,B84
 ability to concentrate, impaired B227
 ability to move B167
 ability, learning A8
 ability, mental A80
 ability, motor B198
 able-bodied B58
 able-bodied adults A50
 able-bodied people A4
 able-bodied, temporarily A40,A41
 able-bodied adult B202
 abnormal electroencephalograms B160
 abnormal electromagnetic environment B16
 abnormalities, immune B171
 aboriginal B181
 aboriginal people A39
 about women, information B35
 about youth, false ideas A27
 abrupt disruption B18
 absence of disease B72
 absence, leaves of A18
 absence, policies for leaves of B93
 absenteeism A63,B77
 absorption A83
 abuse by adults A25
 abuse victims A28
 abuse, alcohol 12,A46,A66,B61,B71,B152,B173
 abuse, child 12,A25,A28,A66,B34,B71,B112,B179
 abuse, drug A46,B173
 abuse, emotional A26,A33,B13,B34,B173
 abuse, illicit drug B192
 abuse, physical B34,B134
 abuse, sexual B13,B34,B134
 abuse, substance 13,A78,B108,B134,B135,B192
 abusive adult behaviour A28
 academic buildings B212
 academic development B150
 academic health classes B76
 academic offices A11,B212
 acceleration A63,B100
 acceleration stress B100
 acceptable care, culturally A39,B7
 acceptance A27,A49
 access buildings, public B73
 access to a window B240
 access to children B86
 access to decent jobs A71
 access to information B144
 access to services B86
 access to services, equitable B8
 access, equal B213
 access, residential B211
 access-oriented problems B211
 accessibility A42
 accessibility, building B128
 accessible 3,A68
 accident in the home A4
 accident incidence on stairs B114
 accident liability B193
 accident prevention B70,B179,B211
 accident statistics B36
 accident, nuclear A67,B61
 accident, traffic B4
 accidental death, residential B4
 accidental injuries B70,B114
 accidental spills B74
 accidents A50,B36,B38,B52,B75,B83,B86,B114,B161,B164,
 B193,B197,B228,B229
 accidents in the workplace B192
 accidents, building related B114
 accidents, childhood B70
 accidents, children's B193
 accidents, domestic B114
 accidents, falling A29,B157,B197,B211,B227
 accidents, home B37,B161,B228
 accidents, household A4
 accidents, nuclear A68
 accidents, office A16,B35
 accidents, risk of B198
 accidents, stairway A20,B10
 accidents, wheelchair B59
 accommodation A46,B2
 accommodation, scarce B63
 accountability B63
 acoustic environment B231
 acetate, methyl cellulose A14,B226
 acetone A14,B23,B226
 achievement, school B213
 achievement, student B84
 achieving health for all B133,B134,B204
 achieving health for all: implications B142
 acid depositions B70
 acid factory, sulphuric A38,B180,B208
 acid precipitation A52,B33,B156
 acid rain A22,A56,B70,B116,B155,B216,B217
 acid, formic B162
 acid, isocyanic B53
 acid, mandelic B191
 acid, sulfuric A8,B69
 acidic aerosols A52,B156
 acidic precipitation A22
 acidic sulfur B187
 acidified lake water A22,A58,B70
 acids A14
 acids, ambient B188
 acids, inhaled B187
 acknowledging difference B26
 acid B199
 acoustic environment B230
 acoustic privacy A60,B230,B231
 acoustic sensibilities B231
 acquired immune deficiency syndrome A47
 acrodynia B89
 acrolein A14
 acrolein vapor B32
 acrylic resins B67
 acrylonitrile B67
 act, child welfare B179
 act, environmental protection B94
 acting through consent, people B133
 action, collaborative B34
 action, environmental B9
 action, group social B213
 action, intersectoral B239
 action, local B83
 action, plan of B123
 action, responsible B162
 action, social B116
 action, support for citizen B144
 actions, environmental B9
 actions, public B139
 actions, sexist A35
 actions, tort B79
 activated carbon B226
 activation B150
 active B183
 active creators B235
 active, socially B204
 activities, community A43
 activities, daily B1
 activities, group B234

- activities, school B243
activities, suitable B84
activity systems B182
activity, brain's electrical A82,B108
activity, educational B222
activity, industrial A51
activity, physical B76,B80
activity, sedentary B17
activity, social B36
activity, spontaneous locomotor B118
acts, antisocial B184
acuity, visual B7
acute B157
acute care A75,B56
acute care, patients in B233
acute health effects B156
acute myelogenous leukemia B186
acute myeloid leukemia B132
acute respiratory conditions B24
acyclic alcohols B36
adaptability 8,A56
adaptable A50
adaptation B16,B107,B111,B112,B149
adaptation to illness B147
adaptation, enzyme B2
adaptation, specific B53
adaptations, psychological B147
adapting the home B122
adaption 6
adaptive A68
adaptive environment B1
adaptive environments B1
adaptive mechanisms B2,B18
add-on exhaust hardware B49
addiction A19,B231
addictive substances B112
addicts, heroin A19,B231
additive B43
additives A55,B14,B53,B127,B161
additives, chemical B47,B210
additives, food A43,A56,A57,A76,B46,B48,B60,B73,B88,
B113,B136,B138,B187,B190,B200,B229,B240
additives, gasoline B116
adequate health care B192
adhesives A13,B6,B14
adipose tissue B188
administration, food and drug B168
administrators B50,B235
administrators, government B185
admission rates, higher B152
admission rates, hospital A69
admission rates, mental hospital B112
admission requirements A38,B44
admission, hospital B15,B192
admissions, hospital 12,A38
admissions, mental health 13
adolescent B173
adolescent crime B112
adolescent homosexuality B173
adolescent, homosexual A46
adolescents B65,B147,B151
adolescents, health disturbances in B151
adolescents, homosexual A49,B156
adsorption, mercury B202
adult B34
adult behaviour, abusive A28
adult population B149
adult support A25
adult values A66,B74
adult, able-bodied B202
adulthood 12,17,B12
adults A27,B74,B77,B144,B193
adults in miniature B213
adults, able-bodied A50
adults, abuse by A25
adults, dependence on A28,B144
adults, education of B22
adults, mentally handicapped B153
adults, needs of B25
adults, older B106
adults, profound distrust of B74
adults, young B147
adults, young unskilled B151
adults, younger B87
adverse drug reactions B86
adverse reaction B169
adverse reactions, sulfite-related B240
advertisements B142
advisory committee, federal/provincial A5
advocacy stance B216
advocacy, mental health A44,B39
aero-allergic reaction B187
aeroallergens A53
aerobic exercise B180
aerosol hair spray A5,B70
aerosol propellants B27
aerosol science B242
aerosol sprays A32,A74,B22,B169,B209
aerosol, infectious B32
aerosols B54,B57,B142,B160,B200,B211
aerosols, acidic A52,B156
aerosols, microbial B82
aesthetic qualities B162
affection in public spaces A49
affection, withholding of A26
affection, withholding of B34
affectional attractions A48
affectional ties A72,B87
affective responses B55
affective shifts B108
affiliation B12
affiliation, need for B12
affinity groups B182
affluent B233
affordability A30,B32
affordable housing B78,B232
aflatoxin B161
aflatoxins A57,B189
against children, prejudice B50
against homosexuality, overt prejudice A49
against homosexuality, prejudices B2
against lesbians and gay men, discrimination B41
against women, discrimination B26,B97
age 3,A2,A23,A68,A80,A84,B1,B26,B112,B131,B157
age discrimination B25,B112
age factors B164
age groups, older B130
age segregation A10,B162
age, bone B214
age, early B86
age, middle B22
age, nuclear B74
age, old B86
age, preschool B52
age-integrated A30,B232
age-related processes B86
age-specific B34
aged B38,B75,B209
aged cerebrovascular disorders B164
ageism 17,B25
ageism, roots of B25
agencies, regulatory B185,B201,B206
agent, noxious B157
agent, sensitizing B228
agents, antistatic A55,B14
agents, chemical A75,B75,B241
agents, environmental B223
agents, etiologic B74
agents, etiological B104
agents, food triggering B171

- agents, immunosuppressive B103
 agents, infectious A24,B75
 agents, physical B75,B241
 agents, sizing A55,B14
 agents, wetting A55,B14
 aggregate co-worker ratings B176
 aggression A27,B216,B243
 aggression, extreme B151
 aggressive A35,B99,B183
 aggressive behaviour A26
 aggressive behaviour, children's B243
 aggressiveness A25,B229
 aging B2,B56,B75,B85,B86,B89,B106,B120,B148,B229
 aging gay people A30
 aging population A74,B224
 aging, attitudes towards B90
 aging, biology of B2
 aging, fear of 13
 aging, gays and lesbians B81
 aging, Ontario longitudinal study of B89
 aging, study on B89
 agreements, collective bargaining B185
 agricultural A51
 agricultural chemicals B27,B60
 agricultural commodities A57,B189
 agriculture B239
 aid B184
 AIDS A47,B2,B4,B55,B201
 AIDS crisis B41
 AIDS, mortality from 13
 aids societies, children's B179
 ailments, serious physical B47
 air changes B227
 air cleaners, electronic B226
 air cleaning B146,B175
 air cleaning devices B207
 air conditioning A54,B54,B181
 air conditioning ductwork B21
 air conditioning equipment B28
 air distribution B18
 air exchange B194
 air filters B21,B113
 air heating system, forced B165
 air indoor pollution B86
 air infiltration B91
 air intakes, school A11
 air ions B123
 air ions, negative A8,A43,B110
 air movement B123,B131
 air pollutants, inhaled B119
 air pollution B5,B15,B18,B19,B23,B25,B28,B36,B38,B41,
 B43,B46,B49,B50,B52,B53,B60,B61,B63,B67,B79,B81,B87,
 B88,B89,B90,B91,B92,B93,B97,B100,B104,B105,B107,B108,
 B109,B111,B115,B116,B117,B118,B119,B124,B133,B137,
 B141,B143,B145,B152,B158,B160,B172,B184,B187,B189,
 B190,B191,B192,B194,B196,B197,B200,B203,B204,B210,
 B213,B214,B223,B224,B229,B237,B241,B242
 air pollution control B184,B223
 air pollution detection B91
 air pollution sampler, domestic B93
 air pollution, domestic B93
 air pollution, indoor 5,B5,B6,B11,B13,B14,B15,B16,
 B17,B19,B21,B22,B27,B28,B29,B35,B38,B42,B43,B46,B47,
 B48,B53,B54,B55,B57,B62,B65,B67,B69,B73,B74,B78,B79,
 B82,B88,B89,B90,B91,B92,B97,B101,B108,B109,B113,B115,
 B120,B123,B124,B126,B127,B128,B135,B136,B137,B139,
 B140,B141,B142,B145,B147,B149,B150,B152,B155,B158,
 B164,B165,B166,B167,B173,B174,B175,B176,B177,B178,
 B181,B184,B186,B188,B189,B190,B194,B196,B197,B198,
 B202,B203,B206,B207,B211,B212,B217,B219,B221,B222,
 B225,B226,B231,B234,B236,B242
 air pollution, outdoor B158,B166,B174
 air pollution, particulate A52,B155
 air pollution, photochemical B104
 air quality B19,B91
 air quality complaints B212
 air quality management B136
 air quality standards B140,B175
 air quality, indoor A4,B3,B6,B18,B21,B51,B56,B66,B73,
 B76,B79,B92,B96,B103,B104,B110,B123,B125,B131,B146,
 B153,B159,B165,B166,B172,B180,B184,B187,B189,B197,
 B199,B201,B205,B206,B207,B209,B214,B215,B220,B226,
 B227,B236,B237,B238
 air sampling devices B184
 air standards, clean B30
 air supply B225
 air supply, fresh B18
 air travel, smoke-free B217
 air, bacteria in indoor B214
 air, classroom B110
 air, clean A51
 air, fresh A11,B158,B172,B212
 air, indoor A53,B215
 air, make-up B21
 air, makeup B175
 air, of indoor B19
 air, outdoor 5
 air, poisoning of B90
 air, polluted indoor 3
 air, pollution-free B119
 air, recirculation of return B19
 air, recycled B165
 air, school B56
 air, viruses in B104
 air-purification devices B220
 air-tight buildings B206
 airborne bacteria B214
 airborne bacteria, health effects of B214
 airborne contaminants, reducing B145
 airborne infection B178
 airborne microorganisms B82
 airborne organisms B177
 airborne pollutants, longrange transport of B70
 airborne radon B88
 airborne route B104
 airborne viruses, indoor B238
 aircleaning systems B210
 aircraft B210
 aircraft noise A60,B77,B139
 airflora B215
 airline flights B217
 airports B77
 airtight B226
 airtight buildings B124
 airtight homes B27
 airtight office buildings B220
 airway hyperresponsiveness B234
 airway irritation, upper B5
 airway obstruction, upper A16
 airway reactivity B38
 airway resistance B5,B6,B153
 airways, upper B6
 alcohol A25,A30,A70,A74,B46,B63,B112,B116,B130,B135,
 B137,B157,B169,B192,B204,B235
 alcohol abuse 12,A46,A66,B61,B71,B152,B173
 alcohol consumption A76,B76,B80
 alcohol consumption, excessive B137
 alcohol intake A15
 alcohol use B61,B76,B130
 alcohol, ethyl B169
 alcohol, methyl A9,B164
 alcoholics A19,B231
 alcoholism B192
 alcohols, acyclic B36
 aldehydes A5,B43,B46,B91
 aldehydes in offices B137
 alert A66
 alienated A72
 alienated individuals B68
 alienation A67,B9

-
- aliens B216
 - aliveness B57
 - alkalis A14
 - all, achieving health for B133,B134,B204
 - all, health for A2,B83
 - all: implications, achieving health for B142
 - allergenic substances B29
 - allergens A26,B211,B220
 - allergens in school buildings B113
 - allergens, environmental B124
 - allergens, isolation from environmental B124
 - allergens, natural B3
 - allergic A53,B48,B80,B159,B173,B239
 - allergic alveolitis B224
 - allergic contact dermatitis A16,B129
 - allergic diseases B170
 - allergic disorders, occupational respiratory B159
 - allergic disorders, respiratory A13
 - allergic population B198
 - allergic pseudo-retardation B127
 - allergic reactions 12,B3,B16,B154
 - allergic responses B51,B90,B215
 - allergic rhinitis B186
 - allergic sensitization B97
 - allergic skin reactions A55
 - allergic students B113
 - allergies A11,A20,B67,B73,B113,B143,B147,B167,B186,
B187,B196,B200
 - allergies, childhood B167
 - allergies, food A76,B127,B150,B160,B167,B178,B187
 - allergies, skin B215
 - allergy A9,A45,B31,B48,B76,B96,B113,B160,B172,B190,
B197,B203
 - allergy tension fatigue syndrome B160
 - allergy testing, sublingual B111
 - allergy, causation of B239
 - allergy, cerebral B127,B160
 - allergy, food B171
 - allergy, unrecognized B167
 - alliance, environmental health B161
 - allies A28
 - allocation of time B44
 - alone, working B28
 - alteration, immune B52
 - alteration, subtle behavioral B54
 - alternaria B189
 - alternating electric fields B102
 - alternative environments A83
 - alternative lifestyles 16
 - alternative social environments B157
 - alternative sources of energy B237
 - alternative, work B37
 - alternatives to health care B85
 - alternatives, housing B42
 - alternatives, income 17
 - alternatives, public policy B203
 - aluminum A52,A56,B23,B218
 - alveolitis, allergic B224
 - alveolitis, fibrosing B124
 - alzheimer's disease B5
 - amalgam filling, dental B94
 - ambient acids B188
 - ambient stressors A85,B31
 - ambiguity, role B44
 - ambulatory care B22
 - ambulatory electrocardiograph recordings,
programmable B191
 - amenities B213
 - american cities B109
 - amitrole B193
 - ammonia A8,A15,B23,B69,B127,B162,B211
 - amoeba proteus B224
 - among people, variability A24
 - among women, illness B143
 - among women, poverty 12
 - amplification B214
 - amplifier, unstable B200
 - amplitude B28
 - an environment issue, diversity as B26
 - anaesthetic gases A32,B209
 - anal intercourse A48
 - analgesia B131
 - analysis B5,B136,B164
 - analysis and evaluation B56
 - analysis, chemical B91
 - analysis, computerized brainwave 10
 - analysis, gas B91
 - analysis, hair B213
 - analysis, quantitative B225,B226
 - analysis, regression B204
 - analytical techniques B136,B238
 - analyzing, infrared B7
 - anaphylactic shock A57,B240
 - anatomical classification B157
 - anemia B167
 - anemic A40,B5
 - energy B232
 - anesthesia B131
 - anger A69,B33,B55,B112
 - angina A53,A56,B5,B171
 - angina pectoris B10
 - angioedema A57,B240
 - anglo-Canadian B7
 - animal experiments B201
 - animal husbandry B239
 - animal life A22
 - animal studies B102,B133,B157,B186
 - animals, diseases in B189
 - animals, experimental B126
 - anishinabek A38,B180
 - annual job hour reductions B149
 - anonymous A75
 - antacids B23
 - antaretic B232
 - anti-gay discrimination B41
 - anti-gay scale 13
 - anti-semitism A46
 - anti-social behaviours B182
 - antiasthmatic medication B58
 - antibody response A83,B52
 - anticarcinogens A57,B4
 - antigens B51,B232
 - antisocial A8,B22
 - antisocial acts B184
 - antisocial behavior A66,B24,B71
 - antisocial behaviour B178
 - antistatic agents A55,B14
 - antistats A55
 - anxiety A33,A43,A64,B42,B50,B54,B86,B90,B101,B103,
B108,B112,B176,B184,B210
 - anxiety levels B54
 - apartment, senior citizens B106
 - apartments B111,B243
 - apartments, high density B183
 - apartments, senior citizens A6
 - apathy 12
 - apathy/despair B112
 - apertures B222
 - appear as normal, desire to B122
 - appliances, combustion B41
 - appliances, gas B73,B86,B219
 - appliances, gas cooking B108
 - appliances, gas-fired kitchen B74
 - appliances, home A62
 - appliances, indoor combustion B73,B219
 - appliances, unvented combustion A5
 - applicants, elderly B35
 - apprentice B202
 - approach, holistic B86
 - approach, individualized curative B213

- approaches, coordinated B222
 approaches, self-interested B34
 appropriate housing B95
 appropriate technology A67,B165
 architects A74
 architectural barriers A42,B122,B128
 architectural design B140
 area, poverty B80
 areas, nonpoverty B80
 areas, recreational B65
 areas, remote B216
 areas, residential B23
 areas, rural B148
 areas, sensitive B70
 areas, smoking B207
 areas, urban A21,B200,B221
 arenas B7
 arenas, ice A19
 arms race A66,B74
 arms race, psychological aspects of the B74
 arms, nuclear A35,B13
 aromatic hydrocarbon, polycyclic A14
 aromatic hydrocarbons B137
 aromatic hydrocarbons, polycyclic A83,B52,B117
 around, school children B41
 arrangements, living B86
 arrangements, work A68
 arrest, false A38,B64
 arrhythmia B38,B75,B180
 arrhythmia, cardiac B27,B169
 arrhythmias, cardiac A83,B171
 arsenic B116,B162,B193
 art A8,B226
 art and craft materials A5
 art materials B84,B225
 art paper A8,B69
 art supplies B47,B167
 art workshops A14
 arthritis, rheumatoid A76,B49,B50,B171
 artificial lighting A16,B69
 artificial ventilation B177
 artists B225
 arts studios A12
 asbestos A9,A15,A20,A53,A58,B6,B25,B43,B57,B100,B115,
 B116,B127,B136,B201,B203,B211
 asbestos, friable B26
 ash, fly B200
 ashamed A35,B114
 ashrae standards B226
 ashrae ventilation standard B227
 asians, southeast B67
 aspects of homes, temporal B230
 aspects of the arms race, psychological B74
 aspects, behavioral B71
 aspects, behavioural B133
 aspergillus B169,B222
 aspergillus fumigatus B224
 assault A46,B67
 assault, sexual A33,B34
 assembly lines, paced B44
 assembly, freedom of B139
 assertive B183
 assessment B209
 assessment of risk B200
 assessment processes, risk B237
 assessment units, geriatric B56
 assessment, environmental impact B141
 assessment, impact B16,B23,B221
 assessment, risk B21,B123,B168,B201
 assimilating A79
 assimilation B2
 assistance, medical B23
 assistance, welfare B63
 associates A72
 association, psychological B60
 associations, nurses' B116
 associations, structure of B133
 associations, symbolic B183
 asthma A9,A36,A42,A51,A75,A77,B29,B31,B38,B96,B107,
 B127,B130,B141,B147,B151,B152,B159,B169,B171,B172,
 B196,B204,B205,B242
 asthma, bronchial A13,B14,B143,B242
 asthma, occupational B82
 asthmatic A41,A45,B5,B58
 asthmatic attacks A57,B240
 asthmatic reactions A13
 asthmatic symptoms B51
 asthmatics A40,B171,B196,B234
 at risk, children B202
 at work, role B44
 at work, stress B44
 athletes B224
 atmospheric B60
 atmospheric emission of metals B241
 atmospheric pollutant B133
 atomic energy B90
 atomic war, fear of B190
 atrophy, cerebral B10
 attacks, asthmatic A57,B240
 attacks, coronary heart B44
 attendance laws, compulsory B162
 attendance, compulsory B162
 attention A65,B32
 attention span A25
 attention span, decreased B150
 attention span, short B167
 attention, deleted from public B58
 attention, lapses of A10
 attention, medical B52
 attentional troubles A60,B139
 attitude B216
 attitude, condescending A33
 attitude, societal B231
 attitudes A17,B12,B41,B233,B243
 attitudes towards aging B90
 attitudes, conflict of B37
 attitudes, favorable B35
 attitudes, homophobic A47
 attitudes, ingrained B50
 attitudes, negative A44
 attitudes, personal A79,A80
 attitudes, public B51
 attitudinal barriers B1
 attitudinal changes 17,A2
 attractions, affectional A48
 attribution therapy B141
 auditory memory B110
 auditory memory skills B150
 auditory stimulus A60
 auditory system A60,B230
 authority A27,B65,B235
 auto exhaust B97
 auto-immune vasculitides B171
 automation A68,B122
 automobile B100
 automobile brakes A20
 automobile emissions B116,B205
 automobile exhaust A8,A20,B33,B137,B143,B167,B199,
 B200,B224
 automobile interiors B243
 automobile traffic dust A22,B217
 automobiles B60
 automotive emissions B36
 automotive exhaust B83
 automotive shops A12
 autonomic nervous system B191
 autonomy A2,A28,A30,B92,B144,B148,B162,B196
 autonomy of the child B162
 autonomy, childhood B162
 availability of services B144

- availability, job B112
 aversion B55
 aversive event A67,B138
 avoid, pitfalls to B225
 avoidance B53,B95,B125,B170
 avoidance measures B18
 avoidance, mould B48
 awareness B8,B74
 awareness of isotrogenic disorders B157
 awareness, professional A28
 awareness, public B83
 awareness, safety B86
 babies B131
 baby A24
 bacilli, gram negative B210
 back pain, chronic low A65,B66
 backdrafting A6
 background, class B183
 background, cultural A38,A39,B86
 background, educational A65
 background, genetic B53
 background, professional B47
 backgrounds, cultural B26
 backpacking stove, gas B41
 backyard phenomenon, not in my B78
 backyard, not in my A7
 bacteria A6,A9,A29,B21,B54,B82,B123,B222,B238
 bacteria in indoor air B214
 bacteria, airborne B214
 bacteria, culturing of A54
 bacteria, health effects of airborne B214
 bacteria, legionella A54
 bacteria, spread of B123
 bacterial A54
 bacterial disease B214
 bacterial infection, enhanced susceptibility to B32
 bacterial infection, susceptibility to A14
 bacterioflora B210
 bacterium A58
 bad weather B21
 balance A4,A29
 ballasts, fluorescent light A54,B124
 bank B233
 banned A57
 banned pesticides B215
 bare floors B165
 bargaining agreements, collective B185
 barrier, radon B46
 barrier-free B1
 barriers A28,B90,B95
 barriers to health care B192
 barriers, architectural A42,B122,B128
 barriers, attitudinal B1
 bars, grab A29
 base on environmental health, data B237
 base, community's economic B37
 base, scientific B91
 baselines 13
 basements B141,B217
 basic design tools B1
 basic human needs B97
 basic human rights B222
 basic needs B97
 basic wellbeing B149
 basis, empirical B130
 bathroom B82
 bathroom ventilation B165
 battered woman B134
 battered women A33,A34,B101
 battered women, shelters for B35
 battering B220
 battering of women 12
 battering, wife B134
 beaten A33
 beatings B13
 beer A58,B240
 behaviour B15,B22,B23,B32,B37,B46,B59,B64,B73,B80,
 B84,B103,B107,B118,B119,B122,B145,B150,B159,B167,
 B187,B198,B199,B209,B223,B229,B235,B243
 behaviour change strategies B22
 behaviour disorders, childhood B71
 behaviour indicators 13
 behaviour modeled by peers, unhealthful B76
 behaviour problems B71,B101,B167
 behaviour problems during childhood B134
 behaviour sciences B59
 behaviour toxicology B73
 behaviour, abusive adult A28
 behaviour, aggressive A26
 behaviour, antisocial A66,B24,B71,B178
 behaviour, children's B243
 behaviour, children's aggressive B243
 behaviour, classroom B110,B162
 behaviour, coping B61
 behaviour, criminal B178
 behaviour, deterioration in A13
 behaviour, discriminatory 15,A80
 behaviour, health B76
 behaviour, healthful A73,B76
 behaviour, human B71,B113,B167,B213,B230
 behaviour, hyperactive A61
 behaviour, non-adaptive classroom B145
 behaviour, oppressive A80
 behaviour, patterns of B182
 behaviour, personal B8
 behaviour, preventive B59
 behaviour, regression to infantile B90
 behaviour, restrictions on B44
 behaviour, safety-related B192
 behaviour, sexist A35
 behaviour, social B12
 behaviour, stress-related B108
 behaviour, type a B22
 behaviour, type-a coronary-prone B50
 behavioural aberration B103
 behavioural alteration, subtle B64
 behavioural aspects B71,B133
 behavioural changes B17
 behavioural competencies B147
 behavioural constraint B181
 behavioural development B64
 behavioural differences B119
 behavioural effects B38,B59,B111,B149,B181,B198
 behavioural factors B22,B61
 behavioural influences B229
 behavioural interventions B22
 behavioural problems B187,B243
 behavioural psychology B162
 behavioural research B144,B234
 behavioural responses B160
 behavioural science B71
 behavioural scientists B163
 behavioural toxicology A76,B7,B229
 behavioural, children's B71
 behaviours that promote health B204
 behaviours, anti-social B182
 behaviours, off-task B97
 behaviours, peer B76
 behaviours, socially interactive B234
 behaviours, wellness B22
 behcet's disease A58,B99
 beliefs, health B76
 beliefs, philosophic B100
 below grade B199
 below-average household incomes B68
 bending A32,B208
 benefits, loss of B78
 benefits, psychological B39
 benefits, unemployment B151
 benzal chloride B178

- benzene A14,A15,B23,B69,B116,B127,B149,B226
benzene hexachloride A58
benzyl chloride B178
benzyl phthalate, butyl B178
beta particles A62,B20
between men and women, equality B13
beverage containers A58
beverages A58
beverages, caffeinated A66,B135
bhc A58
bias crime B67
biases towards homosexuality, societal B16
biases, personal B173
bibliographies B36
bibliography B176,B207
bifida, spina B131
binders A13,B14
binders, formaldehyde-based resin B106
binding materials B137
bioaccumulative substances B180
biochemical effects B158
biochemically predisposed B147
biologic mechanisms B75
biological chain reactions B90
biological effects B16,B57,B60,B102
biological factors B98
biological hazards A75,B88,B241
biological monitoring B129
biological organisms B37
biology of aging B2
biology, cell B67
biology, molecular B67
biopsy, lung B184
biphenyls, polybrominated B188
biphenyls, polychlorinated A54,B180
birth B59,B131
birth defects A24,A58,B99
bisexual A46,A47,B201
bisexuals A48
bisulfite B88
black families B233
blacks A46,B2,B67
blacks, lower-class B9
bladder B231
blame, externalization of B112
bleaches A55,B14
blind men and the elephant A1
blindness A58
blocks, primary B192
blood catecholamine B152
blood cell, white B53
blood circulation A60,B230
blood lead levels A20,B224
blood lead surveys B205
blood levels B20
blood pressure A53,B10,B22,B146
blood pressure, diastolic A70,B146
blood pressure, elevated B152
blood pressure, elevated diastolic B137
blood pressure, high A69,B47,B189
blood pressure, increased B44
blood sugar B187
blood sulfhydryl group A63
blood sulfhydryl B100
blood synthesis, red B116
blood tests B171
blood transfusions B100
bloodstream B188
blue collar workers 6
blue-collar unemployed A69,B47
blurred vision A9
board of education B113
board of education, Waterloo county B216
board, particle A53,B5,B79,B152
board, prefabricated woodchip B160
boards, particle A6
boards, school A76
bodily fluids A48
body burdens B180,B188
body defense mechanisms B119
body lead burdens B205
body odour B62
body of knowledge, core B176
body tissues A82
body/mind interaction B54
bone age B214
bone fractures B30
bone strength A29
bone-joint problems A69,B47
bones A52
bonus money B39
books A8,B69
boss B44
bottom income groups B8
boundaries, inside/outside B113
boys B21
boys' domain B84
brain A8,A25,B119,B125,B159,B199
brain damage B83,B223
brain damage, minimal B167
brain deterioration B5
brain development B33
brain disfunction, minimal B167
brain disorders, organic B160
brain dysfunction syndrome, minimal B113
brain dysfunction, minimal B33,B167,B229
brain electrophysiology A83
brain function A8,B108,B170,B198
brain imaging B108
brain malfunction A75,B178
brain research B108
brain synaptic function A62,B102
brain tissue B232
brain tissues B102
brain's electrical activity A82,B108
brain's state of health B108
brainwave analysis, computerized 10
brainwave measurement technology A82
brainwaves B214
brake emissions B101
brakes B100
brakes, automobile A20
breadwinner A36,B39
breakdown, family B112
breakup, marriage A47,B86
breast milk A57,B136
breast milk samples B136
breath, shortness of A6,B50
breathing A60,B230
bridging programs A38,B44
broader society B212
brokers, real estate B233
bronchial asthma A13,B14,B143,B242
bronchial sensitivity B153
bronchiectasis A75,B169
bronchiolitis A40,B172
bronchitis A75,A77,B29
bronchitis, inflammatory A13,B14
bronchitis, recurrent B169
brother B211
brutality A38
brutality, policy B64
budgets, reduced cleaning B76
buffers A72
building B19
building accessibility B128
building characteristics B114
building code B85
building code, municipal B78
building design 17,B13,B85,B114,B157,B222,B238

- building design, low-pollution A7,B199
- building envelope, tightly-sealed B199
- building illness B178,B206,B207
- building managers A74
- building material B92,B137,B181
- building material, synthetic B137
- building materials 5,15,A6,A15,A25,A54,A77,B6,B20,
B22,B31,B62,B79,B83,B89,B97,B103,B104,B106,B137,B139,
B140,B203,B220,B221
- building performance, total A10,B75
- building products A6
- building related accidents B114
- building shell B222
- building syndrome, sick A9,A17,A82,B3,B35,B73,B76,
B140,B153,B177,B237,B238
- building syndrome, tight B18,B149
- building user complaints B3
- building ventilation B211
- building, community school B43
- building, energy efficient B92
- building, office B115,B149
- building, tightly-sealed B16
- building-related illness A9,B172
- buildings B21,B66,B76,B89,B91,B92,B115,B124,B146,
B194,B221
- buildings, academic B212
- buildings, air-tight B206
- buildings, airtight B124
- buildings, airtight office B220
- buildings, allergens in school B113
- buildings, commercial B20,B113
- buildings, deep-core B85
- buildings, emergency evacuations of B103
- buildings, energy efficient B89
- buildings, european B227
- buildings, falls in B198
- buildings, high-rise A22,B21,B183
- buildings, hospital A16,B207
- buildings, large B184
- buildings, low-pollution B198
- buildings, maintenance of B215
- buildings, office B131,B206,B217,B220
- buildings, public A19,B43,B54,B209
- buildings, public access B73
- buildings, renovating B9
- buildings, residential A74,B19,B57,B184,B242
- buildings, school A11
- buildings, sealed B171,B206,B217
- buildings, sealed office A16,B35,B207
- buildings, tight B62
- built environment B85
- built environment, health and B85
- built environment, urban B107
- built-in furniture B106
- burden of family B34
- burden of home responsibilities B209
- burdens B123
- burdens, body B180,B188
- burdens, body lead B205
- bureaucrats B188
- burglarized A7
- burglary, people's reactions to B113
- burning B241
- burning eyes A6
- bus depots A19
- buses B210
- buses, school A9,B22,B177
- business B188
- business, computer-based A18,B37
- butyl benzyl phthalate B178
- byproducts B37
- byproducts, combustion B60
- cadmium A13,A15,A20,A25,A52,A58,B18,B38,B72,B116,
B189,B190,B200,B213
- caffeinated beverages A66,B135
- caffeine B39,B116
- calcine B181
- calcium A56
- calcium exchange A62,B102
- calibrated instruments B3
- cameras, time-lapse A10
- campaigns, safety B193
- campers B41
- can, toxins A57
- Canada B8,B128,B131,B142
- Canada fitness survey B137
- Canada health survey B46,B137
- Canada's cities B192
- Canada's food guide A56
- Canada, eastern B70
- Canada, sex and law in B112
- Canada, statistics B194
- Canadian centres of research B91
- Canadian faculties of medicine B204
- Canadian health and social policies B142
- Canadian health care B90
- Canadian health care system B224
- Canadian homes B132
- Canadian law B212
- Canadian medical schools B91
- cancer 12,A53,A57,A76,B20,B31,B42,B46,B59,B62,B74,
B88,B99,B112,B132,B146,B152,B153,B215,B217,B231,
B241
- cancer chemotherapy B3
- cancer incidence B74
- cancer prevention B46
- cancer promoters, weak B132
- cancer rates A58
- cancer risk B59
- cancer, childhood B74,B165
- cancer, etiology of childhood B74
- cancer, formaldehyde-induced B83
- cancer, human B174
- cancer, lung A6,A14,B11,B42,B120,B121,B132,B146,B174,
B223
- cancer, pediatric B74
- cancer, prevention of B46
- cancer, skin A61
- cancers A58,B192
- cancers, lymphatic B132
- cancers, smoking-related B231
- capabilities, language B120
- capabilities, perceptual A50,B209
- capacitive coupling B102
- capacity of the environment B222
- capacity to cope B107
- capacity, child's A28
- capacity, forced vital B6
- capacity, mental B63
- capacity, vital B202
- car exhaust A74,B169
- carbohydrate diet, high protein-low B39
- carbon dioxide A5,A19,B43,B124,B162,B227
- carbon disulfide B235
- carbon monoxide A5,A8,A15,A16,A19,A24,A52,A53,A76,B5,
B7,B19,B27,B29,B33,B41,B42,B43,B49,B57,B63,B80,B88,
B91,B92,B101,B107,B118,B121,B136,B141,B143,B155,B184,
B191,B195,B203,B211,B219,B228,B232
- carbon monoxide exposure, low-level B7
- carbon monoxide poisoning B221
- carbon paper B69
- carbon tetrachloride B27
- carbon, activated B226
- carbonless copy A16
- carbonless copy paper B30,B129
- carbonless paper A8,B69
- carboxyhemoglobin A5,A20,A41,A53,B195,B221
- carcinogen B27,B80,B149
- carcinogen, epigenetic B83
- carcinogenesis B75

- carcinogenic B6,B16,B52,B117,B159,B175,B221
 carcinogenic risk B175
 carcinogenic substances B12
 carcinogenicity A14,A53,B83,B146,B156,B228
 carcinogens A15,A22,A57,B42,B46,B75,B79,B120,B168,
 B191
 carcinogens, chemical B14
 carcinogens, dietary B4
 cardiac arrhythmia B27,B169
 cardiac arrhythmias A83,B171
 cardiac problems B198
 cardiovascular A52,A55,B17
 cardiovascular disease A53,A70,B67,B137,B171,B174
 cardiovascular disease, diet and B142
 cardiovascular diseases B152,B171
 cardiovascular diseases, inflammatory A56,B171
 cardiovascular distress 12
 cardiovascular functions B225
 cardiovascular reactivity to mental stress B146
 cardiovascular stress A41
 cardiovascular system B94,B171,B191
 care settings, tertiary B233
 care units, environmental B190
 care, acute A75,B56
 care, ambulatory B22
 care, child A34,B25,B35,B39
 care, community-based B64,B156
 care, culturally acceptable A39,B7
 care, day A36,B25
 care, extra-parental B25
 care, institutional B219
 care, patients in acute B233
 care, prenatal B131
 care, primary B63
 care, quality of B156
 care, special B86
 caregivers B54
 caregiving professionals B54
 caretaking, participating in B7
 caring neighbourhoods, integrated B119
 carpet manufacture B46
 carpet shampoos B115
 carpet sweeper, manual B165
 carpeting B96
 carpetlayers B10
 carpets B76,B137
 carpets, needle-felt A9,B76
 carpets, wool B46
 carrying B208
 cars B210,B229,B241
 case studies B200
 caesarean delivery B131
 castles B50
 cat hypothalamus B186
 catalysts A55,B14
 catalytic gasoline exhaust B32
 catecholamine, blood B152
 categories, occupational B208
 caulking compounds B137
 causal connection B112
 causal relationship B243
 causation of allergy B239
 cause of disease B142
 cause of sexism B13
 causes of ill health, socioeconomic B133
 cell biology B67
 cell membrane B28
 cell, white blood B53
 cell-mediated immunity B232
 cell-mediated responses B232
 cellulose A54,B103
 cellulose acetate, methyl A14,B226
 cement, portland B162
 cement, self-leveling B82
 censorship B4
 central nervous system A8,A76,B23,B80,B119,B125,B157,
 B164,B223,B225,B235
 central nervous system damage A15,A57,B215
 central nervous system depressants B172
 centre, day-care B115
 centres of power B25
 centres of research, Canadian B91
 centres of research, national B90
 centres, day care A19
 centres, health science B91
 centres, pollution relief A11
 centres, rape crisis A34,B35
 centres, urban B86,B191
 century disease, twentieth B216
 ceramics A5,B84
 cerebral allergy B127,B160
 cerebral atrophy B10
 cerebral disturbances B106
 cerebrovascular disease B108
 cerebrovascular disorders, aged B164
 chaetomium B189
 chain reactions, biological B90
 chairs, plastic B195
 chalk dust A8,B199
 challenges, incitant A83,B169
 chamber, climate B124
 chamber, environmental B219
 change B69
 change strategies, behavior B22
 change, dietary B39
 change, economic B140
 change, efficacy of lifestyle B76
 change, physiological B230
 change, self-directed B109
 change, social B133,B163
 change, strategies for B35
 change, technological A82
 changes in memory B56
 changes, air B227
 changes, attitudinal 17,A2
 changes, behavioural B17
 changes, dietary A56
 changes, economic A69,B112
 changes, institutional A85
 changes, lifestyle B151
 changes, massive B134
 changes, neurologic A58
 changes, personality A13
 changes, structural A71
 changes, visual B56
 changing human service needs B224
 changing rooms A12
 changing shift, frequently B28
 changing values B120
 characteristics, building B114
 characteristics, chemical B158
 characteristics, demographic B43
 characteristics, environmental B106
 characteristics, human A1
 characteristics, maternal B131
 characteristics, minority 16
 characteristics, personal B148
 characteristics, physical B158
 characterization, stereotyped B50
 charged conductors B102
 charter of rights and freedoms B218
 chauvinistic male physicians B136
 chemical B168
 chemical additives B47,B210
 chemical agents A75,B75,B241
 chemical analysis B91
 chemical carcinogens B14
 chemical characteristics B158
 chemical companies A57
 chemical composition B156

- chemical contaminants B53
chemical evaporants B220
chemical exposure B150
chemical exposure, massive B125
chemical exposure, toxic B109
chemical exposures B73,B171
chemical fumes B125
chemical hazards B162
chemical hypersensitivity A55,A82,B81
chemical hypersusceptibility A17,B172
chemical intolerance B105
chemical irritants B130,B190
chemical laboratories B172
chemical manufacturing facilities B193
chemical overexposure B168
chemical patterns B19
chemical pollutants B18,B229
chemical pollution B33,B127,B172
chemical sensitivities B47,B167
chemical sensitivities, food or B168
chemical sensitivity B50,B53,B111,B138,B149,B150,
B160,B167
chemical sensitization B97
chemical substances B47
chemical substances, harmful B215
chemical susceptibilities B209
chemical susceptibility A6,B22,B126,B160,B161,B166,
B168,B189
chemical toxicity B180
chemical wastes B23
chemical workers B114
chemically hypersensitive 6
chemically sensitive A62
chemically sensitive children B150,B188,B189
chemically sensitive patients B171
chemically susceptible B199
chemically susceptible individuals A11,B212
chemically susceptible population B198
chemically-induced personality disorders B9
chemically-less-contaminated environment B171
chemicals, agricultural B27,B60
chemicals, cleaning A8,B199
chemicals, controversial B116
chemicals, environmental B229
chemicals, hypersensitivity to non-physiologic B193
chemicals, industrial B3,B15,B45
chemicals, janitorial B167
chemicals, man-made B16
chemicals, overload of synthetic B170
chemicals, pungent B209
chemicals, residues of B215
chemicals, sensitizing B3
chemicals, toxic B45,B47,B94,B177
chemicals, volatile organic 15,B5
chemicals, xenobiotic B72
chemistry, reactive B155
chemists A74
chemotherapy, cancer B3
chest pain A83,B169
chest tightness A9
child abuse 12,A25,A28,A66,B34,B71,B112,B179
child as citizen B212
child care A34,B25,B35,B39
child care workers A32,B209
child custody A47,B2
child development B214
child health B138
child health and safety B83,B130
child poverty B149
child victims of welfare B45
child welfare act B179
child's capacity A28
child's competence B202
child's environment B71
child's functional requirements B122
child's mental health B71
child's perceptions A50
child's perspective B193
child, autonomy of the B162
child, chronically ill B7
child, curious B222
child, disabled B122
child, exploitation of a A26
child, hearing impaired B153
child, ill-treatment of a B34
child, punishment of a B34
child-care facilities A38,B44
child-centered view B25
child-centred A37
childbirth B136
childhood B71,B173
childhood accidents B70
childhood allergies B167
childhood autonomy B162
childhood behavior disorders B71
childhood cancer B74,B165
childhood cancer, etiology of B74
childhood household injuries B52,B53
childhood infection B172
childhood risk B209
childhood, behaviour problems during B134
children A4,A22,A36,A37,A50,A51,A53,A75,B12,B20,B21,
B22,B24,B25,B33,B41,B45,B52,B59,B64,B65,B66,B67,B70,
B71,B73,B74,B77,B84,B86,B92,B97,B104,B108,B110,B112,
B113,B115,B121,B122,B125,B127,B130,B132,B135,B138,
B139,B144,B145,B147,B151,B153,B159,B162,B172,B177,
B179,B183,B184,B193,B194,B202,B203,B205,B209,B210,
B212,B213,B216,B217,B218,B222,B225,B229,B234,B242,
B243
children around, school B41
children as people B235
children as persons B212
children at risk B202
children in design, involving B235
children of smokers A5
children of the poor A70
children's accidents B193
children's aggressive behaviour B243
children's aids societies B179
children's behavioral B71
children's behaviour B243
children's environments B235
children's health B183
children's hospital B65
children's knowledge B243
children's mental health B183
children's performance in traffic B202
children's rights B25,B212
children's risk B165
children, access to B86
children, chemically sensitive B150,B188,B189
children, circumstances of B213
children, disabled B151
children, elementary school B222
children, handicapped B122
children, harm to B70
children, health status of B138
children, hyperactive A9
children, incapacity of B144
children, japanese B216
children, kindergarten A63,B77
children, learning impaired B110
children, low income B183
children, needs of B212
children, participatory roles of B25
children, pre-school B205
children, prejudice against B50
children, protection of B179
children, reducing risks to B209
children, research involving B144

- children, respiratory symptoms in B108
 children, school B92,B107,B109,B137,B158,B214
 children, vulnerable A26,B243
 children, welfare of the B144
 children, young B84,B199,B205,B212,B238
 chinese, immigrant B7
 chipboard B5
 chlordane A57,B161
 chloride resins, vinylidene B67
 chloride, benzal B178
 chloride, benzyl B178
 chloride, methylene A41,B27
 chloride, vinyl A15,B27,B116,B243
 chlorinated hydrocarbon B136
 chlorinated hydrocarbon pesticides A82,B117
 chlorinated hydrocarbons B136,B170,B232
 chlorinated naphthalenes B136
 chlorinated pesticides B188
 chlorinated solvents A26,B165
 chlorine A8,B69,B169,B211
 chloroform B116
 chlorophenols B192
 choice A7,A81
 choice discrimination errors B7
 choice, consumer A81
 choice, environmental 3,17,A79
 choice, freedom of A65,B163
 choice, individual B142
 choice, informed B63
 choices, occupational A76
 choices, social B148
 choices, worklife A18,B87,B93
 cholesterol level, elevated serum B137
 cholesterol levels, high B44
 cholesterol, serum A70,B152
 cholinesterase B154
 chromatographic study, gas B56
 chromatography, gas B117,B243
 chromatography, high-resolution gas A82
 chromatography-mass spectrometry, gas A82
 chromatography/mass spectrometry, gas B227
 chronic B157,B207
 chronic care environments A29
 chronic cough A5,B87
 chronic disease B120
 chronic diseases B229
 chronic effects B90
 chronic exposure B72
 chronic fatigue A6
 chronic illness B122
 chronic low back pain A65,B66
 chronic low-level exposures B49
 chronic mental disorders B216
 chronic occupational exposure B69
 chronic unemployment picture B33
 chronic under-nutrition B218
 chronically ill child B7
 chronically mentally ill B207
 cigarette B176
 cigarette smoke A74,B83,B169,B217,B220
 cigarette smoke, sidestream B73
 cigarette smoking A70,B22,B44,B54,B74,B87,B89,B93,
 B94,B97,B132,B135,B137,B144,B240
 cigarettes A50,A66,B10,B24,B179
 circulation, blood A60,B230
 circumstances of children B213
 circumstances, economic B213
 cities A19,A21,A38,A66,B12,B58,B94,B183,B230,B233
 cities, american B109
 cities, Canada's B192
 cities, livable B162
 cities, masculine B183
 citizen action, support for B144
 citizen and community, roles of B133
 citizen participation A67,B9,B23
 citizen, child as B212
 citizens apartment, senior B106
 citizens apartments, senior A6
 citizens groups B147
 citizens, second-class corporate B39
 citizens, senior B90,B204
 citizenship A44
 citizenship status B95
 citizenship status, true B212
 city B68,B229
 city life, stressors of B12
 city, health in the B229
 city, healthy B58
 city-dwellers B217
 civic rights A49,B139
 civil disorders B109
 civil rights A49,B139
 civil rights statutes B79
 civil suits A38
 civilian A66
 civilization B119
 claim, reasonable B79
 claims B135
 claims, health B142
 clandestinely B61
 clarity, job B66
 class B183
 class background B183
 class homogeneity, social B233
 class inequities, social A78,B134
 class, middle B181
 class, social B135,B202
 class, white middle B120
 class, working B114
 classes, academic health B76
 classes, remedial A70,B45
 classes, university conditioning B76
 classification, anatomical B157
 classroom air B110
 classroom behavior, non-adaptive B145
 classroom behaviour B110,B162
 classroom performance B145
 classroom, ecology A11
 classroom, experimental B97
 classroom, low-pollution B216
 classroom, well-conducted B222
 classrooms B166,B212
 classrooms, ecological B113
 classrooms, portable B210
 classrooms, science B24
 classrooms, windowless B154,B222
 claviceps B189
 clean air A51
 clean air standards B30
 clean water B229
 clean water supply B116
 cleaner materials 3
 cleaner, vacuum B165
 cleaners, electronic air B226
 cleaning budgets, reduced B76
 cleaning chemicals A8,B199
 cleaning devices, air B207
 cleaning products, scented B166
 cleaning, air B146,B175
 cleanup, environmental B94
 clearance system, nonspecific B188
 clerical B240
 clerical women B39
 clerical workers A32,B31,B69,B208
 clienthood B133
 clients B95,B216
 climate B44
 climate chamber B124
 climate complaints, indoor B76
 climate control B124

- climate, emotional B211
- climate, indoor B238
- climate, perceived B98
- climate, social B176
- climate, thermal B123
- climates, cold B232
- climatic pathogens B177
- clinical and sub-clinical effects B111
- clinical ecology B18,B27,B53
- clinical histories B170
- clinical manifestations B240
- clinical payoff B31
- clinical programs B204
- clinical trials B22
- clinics, independent occupational health B241
- close relationships A35,B99
- close ties, number of B87
- closed offices B131
- closed spaces B131
- closeness A35
- closeness to nature B183
- closeting 13
- closets B71
- closings, plant B140
- clothing A55,B151
- clothing, cotton B191
- clothing, synthetic A32,A55
- clouds, radioactive B61
- clutches B100
- co-ordination A4
- co-resident distrust B213
- co-worker ratings, aggregate B176
- coal A22,B217
- coaster, roller B138
- code, building B85
- code, criminal B179
- code, municipal building B78
- Code, Ontario Human Rights B41
- code, rigid no-noise B231
- codes and standards B128
- codes, human rights A47,B112
- coercion A28,B162
- coffee A66
- cognitive B202
- cognitive development B64
- cognitive factors B76
- cognitive impairment B87
- cognitive influences B76
- cognitive overload B181
- cognitive reactions B168
- cohesion, peer A65,B66
- coil units, fan A54
- coil, fan B82
- cold A63,B100
- cold and flu symptoms B103
- cold climates B232
- collaborative action B34
- collagen A74,B170
- collagen vasculitis B171
- collar workers, blue 6
- colleagues B44
- collective bargaining agreements B185
- collective control B196
- collective determinants of lifestyles B8
- collective visions B183
- college, community B44
- color B84
- colour B84
- colour rendering A10
- colour, skin 13
- combustion appliances B41
- combustion appliances, indoor B73,B219
- combustion appliances, unvented A5
- combustion byproducts B60
- combustion devices 5,15
- combustion products B91,B166,B220
- combustion sources A52
- combustion, oil A22,B217
- combustion, wood B141
- combustionproducts B120
- comfort B17,B62,B136,B206,B238
- comfort complaints B207
- comfort, physical B66
- comfortable B29
- comforts, household B33
- coming out A47,B86
- commercial B231
- commercial buildings B20,B113
- commercial inks B47
- commercial settings A16
- commitment to the future B63
- committee, federal/provincial advisory A5
- committees, joint health and safety B185
- commodities, agricultural A57,B189
- common illnesses, mimicry of B5
- common social environment B176
- common space B43
- communicable diseases A32,B209
- communicating values B111
- communication B8
- communication, honesty in A68
- communications B117
- communities B147,B204,B222
- communities, rural B86
- communities, semirural A72,B68
- communities, unprepared B156
- community A72,B26,B37,B58,B68,B86,B133,B148
- community activities A43
- community college B44
- community development A67,B119
- community disintegration B61
- community economy B140
- community education A28,B162
- community fitness program B76
- community focus B63
- community health B36,B109
- community health care workers, native B96
- community integration B219
- community investment A44
- community involvement B148
- community life B95
- community mental health B140
- community mental health variables B140
- community noise A65,B32
- community participation B36,B95,B239
- community planning B231
- community psychiatry B79
- community reinvestment B156
- community resistance A7,B78
- community resources B127
- community response B23
- community responsibility A78
- community school B42
- community school building B43
- community settings B146
- community's economic base B37
- community, energy conserving B71
- community, human B37
- community, medical B46
- community, native B96
- community, regenerating B133
- community, roles of citizen and B133
- community, social B71
- community, vision for B133
- community-based care B64,B156
- community-based public health B82
- community-based rehabilitation A44
- commuters B183
- commuting time A18,B37
- companies, chemical A57

- companionship A35
 comparison populations 9
 comparisons, epidemiological A83
 compelled to obey A27,B50
 compensation B241
 compensation legislation, workers B185
 compensation policy B241
 compensatory damages B79
 competence A22,A56,B21,B54,B99,B184,B209
 competence, child's B202
 competence, feeling of B85
 competence, functional A66
 competencies, behavioral B147
 competencies, range of B202
 competition, industrial A85
 complaint B73
 complaints B227
 complaints, air quality B212
 complaints, building user B3
 complaints, comfort B207
 complaints, employee health B131
 complaints, health B35
 complaints, indoor climate B76
 complaints, somatic B101
 complete health A1
 complex, hermetically sealed office B18
 complex, three-tower government office B131
 compliance A46,B2,B134
 compliance, maternal B131
 compliance, program B76
 composite B70
 composition, chemical B156
 compound emissions, organic B137
 compound, organic B227
 compounds, caulking B137
 compounds, fat-stored foreign B188
 compounds, industrial B157
 compounds, inorganic B42
 compounds, nitrogen B91,B92
 compounds, organic B140,B184
 compounds, organochlorine A58
 compounds, organophosphorous B232
 compounds, resin B14
 compounds, sulfur B91
 compounds, synthetic B168
 compounds, total volatile organic B227
 compounds, toxic B168
 compounds, volatile organic A54,B57,B140,B142,B243
 comprehension, reading A8,B22,B150
 comprehensive environmental control B166
 comprehensive standards B35
 comprehensiveness B63
 compromise B231
 compulsive, sexually A35,B99
 compulsory attendance B162
 compulsory attendance laws B162
 compulsory schooling A10,B162
 computer B37
 computer-based business A18,B37
 computer-mediated home-based work B39
 computerized brainwave analysis 10
 computerized diagnostic system A82,B108
 computerized eeg B214
 computerized imaging B108
 concentrate B1
 concentrate, impaired ability to B227
 concentrate, inability to B50
 concentrating, difficulty in B64
 concentration A58
 concentration difficulty B72
 concentration, hemoglobin B83
 concentrations, threshold B21
 concepts, philosophical B142
 conceptualizing B184
 condensation B199
 condensers, evaporative B236
 condensing altitude A33
 conditioned A60
 conditioning classes, university B76
 conditioning ductwork, air B21
 conditioning equipment, air B28
 conditioning, air A54,B54,B181
 conditioning, social B221
 conditions of exposure B154
 conditions of work B33
 conditions, acute respiratory B24
 conditions, domestic stair B114
 conditions, environmental B206,B232
 conditions, growth B103
 conditions, living B182
 conditions, moist B215
 conditions, safety B177
 conditions, skin B209
 conditions, social A64
 conditions, unhealthy working A38
 conditions, unpleasant work B44
 conditions, weather A6
 conditions, working B44,B206
 conduct, discriminatory A38,B79
 conduct, racist B79
 conduction, nerve B157
 conductors, charged B102
 conference proceedings B147
 confidence A27,A35,B84,B92,B220
 confidence, low self B44
 configuration B182
 confined environments B53
 conflict B142
 conflict of attitudes B37
 conflict, family A65,B66
 conflict, role B44
 conflicts, environmental B9
 conflicts, freedom from intergroup B164
 conflicts, frequent B183
 conflicts, intergroup A66
 conformity B235
 confused A44,A66
 confusion A8,B166
 confusion, mental A8
 congenital malformations B193
 congestion, nasal A16,B30,B31
 congestion, traffic A21,B109
 congregate housing A31,B111
 congregate housing for elderly B111
 conjunctivitis B5,B31
 connection, causal B112
 connection, environmental/social B9
 connections B9,B43
 consciousness raising B85
 consensus B54
 consent, people acting through B133
 consent, process of A28
 consequences of stress B40
 consequences, environmental B222
 consequences, injury B135
 consequences, social B182
 conservation B237
 conservation strategies B120
 conservation, energy A5,A19,A77,B16,B19,B28,B57,B66,
 B91,B92,B174,B194,B197,B211,B219,B242
 conserving community, energy B71
 consideration B34
 consideration, genuine B231
 considerations, economic A15,B185
 constantly, judged B50
 constitutional factors B37
 constraint, behavioral B181
 constraints B167
 constraints, economic B37
 constraints, environmental B37

- construction B19,B76,B222
construction industry B134
construction material B5
construction materials B6
construction sites B134
construction workers A14,B134
construction, floor B106
construction, interior B199
consultation, effective B44
consumer choice A81
consumer guide B47
consumer market B213
consumer organizations B63
consumer perspective B95
consumer products A50,B47,B97,B179
consumer responsibility B8
consumer status B133
consumers A57
consumption, alcohol A76,B76,B80
consumption, excessive alcohol B137
contact dermatitis B151
contact dermatitis, allergic A16,B129
contact lenses A53,B173
contact with individuals, direct 12
containers, beverage A58
containers, polyethylene food B105
contaminants B14,B136
contaminants, chemical B53
contaminants, low-level B27
contaminants, microbial B74
contaminants, reducing airborne B145
contaminants, volatile organic A77
contaminated soil A25
contamination, drinking water B200
contamination, fungal B184
contemporary science B91
content, low nitrogen B103
content, nitrogen A54
contentment B115
context of the family B212
context, human B93
continuity A30,B42
continuous low-speed fan B165
contraceptives, oral A70,B137
contractions, premature ventricular A55,B191
contractors, independent A18,B39
contradictory realities B183
contribution A43
control 11,17,19,A27,A44,A66,A68,A75,A79,A81,B99,
B138,B184
control in the physical environment B65
control of people B133
control over one's life 5
control strategies B137
control, air pollution B184,B223
control, climate B124
control, collective B196
control, comprehensive environmental B166
control, dust B48
control, environmental A76,A81,A83,B68,B71,B170
control, family B66
control, individual 3,B172
control, injury B179
control, little job B209
control, need for A67,B99
control, perceived A65,B32
control, personal 7,A30,B106,B233
control, pest 5
control, pollution B27
control, sense of B111
control, source B146
control, worker's B122
controls B95
controls, dietary A76
controls, rigid environmental B170
controversial chemicals B116
cooking B220
cooking appliances, gas B108
cooking devices B91
cooking equipment B22
cooking fuel, gas B55
cooking fumes B55
cooking stoves, gas B155
cooking stoves, gas fired B73
cool-mist humidifiers B224
cooling towers A54,B21,B236
cooperation, international A22,B222
cooperation, interprofessional B179
cooperative measures B34
coordinated approaches B222
coordination A29,A52,B187
coordination, disturbance of B7
coordination, impaired B87
coordination, impaired motor B23
coordination, policy B144
cope, capacity to B107
cope, inability to B167
copier paper B69
coping A72,B97,B195
coping behaviour B61
coping efforts, health-damaging B61
coping mechanisms B61,B68
coping repertoires B55
coping resources A65,B32
coping skills A64,B54,B71
coping strategies A46
coping with physical disabilities B148
copper A22,A58,B70
copy paper, carbonless B30,B129
copy, carbonless A16
copying, dyeline A61,B62
core body of knowledge B176
cornerstones of life B57
coronary disease B22
coronary heart attacks B44
coronary heart disease B22,B44
coronary spasm A56,B171
coronary-prone behavior, type-a B50
corporate citizens, second-class B39
corporation, profit-oriented B188
correcting mistakes B92
cortical inhibition processes B24
cosmetics B22,B81
cosmetics, scented A44
cost, health B27
cost-effectiveness B63
cost/benefit B131
costs of unemployment, human B112
costs, housing A71,B149
costs, human B112
costs, ventilation B49
cotton A55
cotton clothing B191
cotton fabric B191
cotton garments B191
cough A51,B115
cough, chronic A5,B87
cough, hacking A12
coughing A53,B80,B173,B178
Council on Health Strategy, Premier's B64
Council, Swedish National B123
counseling B178
counselling, health management B204
countries, nordic B83
county board of education, waterloo B216
couples, gay A47
coupling, capacitive B102
courts B79
coverage, press B129
craft materials B84

- craft materials, art and A5
 crash involvement B130
 crashes, road B130
 cravings, food A77,B29
 creative ventilation A7
 creativity B196
 creators, active B235
 credibility B12
 credits, income B149
 crime A21,B109,B147
 crime and delinquency B147
 crime, adolescent B112
 crime, bias B67
 crime, fear of A66,B147
 crime, violent B178
 crime-prevention B147
 crimes, hate B67
 criminal behaviour B178
 criminal code B179
 criminal justice system B67,B147
 criminal prosecution B185
 criminology B187
 crises, domestic A66
 crises, environmental B58
 crises, life B184,B195
 crisis B54,B122
 crisis centres, rape A34,B35
 crisis, aids B41
 crisis, employment B148
 crisis, environmental B163
 crisis, families in A78
 crisis, rental housing B62
 crisis, world economic B138
 crowded freeways A65
 crowded urbanites A67,B12
 crowding A21,A65,B12,B32,B68,B109,B140,B158,B181,
 B182
 crowding, residential B183
 cues, familiar environmental B65
 cultural background A38,A39,B86
 cultural backgrounds B26
 cultural differences 3,A38,A39,B7,B192,B216
 cultural diversity A38,B44
 cultural minorities A24,B50
 culturally acceptable care A39,B7
 culture 3,A2,A29,A80
 culture medium B28
 culture of prejudice B211
 cultures A7,B4
 culturing of bacteria A54
 cumbersome B122
 cumulative exposure A6,B227
 cumulative exposures B166
 cupboards B71,B128
 curative approach, individualized B213
 cure B233
 curious child B222
 current levels B156
 current medical model B204
 currents B102
 currents, internal B102
 curricula, medical school A49,B16
 curricula, school A26
 curricula, undergraduate B204
 curriculum design B162
 curve, dose-response B45
 custody rights B86
 custody, child A47,B2
 customary lifestyle, loss of B18
 customs A39
 customs for managing illness B7
 customs guidelines B4
 cutting fumes B123
 cutting oil A26,B165
 cyanide, hydrogen B121
 cycle of poverty A33,B134
 cynicism A43,B95
 cystic fibrosis A40,B172
 daily activities B1
 daily use, products in B47
 damage, brain B83,B223
 damage, central nervous system A15,A57,B215
 damage, environmental B70,B236
 damage, environmentally-induced B171
 damage, irreversible lung B73
 damage, kidney A13
 damage, lung B205
 damage, minimal brain B167
 damage, property B192
 damage, water A9,A54,B76
 damages to health B12
 damages, compensatory B79
 damages, social B157
 damp mop B165
 dangerous habits A4
 dangerous machinery A15,B73
 Danish National Inventory B6
 Danish research B139
 data 14
 data base on environmental health B237
 data systems B138
 data, evaluated B19
 data, experimental B91
 data, health B8
 data, health survey B143
 data, morbidity/mortality B237
 data, mortality B132
 daughter nuclides, radon A53
 daughter products A6
 daughter products, radon A54
 daughters, radon B89
 day care A36,B25
 day care centres A19
 day-care A37
 day-care centre B115
 daylight B240
 daylight, natural A10,B154
 days, disability B24
 dbcp A57,B116,B193
 DDT A57,A58,B99,B106,B116,B168,B193,B215
 de-institutionalized patient B216
 deactivators A55
 deadlines A32
 deadlines, unrealistic B209
 death A4,A29,A50,A57,A58,B89,B114,B134,B143,B149,
 B179,B215,B229,B240
 death syndrome, sudden infant B5
 death, increased risk of B80
 death, infant A38,B192
 death, life and A47,B54
 death, residential accidental B4
 death, risk of A70
 death, sudden infant A40
 death, work-related B241
 deaths A25,B130,B168,B179
 deaths, multiple B185
 debilitating B73,B219
 decay of physical settings B163
 decay products, radon B89
 decent jobs, access to A71
 decision overload A65
 decision processes B209
 decision-making B162
 decision-making process B43,B44
 decision-making roles B35
 decision-making, sexual B173
 decisional overload B183
 decisions from numbers B201
 decisions, design B1
 decisions, environmental B9

- decisions, physical planning B9
 declining influence of the generalist B135
 decongestants, solvents, pollutants, nicotine B48
 decorative possessions B112
 decreased attention span B150
 decreased immunological responsiveness B232
 decreased interest in reading B243
 decrement, performance B118
 deep-core buildings B85
 defects, birth A24,A58,B99
 defence mechanisms A78,B237
 defense mechanisms, body B119
 defense mechanisms, pathological B90
 defense mechanisms, respiratory tract B187
 defensive functions B56
 deficiencies, design B126
 deficiencies, nutritional A56,B23,B73
 deficiency syndrome, acquired immune A47
 deficiency, nutritional A41,B126
 deficiency, sulfite oxidase B88
 deficit, neurological B223
 deficits, learning B59
 deficits, memory B150
 deficits, neurological A25
 deficits, neuropsychologic B145
 deficits, performance A76
 deficits, skill B150
 defiled B114
 degenerative disease B143
 dehumanization A65,B122
 dehumanization of men B99
 dehumanized A35
 dehumanizing B229
 dehumanizing environments B44
 delayed effects B235
 delayed health hazards A58,B192
 delayed responses A64
 delayed responses to stressors B40
 delayed-type hyper-sensitivity B14
 deleted from public attention B58
 deleterious reproductive outcomes B192
 delinquency B147,B178,B187
 delinquency, juvenile B73
 delinquents B147
 delinquency, crime and B147
 delivery of services B86
 delivery systems, health care 3,B156
 delivery, cesarean B131
 delivery, health care B16
 delivery, home B131
 demand/supply dilemma B156
 demands B107
 demands, job B44
 demeaned B211
 dementia A72,B87
 dementia, established B87
 democratic freedoms A66,B164
 democratic society B163
 democratization B185
 democratization, workplace A15
 demographic B37,B111
 demographic characteristics B43
 demographics A64,B118
 demoralizing A44
 denial B90
 dense urban neighborhoods A67,B12
 density A67,B68,B140
 density apartments, high B183
 density environments, high B182
 density environments, lower B182
 density measurement B228
 density, effects of B12
 density, high B182
 density, higher B183
 density, magnetic flux B15
 density, residential B183
 density, urban B12
 dental amalgam filling B94
 deodorants B23
 dependence A43,A67,B95,B119
 dependence on adults A28,B144
 dependence, minimal B156
 deploia B189
 depositions, acid B70
 depositions, wet sulfate B70
 depots, bus A19
 depressants A9
 depressants, central nervous system B172
 depressed A8,A72
 depression A9,A33,A65,B66,B80,B87,B101,B112,B134,
 B153,B176,B211
 depressive reactions A58,B72
 deprivation A47
 derailing B25
 dermatitis A8,A13,B14,B31,B69,B103,B129
 dermatitis, allergic contact A16,B129
 dermatitis, contact B151
 dermatitis, paper B69
 deschoolers B162
 deschooling B162
 desensitization B186
 design B91,B163
 design decisions B1
 design deficiencies B126
 design for health B85
 design for the elderly B56
 design of hospital environments B233
 design of programs B8
 design process B234
 design researchers B26
 design tools, basic B1
 design, architectural B140
 design, building 17,B13,B85,B114,B157,B222,B238
 design, curriculum B162
 design, domestic stair B114
 design, environmental B1,B26,B36,B42,B69,B85,B100,
 B111,B118,B122,B163,B234
 design, housing B42,B165,B199,B206,B226
 design, insensitive and elitist B1
 design, institutional B56
 design, involving children in B235
 design, job B185
 design, low pollution B184
 design, low-pollution building A7,B199
 design, participating in A67
 design, participation in A84
 design, physical B107
 design, product A50
 design, residential B32,B114
 design, safer stair B114
 design, space B183
 design, stair A4,B114
 design, urban B230
 designated substance B125
 designed furniture, poorly B208
 designers B1,B9,B235
 designs, traditional window B222
 desire to appear as normal B122
 desire, parents B122
 despair A69,B33
 destruction, mass A22,B222
 destructive lifestyle B152
 destructive rigidities B149
 detection, air pollution B91
 detective process 12
 detector, track-etch B89
 detergent dust B115
 detergent residue A19
 detergent residue, dried B115
 detergents B115

- detergents, soaps B60
 deteriorating paintwork B137
 deterioration in behavior A13
 deterioration, brain B5
 deterioration, structural A6,B38
 determinants of lifestyles, collective B5
 determined susceptibility, genetically B75
 detoxification treatment B188
 detroit B161
 devaluation A23,A79
 devalued A2,A40
 devaluing 3,6
 devaluing of women A32
 developing nations B4
 development B236
 development of technology B190
 development planning B222
 development, academic B150
 development, behavioral B64
 development, brain B33
 development, child B214
 development, cognitive B64
 development, community A67,B119
 development, economic B120
 development, human B131
 development, infant B131
 development, knowledge B144
 development, legislative B185
 development, post-industrial A69
 development, psychological B21,B183
 development, scientific research and B222
 development, self-help B39
 development, skills A26
 development, social B21,B222
 development, urban B162
 developmental effects B217
 developmental psychology B118
 developmentally disabled A43,B234
 deviant B27
 deviants B2
 devices, air cleaning B207
 devices, air sampling B184
 devices, air-purification B220
 devices, combustion 5,15
 devices, cooking B91
 devices, health monitoring 15
 devices, measurement 15
 devices, medical B86
 devoid of vegetation B208
 di-isocyanate, toluene A14
 diabetes A70,B131
 diabetes mellitus B137
 diagnostic B134
 diagnostic system, computerized A82,B108
 diagnostic systems B108
 diagnostic techniques, therapeutic and B53
 diarrhea A57,B103,B240
 diastolic blood pressure A70,B146
 diastolic blood pressure, elevated B137
 diesel engine exhaust B158
 diesel exhaust A14,B32,B48,B49,B117,B118,B223
 diesel exhaust emissions B117
 diet A26,A28,A76,B46,B142,B167,B180
 diet and cardiovascular disease B142
 diet and health B142
 diet, feingold B127
 diet, high protein-low carbohydrate B39
 dietary carcinogens B4
 dietary change B39
 dietary changes A56
 dietary controls A76
 dietary guidelines B142
 dietary intake, maximum permissible B72
 dietary manipulation therapy B49,B50
 dietary sulfites B88
 dietary therapy A76,B50
 diets A56
 diets, junk food B187
 difference, acknowledging B26
 difference, gender A33,B211
 difference, urban-rural B11
 differences B143
 differences in illness, sex B143
 differences in smoking rates, socioeconomic B145
 differences, behavioural B119
 differences, cultural 3,A38,A39,B7,B192,B216
 differences, exploring B26
 differences, gender B40,B88
 differences, geographic B74
 differences, hiding A79
 differences, individual B61,B209
 differences, sex B84,B143
 differences, tolerance of A17
 different B234
 differentness, tolerance of B34
 difficulties with partners B40
 difficulties, financial A69
 difficulties, gastrointestinal B29
 difficulties, learning B150,B187
 difficulties, practical B122
 difficulties, psychological B122
 difficulty in concentrating B64
 difficulty, concentration B72
 diffused light B62
 digit span test B11
 dignified shelter B78
 dignifying paid work B219
 dignity A7,A71,B12,B95,B163
 dignity, human B163
 diisocyanate, toluene B38,B151
 dilemma, demand/supply B156
 dilution B27
 dioxide emissions, sulphur B27
 dioxide, carbon A5,A19,B43,B124,B162,B227
 dioxide, nitrogen A5,A19,A41,A52,A53,B5,B7,B18,B19,
 B20,B41,B43,B57,B60,B65,B73,B74,B92,B93,B105,B109,
 B121,B135,B142,B147,B191,B196,B197,B202,B203,B211,
 B219
 dioxide, sulfur B18,B46,B65,B88,B93,B105,B121,B133,
 B141,B142,B154,B184,B191,B234
 dioxide, sulphur A5,A20,A22,A51,A52,B116,B196,B216
 dioxin A50,B116,B242
 dioxins B192
 direct contact with individuals 12
 direct venting B207
 disabilities B26,B234
 disabilities, coping with physical B148
 disabilities, learning 12,A52,B33,B51,B73,B110,B113,
 B118,B127,B149,B150,B167,B214
 disabilities, mental B39,B133,B156
 disabilities, people with mental B39
 disabilities, sensorymotor B108
 disabilities, severe mental B219
 disability B47,B59,B122,B128,B179
 disability days B24
 disability, individuals with a B95
 disability, learning B160
 disability, long-term A69
 disability, mental B78
 disability, mental and physical B234
 disability, permanence of the B122
 disability, permanent A29,A50,B89
 disability, physical A79
 disabled A50,B103,B211
 disabled child B122
 disabled children B151
 disabled persons, non-ambulant B103
 disabled, developmentally A43,B234
 disabled, employment for the A18,B37
 disabled, mentally A70

- disabled, physically A80
 disabling professions B96
 disadvantaged B138
 disadvantages, social B182
 discharge of toxic substances B222
 discharges, spark B102
 discipline techniques B147
 disciplines, relevant B91
 disclosure, product B33
 discomfort B6, B173, B203
 discomfort, minor B47
 discord, family A46, B173
 discrimination 3, 6, A7, A21, A38, A40, A47, A79, B2, B41, B86, B138, B167, B192, B218
 discrimination against lesbians and gay men B41
 discrimination against women B26, B97
 discrimination errors, choice B7
 discrimination in the private sector B41
 discrimination procedures, drug B172
 discrimination, age B25, B112
 discrimination, anti-gay B41
 discrimination, overt B55
 discrimination, racial B44, B109, B233
 discrimination, sex A33
 discrimination, social B139
 discriminative effects, pentobarbital-like B172
 discriminatory behaviour 15, A80
 discriminatory conduct A38, B79
 disease B96
 disease mimicry B5
 disease patients, infectious B119
 disease prevention A67, A77, B46, B119, B213
 disease susceptibility A64, B40
 disease susceptibility, worker B129
 disease, absence of B72
 disease, alzheimer's B5
 disease, bacterial B214
 disease, behcet's A58
 disease, cardiovascular A53, A70, B67, B137, B171, B174
 disease, cause of B142
 disease, cerebrovascular B108
 disease, chronic B120
 disease, coronary B22
 disease, coronary heart B22, B44
 disease, degenerative B143
 disease, diet and cardiovascular B142
 disease, genetic B37
 disease, heart A58, B22, B100, B112, B137
 disease, industrial A15, B185
 disease, inflammatory B168, B170
 disease, legionnaires A58, B21, B178, B236
 disease, lung B197
 disease, maternal B131
 disease, meniere's A75, B169
 disease, mental B94
 disease, occupational A75, B241
 disease, pulmonary A53, B174, B187, B188
 disease, raynaud's A56, B171
 disease, respiratory B51, B67, B109, B158, B159, B173, B231, B241
 disease, sexually transmitted B131
 disease, susceptibility to heart B76
 disease, twentieth century B216
 disease, work-relatedness of B241
 diseases B91
 diseases in animals B189
 diseases, allergic B170
 diseases, cardiovascular B152, B171
 diseases, chronic B229
 diseases, communicable A32, B209
 diseases, environmentally induced B243
 diseases, hypersensitivity A45
 diseases, iatrogenic B136
 diseases, infectious B112
 diseases, inflammatory A75, B169
 diseases, inflammatory cardiovascular A56, B171
 diseases, inflammatory type A55
 diseases, lung B233
 diseases, occupational A13, B28, B36, B135
 diseases, psychosomatic B152
 diseases, respiratory A20, A63, B41, B80, B90, B119, B185
 diseases, respiratory tract B202
 diseases, sexually transmitted 13, B2, B173
 diseases, sexually-transmitted A47
 diseases, work-related A75
 disfunction, minimal brain B167
 disgust B55
 disinfectants B243
 disinfection B178
 disintegration, community B61
 disinterest in learning B160
 dislocation A69, A71
 disorder, mental B61
 disorders, aged cerebrovascular B164
 disorders, awareness of iatrogenic B157
 disorders, chemically-induced personality B9
 disorders, childhood behavior B71
 disorders, chronic mental B216
 disorders, civil B109
 disorders, gastrointestinal B5
 disorders, genetic B108
 disorders, mental B22, B152, B229
 disorders, neurological B23
 disorders, neuropsychiatric B10
 disorders, nonspecific neuropsychiatric B10
 disorders, occupational respiratory allergic B159
 disorders, organic brain B160
 disorders, pediatric B52
 disorders, perceptual A52
 disorders, psychiatric A72, B45, B52, B72
 disorders, respiratory A51, B143
 disorders, respiratory allergic A13
 disorders, somatic and psychosomatic B152
 disorders, vision B38
 disparaging A43
 disparities B239
 disparities, socioeconomic B137
 disparity, female/male income 12
 dispersal, odour B165
 display terminals, video A16, A32, A62, B126, B128, B206
 disregard by medical personnel B122
 disruption, abrupt B18
 dissatisfaction, job B44
 distance B86
 distance from the highway B143
 distance runner B49
 distance runners A20
 distances, long B21
 distension, gaseous B50
 distracted A61
 distractibility B167
 distraction A65, B66
 distractions B222
 distress B47
 distress, cardiovascular 12
 distress, emotional B39
 distress, psychological A69, B47, B66
 distress, spiritual B34
 distressed profile B39
 distribution B28
 distribution of property A47, B2
 distribution workers B28
 distribution, air B18
 distribution, humidity B165
 distributions of work A71, B149
 distrust of adults, profound B74
 distrust, co-resident B213
 disturbance of coordination B7
 disturbances A6
 disturbances in adolescents, health B151

- disturbances, cerebral B106
 disturbances, neurological B159
 disturbances, psychological A66
 disturbances, speech A52
 disturbances, visual A9,B87,B106
 disturbed, emotionally A43,B160,B234
 disulfide, carbon B235
 diversification, physical B233
 diversity 3,6,7,8,11,15,17,19,A1,A19,A35,A48,A79,A81,
 A84,A85,B26,B27,B58
 diversity as an environment issue B26
 diversity of needs A78
 diversity, cultural A38,B44
 diversity, revealing B26
 divorce B33,B78
 divorced A33,B122
 dizziness A6,A9,A13,B9,B22,B38,B74,B75,B128,B153,
 B160,B180,B221
 doctoral students B53
 dodecyl sulfate, sodium B115
 domain, boys B84
 domains B202
 domestic accidents B114
 domestic air pollution B93
 domestic air pollution sampler B93
 domestic crises A66
 domestic life, responsibility for B183
 domestic stair construction B114
 domestic stair design B114
 domestic violence B147
 domestic water taps B21
 domestic work A34,B183
 domesticity B183
 dominance, professional B97
 dominant groups B27
 dominant theory B142
 dominate, drive to B25
 dominated peoples B2
 dominating group B2
 domination, survival of B2
 doors B128
 dopiness A8,B166
 dosage levels B144
 dose, low B14
 dose-response curve B45
 dose-response relation B140
 dose-response relationship B10
 dose-response relationships B65,B120,B124,B174
 doses of toxic substances, low B15
 double B86
 double standard B176
 double-blind A55,A83
 down's syndrome B131
 downturns, economic B140
 dioxide, nitrogen B155
 dried detergent residue B115
 drinking B76
 drinking and driving B130
 drinking drivers A25
 drinking water 5,A58,B27,B115,B201,B236
 drinking water contamination B200
 drinking, heavy B192
 drinking, social B235
 drinks, soft A58
 drive to dominate B25
 drivers, drinking A25
 drivers, young B130
 driving B130
 driving, drinking and B130
 driving, risky B61
 drop-outs B89
 droplet nuclei B178
 drowsiness A16,B131,B195
 drug B152
 drug abuse A46,B173
 drug abuse, illicit B192
 drug administration, food and B168
 drug discrimination procedures B172
 drug reactions, adverse B86
 drug residue A57,B161
 drug therapy, iatrogenic B3
 drug use, illicit B131
 drug, illicit A24
 drug, medicinal 12
 drugs B3,B38,B192
 drugs, effects of B22
 drugs, illicit A66,B135
 drugs, labelling of B86
 drugs, prescription A15
 drugs, psychotropic B22
 dry environment, hot and B100
 dry mucous membranes B140
 dry throat A12,B115
 dry-cleaning fumes A32
 dryers, hair A62
 dryness in the nose B6
 dryness, skin B131
 ductwork, air conditioning B21
 dump, toxic waste A39,B181
 dumping A57
 dumping, pesticide B215
 duplicating machines B81
 duplicating, spirit B81
 duplicators, spirit A9,B164
 durable press treatments B14
 durable-press treatments A55
 during childhood, behaviour problems B134
 dust B28,B76,B115,B187,B200,B215
 dust abatement B46
 dust control B48
 dust mite B48
 dust mite, house B186
 dust mites, house B186
 dust suppressant B223
 dust, automobile traffic A22,B217
 dust, chalk A8,B199
 dust, detergent B115
 dust, house B5
 dust, inert B5
 dust-free room B242
 dusts A13
 dusts, mineral A14
 dusts, organic B73
 dwellers, urban A19,A66,B184
 dwelling B114
 dwelling, self-contained B32
 dwelling, use of the B113
 dwellings B11,B146,B181,B243
 dyeline copying A61,B62
 dyes A32,A55,B14,B60,B165,B209
 dying A42,B189
 dysfunction syndrome, minimal brain B113
 dysfunction, hippocampal B159
 dysfunction, inner ear A13
 dysfunction, minimal brain B33,B167,B229
 dysfunction, nervous system B51
 dysfunction, vascular B171
 dysfunction, vestibular B9
 dysfunctions, perceptual motor A52
 ear B169,B170
 ear and hearing problems B29
 ear dysfunction, inner A13
 early age B86
 early warning sign A74
 earners, principal family B47
 eastern Canada B70
 ecologic management B125
 ecological classrooms B113
 ecological illness B95,B125,B149,B160
 ecological irritant B130

- ecological paradigm B37
ecological pathways B60
ecological perspective B106
ecology B12,B60
ecology classroom A11
ecology of health B142
ecology, clinical B18,B27,B53
ecology, human B17,B166
ecology, marriage of economy and B236
economic A79,B8,B37,B116,B156
economic base, community's B37
economic change B140
economic changes A69,B112
economic circumstances B213
economic considerations A15,B185
economic constraints B37
economic crisis, world B138
economic development B120
economic downturns B140
economic exploitation A32,B13
economic factors B142
economic foundations B148
economic future B142
economic goals A69
economic handicaps B25
economic hardship B34
economic incentive B134
economic levels B194
economic reversals B138
economic role of women B218
economic spinoffs 18
economic status B35,B140
economic status, women's B33
economic stress 9,B149
economics B26
economy B37,B112
economy and ecology, marriage of B236
economy, community B140
ecosystem A22,A56,B119,B168
ecosystem, urban B230
eczema A15,B157
edema, laryngeal A75,B169
edra B27
educated B8
education A8,B26,B35,B41,B44,B45,B75,B84,B116,B162,
B198
education of adults B22
education program, health B53
education system A10,B216
education, board of B113
education, community A28,B162
education, environmental B120
education, health B53,B239
education, level of B137
education, lighting for B239
education, medical A78,B204
education, poverty and B46
education, public 19,B216
education, road safety B202
education, safety A50
education, sex B100
education, special B110
education, waterloo county board of B216
educational activity B222
educational background A65
educational environment B66
educational facilities B235
educational institutions B50,B91
educational intervention B53
educational measures B193
educational policies, equity-oriented B239
educational practices B235
educational programs B204
educational system A27
educators A74,B142
educators, professional B222
eeg, computerized B214
effect, suntan B84
effect, synergistic B60,B191
effective consultation B44
effective ventilation A17
effectiveness of filtration B21
effects in tissues, thermal B17
effects of airborne bacteria, health B214
effects of density B12
effects of drugs B22
effects of pollution, health B77
effects of unemployment B112
effects, acute health B156
effects, behavioral B38,B149,B181
effects, behavioural B59,B111,B198
effects, biochemical B158
effects, biological B16,B57,B60,B102
effects, chronic B90
effects, clinical and sub-clinical B111
effects, delayed B235
effects, developmental B217
effects, electromagnetic B154
effects, health B5,B6,B14,B15,B23,B29,B40,B42,B49,
B53,B55,B60,B61,B65,B77,B79,B85,B92,B98,B100,B101,
B104,B105,B107,B108,B111,B112,B115,B117,B118,B119,
B121,B129,B131,B132,B133,B135,B136,B137,B139,B140,
B143,B146,B149,B151,B152,B155,B158,B160,B161,B166,
B167,B175,B177,B180,B184,B186,B187,B191,B192,B193,
B194,B195,B196,B197,B203,B210,B214,B215,B223,B224,
B225,B226,B228,B230,B231,B234,B237
effects, interaction B123
effects, interactive B43,B232
effects, mental health B68
effects, metabolic B158
effects, nervous system B7
effects, neuro-behavioural B179
effects, neurological B59
effects, neuropsychologic B145
effects, non-visual B84
effects, nonthreshold B168
effects, pentobarbital-like discriminative B172
effects, performance 15
effects, physiological B149,B232
effects, psychological B42,B74,B123
effects, pulmonary B191
effects, respiratory B197
effects, side B186
effects, synergistic 7
effects, teratogenic B20,B238
effects, threshold B168
effects, unhealthy side B220
efficacy B144
efficacy of lifestyle change B76
efficiency, energy A17,B126,B137,B172,B184
efficiency, medium B165
efficiency, removal B21
efficiency, work B119
efficient building, energy B92
efficient buildings, energy B89
efficient, energy B19
effluent, pulp and paper mill B208
effort, physical B44
efforts, health-damaging coping B61
egocentric thinking A50,B209
elderly 13,A5,A28,A40,A50,A72,A80,B5,B22,B30,B32,B35,
B38,B42,B49,B56,B84,B87,B90,B106,B111,B120,B122,B128,
B148,B164,B166,B198,B204,B227,B232,B238
elderly applicants B35
elderly men B87
elderly poor B194
elderly women B87
elderly, congregate housing for B111
elderly, design for the B56
elderly, frail B148

- elderly, health of the B238
elderly, homes for the A66
elderly, housing for the B32
elected officials B188
electric field B15
electric field, low-frequency B102
electric fields A62,B28,B132
electric fields, alternating B102
electric stoves B202
electrical activity, brain's A82,B108
electrical and electronic engineers B186
electrical fire A50
electrical potentials B230
electrical power transmission A61,B28
electrical transmission B153
electricity A66
electricity transmission facilities B132
electro-magnetic fields B132
electro-plating A13
electrocardiograph recordings, programmable ambulatory B191
electrocution B4
electroencephalograms, abnormal B160
electroencephalography B23,B191
electromagnetic signals B17
electromagnetic effects B154
electromagnetic energy, frequencies of B200
electromagnetic energy, man-made B17
electromagnetic environment B200
electromagnetic environment, abnormal B16
electromagnetic exposure B186
electromagnetic field B59,B102,B186
electromagnetic fields A16,B15,B16,B28,B59,B123,B185, B186,B200
electromagnetic fields, low frequency B153
electromagnetic fields, low-frequency A61,B15
electromagnetic fields, nonthermal B17
electromagnetic radiation B16,B28,B97,B102,B132
electromagnetic spectrum A60
electromagnetically sensitive A62,B171
electromagnetism B16
electroneuromyography B191
electronic air cleaners B226
electronic engineers B186
electronic engineers, electrical and B186
electronic products B63
electrophysiology, brain A83
electrophysiology, quantitative B108
elementary school B158,B177
elementary school children B222
elementary school students A60
elementary schools B242
elements, trace B200
elephant, blind men and the A1
elevated blood pressure B152
elevated diastolic blood pressure B137
elevated particulate levels B94
elevated serum cholesterol level B137
elf B59,B153
elf fields B59
eliminating prejudice 15
elimination of sexism B13
elitist design, insensitive and B1
embryo A24,B131
embryonic A52
emergencies A42
emergency A50,B78
emergency evacuations of buildings B103
emerging technologies 3
emission B100
emission of metals, atmospheric B241
emission rate B128
emission sources, indoor 5
emission, formaldehyde B5
emission, low frequency B128
emissions B70,B106,B137
emissions from products B75
emissions, automobile B116,B205
emissions, automotive B36
emissions, brake B101
emissions, diesel exhaust B117
emissions, industrial A8,B199
emissions, organic compound B137
emissions, particulate B73
emissions, sulphur dioxide B27
emissions, waste incineration A21
emitters, x-ray B20
emotional abuse A26,A33,B13,B34,B173
emotional and physical health B101
emotional climate B211
emotional distress B39
emotional factors A42,B147
emotional functioning B119
emotional independence B35
emotional lability B64,B160
emotional reactions B90
emotional resources B33
emotional stability A64,B54
emotional state B1,B39
emotional states B54
emotional stress B39
emotional suffering B79
emotional support A65
emotional toll A42,A69,B147
emotional well-being B47
emotionally disturbed A43,B160,B234
emotions A35,A65,B32
empirical basis B130
employed B47
employee B37,B93
employee health complaints B131
employee rights A18
employees A16,B18,B34,B115,B176,B206
employer B93
employers B34,B185
employment A66,B34,B35,B39,B112
employment crisis B148
employment for the disabled A18,B37
employment income B62
employment opportunities B85
employment patterns A69,B34
employment status B80
employment, full A69,A71,B149,B164
employment, non-discrimination in B139
employment, universal B148
empowering A81
empowerment 17,B90
empty vessels B162
encephalopathy A13,B191
encephalopathy, toxic B9
endemic social problem B193
endings, trigeminal nerve B11
endocrine B17
endotoxins B104,B214
endurance A35,B114
energy B116
energy conservation A5,A19,A77,B16,B19,B28,B57,B66, B91,B92,B174,B194,B197,B211,B219,B242
energy conserving community B71
energy efficiency A17,B126,B137,B172,B184
energy efficient B19
energy efficient building B92
energy efficient buildings B89
energy management B239
energy saving B11
energy sources B237
energy use A67,B119
energy, alternative sources of B237
energy, atomic B90
energy, frequencies of electromagnetic B200

- energy, man-made electromagnetic B17
energy, nuclear B90,B237
energy, renewable B237
energy-efficient A74,B177
energy-efficient houses B220
energy-related toxicants B238
enforcement officials, law A38
engine exhaust B7
engine exhaust, diesel B158
engine, gasoline A21
engineering methods, modern B231
engineering practices B238
engineers A74
engineers, electrical and electronic B186
engineers, electronic B186
english, learning B44
enhanced susceptibility to bacterial infection B32
enriched pension entitlements B149
ensuing suffering B79
enterprises B222
entitlements, enriched pension B149
entrance, university B46
enumerable A47
envelope, tightly-sealed building B199
environment B3,B57,B60,B66,B74,B82,B85,B98,B106,B118,
B119,B120,B140,B148,B161,B163,B167,B172,B178,B190,
B202,B213,B229,B236,B237,B239
environment issue, diversity as an B26
environment survey, work B206
environment, abnormal electromagnetic B16
environment, acoustic B231
environment, acoustic B230
environment, adaptive B1
environment, built B85
environment, capacity of the B222
environment, chemically-less-contaminated B171
environment, child's B71
environment, common social B176
environment, control in the physical B65
environment, educational B66
environment, electromagnetic B200
environment, eyes to the outside B222,
environment, family B211
environment, health and built B85
environment, health and the B237
environment, healthy working B13
environment, home B36,B37,B111,B167,B175,B182
environment, hostile B18
environment, hot and dry B100
environment, hot and wet B100
environment, human B12
environment, indoor B4,B62,B227,B238
environment, natural 3,B21
environment, office B69
environment, oppressive A46
environment, optimum B98
environment, outdoor B21,B84
environment, particle-free A83,B168
environment, physical B9,B36,B58,B84,B107,B140,B148,
B233
environment, proximal individual social B176
environment, residential B231
environment, school B84
environment, social B8,B98,B107,B176
environment, therapeutic B65,B157
environment, typical office B127
environment, underground B49
environment, unfamiliar B65
environment, urban B12,B36,B58,B109,B119,B230
environment, urban built B107
environment, work B66,B83,B121,B176
environment, working B37
environment-behaviour interactions B21
environmental B126,B161,B193
environmental action B9
environmental actions B9
environmental agents B223
environmental allergens B124
environmental allergens, isolation from B124
environmental care units B190
environmental chamber B219
environmental characteristics B106
environmental chemicals B229
environmental choice 3,17,A79
environmental cleanup B94
environmental conditions B206,B232
environmental conflicts B9
environmental consequences B222
environmental constraints B37
environmental control A76,A81,A83,B68,B71,B170
environmental control, comprehensive B166
environmental controls, rigid B170
environmental crises B58
environmental crisis B163
environmental cues, familiar B65
environmental damage B70,B236
environmental decisions B9
environmental design B1,B26,B36,B42,B69,B85,B100,
B111,B118,B122,B163,B234
environmental education B120
environmental evaluation B243
environmental factors B46,B74,B75,B112,B164
environmental hazard law B188
environmental hazards B23,B53,B71,B188
environmental health B30,B82,B237
environmental health alliance B161
environmental health policies B237
environmental health sciences B229
environmental health, data base on B237
environmental health, experimental 10
environmental hydrocarbons B186
environmental illness B18,B29,B54,B180,B188,B189,
B190
environmental impact B28,B223
environmental impact assessment B141
environmental improvements B69
environmental influences B3
environmental irritants B171
environmental laws B188
environmental maladaptation B160
environmental medicine B27,B50,B67,B72,B94,B127,
B171
environmental movement B61
environmental overload B169
environmental perspectives B144
environmental pollutants, toxicity of B126,B192
environmental pollution B126,B131,B134,B136,B161,
B190,B204,B232
environmental problems, prevention of B71
environmental programs B199
environmental protection B71
environmental protection act B94
environmental psychology B65,B107,B233
environmental relationships, people B235
environmental requirements, special B239
environmental research B26
environmental resources A1
environmental security B184
environmental sensitivity A44,B27,B117,B170
environmental settings B163
environmental skills B202
environmental stimulation A64,A66,A75,B107,B210,B233,
B234
environmental strain B17
environmental stresses B45
environmental stressor B40
environmental stressors B31,B40,B66,B108,B141,B183,
B210
environmental survey B131
environmental toxicity B94

- environmental toxicology B60
 environmental toxins B103,B214
 environmental variables B98,B140
 environmental, remedial B128
 environmental/social connection B9
 environmentally induced diseases B243
 environmentally induced illness B180
 environmentally susceptible individuals B58
 environmentally-induced damage B171
 environmentally-sensitive people A22
 environmentally-triggered immunopathology B3
 environments B16,B57,B68,B84,B182
 environments, adaptive B1
 environments, alternative A83
 environments, alternative social B157
 environments, children's B235
 environments, chronic care A29
 environments, confined B53
 environments, dehumanizing B44
 environments, design of hospital B233
 environments, habitable urban B230
 environments, healthy B85,B130
 environments, high density B182
 environments, home B4,B9,B113,B167,B168,B213,B230
 environments, indoor B182,B222,B239
 environments, industrial A13
 environments, institutional B65
 environments, job A65,B66
 environments, learning B235
 environments, less-contaminated A44
 environments, living B227
 environments, lower density B182
 environments, man-made B222
 environments, mine B49
 environments, mixed-use residential B183
 environments, modern job B193
 environments, natural 5,A22,B156
 environments, office B90
 environments, office work B207
 environments, polluted indoor A74
 environments, public 5
 environments, safe B83
 environments, safer B147
 environments, school B149
 environments, self-tailored A1
 environments, small town B68
 environments, social B229
 environments, stress-inducing B183
 environments, stress-reducing A68,B183
 environments, suburban B12
 environments, therapeutic B65
 environments, urban B12,B107,B229
 environments, urban physical B164
 environments, windowless B85
 environments, work B39,B170
 environments, working B222
 enzyme adaptation B2
 enzymes, nasal B48
 epidemic B144,B177,B193
 epidemics B12,B157
 epidemiologic evidence B192
 epidemiologic studies B192
 epidemiologic study B203
 epidemiological 8
 epidemiological comparisons A83
 epidemiological evaluations B123
 epidemiological studies B65,B102
 epidemiological study B61
 epidemiology B15,B80,B177
 epidemiology of fungi B184
 epidemiology, human B201
 epigenetic carcinogen B83
 epilepsy B108
 epoxy B6
 epp report, the B133
 epstein-barr virus A24,B75
 equal access B213
 equal opportunity A32
 equal pay A32
 equal relationships A35
 equal status for women 3
 equality B33
 equality between men and women B13
 equality for women 19
 equality of opportunity, need for B212
 equality of rights B218
 equality, full A79
 equalization, temperature B165
 equilibrium B22
 equipment B152
 equipment, air conditioning B28
 equipment, cooking B22
 equipment, office B74
 equipment, protective B152
 equipment, sports A50,B179
 equitable access to services B8
 equitable responsibility B222
 equity B8,B63
 equity-oriented educational policies B239
 eraser, liquid B149
 ergonomic factors A75,B241
 ergonomics B128
 errors A27
 errors, choice discrimination B7
 esophagus B231
 established dementia B87
 estate brokers, real B233
 estates, housing B21
 esteem, low self B44
 ethanol A15,B127
 ethical principles B144
 ethics B100
 ethnic minority groups B44
 ethnic origin A38,A80,B218
 ethnic/racial groups B26
 ethnicity B7,B120
 ethyl alcohol B169
 ethyl ketone, methyl A26,B165
 etiologic agents B74
 etiological agents B104
 etiology B51,B164
 etiology of childhood cancer B74
 european buildings B227
 euthanasia, voluntary B189
 evacuation B103
 evacuations A42
 evacuations of buildings, emergency B103
 evaluated data B19
 evaluation, analysis and B56
 evaluation, environmental B243
 evaluations, epidemiological B123
 evaporants, chemical B220
 evaporative condensers B236
 event, aversive A67,B138
 events, external A64,B138
 events, interpersonal B230
 events, life A72,B52,B55,B66
 events, negative life B195
 events, stressful B61
 events, stressful life B195
 evidence, epidemiologic B192
 evoked potential, visual B24
 examination B53
 examination stress A10,A64
 examination, physical B128
 excess mortality, socioeconomic status and B80
 excess sunshine B46
 excessive alcohol consumption B137
 exchange, air B194
 exchange, calcium A62,B102

- excipients B53
excitability, membrane B102
excited A8
excluded sectors B25
exclusiveness, racial B183
excretion of xenobiotics, mobilization and B188
exercise A20,B76,B105,B204,B234
exercise frequency, low B76
exercise program B76
exercise, aerobic B180
exercise, intermittent B105
exhaust B172
exhaust emissions, diesel B117
exhaust fans, kitchen B165
exhaust fans, noisy B165
exhaust hardware, add-on B49
exhaust of pollutants B199
exhaust, auto B97
exhaust, automobile A8,A20,B33,B137,B143,B167,B199,
B200,B224
exhaust, automotive B83
exhaust, car A74,B169
exhaust, catalytic gasoline B32
exhaust, diesel A14,B32,B48,B49,B117,B118,B223
exhaust, diesel engine B158
exhaust, engine B7
exhaust, spillage of A6
exhausted B138
exhaustion A35,B114,B131
exile A44
exiled people B133
exotic pets B213
expectancy, life A53,B8,B61,B88
expectancy, loss of life B174
expectations B37,B196
expectations for health care, public B63
expectations, unrealistic B195
expendable A35,B99
experience, illness B7
experience, traumatic B125
experience, user B95
experience, work B26
experiences, high-density B182
experiences, real-life A10
experiential learning B162
experiment, peckham B157
experimental animals B126
experimental classroom B97
experimental data B91
experimental environmental health 10
experiments, animal B201
expertise, exportable 18
experts, psychiatric B130
experts, ventilating A74
experts, world A79,A85
expiratory flow, peak B41
expiratory volume, forced B6,B110,B202
exploitation A35,B34
exploitation of a child A28
exploitation, economic A32,B13
exploitation, physical B114
exploring differences B26
export markets A85
exportable expertise 18
exposure guidelines B43
exposure guidelines, residential 13
exposure levels, radiation B20
exposure limit guidelines B43
exposure measurements B129
exposure monitoring, personal B155
exposure rates B174
exposure reduction B6
exposure standards, pollutant B30
exposure to heat B100
exposure, chemical B150
exposure, chronic B72
exposure, chronic occupational B69
exposure, conditions of B154
exposure, cumulative A6,B227
exposure, electromagnetic B186
exposure, fibrous glass B147
exposure, human B83
exposure, lead B159
exposure, long-term B158
exposure, low-level carbon monoxide B7
exposure, massive chemical B125
exposure, noise B230
exposure, occupational B87,B185,B186
exposure, outdoor/indoor B147
exposure, parent's workplace B165
exposure, passive B200
exposure, periods of B232
exposure, personal B147
exposure, pesticide B192
exposure, prenatal B64,B75
exposure, solvent B10
exposure, styrene B11,B123
exposure, toxic chemical B109
exposure, toxic substance B108
exposure, worker B185
exposure, workplace B5
exposure-response relationships B210
exposures to particles, personal B156
exposures, chemical B73,B171
exposures, chronic low-level B49
exposures, cumulative B166
exposures, multiple B29
exposures, outdoor B227
exposures, parents B165
exposures, pesticide A62
exposures, radiation B63
exposures, workplace B177
expression, outlet for B9
exterior spaces B213
external events A64,B138
external load factors B29
externalization of blame B112
extra-parental care B25
extreme aggression B151
extremely low frequency B15
extremely low-frequency B102
eye irritation A8,A19,B131
eyes to the outside environment B222
eyes, burning A6
eyes, pupils of the A60,B230
eyestrain B240
fabric A55
fabric filter A7,B165
fabric, cotton B191
fabric, synthetic B191
fabrics B14,B97
fabrics, treated A5
fabrics, urea formaldehyde in B83
facilities B182
facilities, chemical manufacturing B193
facilities, child-care A38,B44
facilities, educational B235
facilities, electricity transmission B132
facilities, health care A75,B192
facilities, medical A19,B210
facilities, medical care B113
facilities, play B65
facilities, psychiatric B233
facilities, sewage treatment B21
facility, psychiatric B234
factors, age B164
factors, behavioral B22,B61
factors, biological B98
factors, cognitive B76
factors, constitutional B37

- factors, economic B142
factors, emotional A42,B147
factors, environmental B46,B74,B75,B112,B164
factors, ergonomic A75,B241
factors, external load B29
factors, genetic B74
factors, interdependent B235
factors, personality B54,B230
factors, physical B206
factors, political B119
factors, predisposing A40
factors, regional geological A6
factors, risk B30,B71,B137,B233
factors, socioeconomic B80,B202
factors, stress B232
factors, thermal B238
factory smokestack B176
factory, sulphuric acid A38,B180,B208
faculties of medicine B192
faculties of medicine, Canadian B204
failure to provide B34
faint-dizziness A69
faint-dizziness, spells of B47
fall injuries, slip and B211
fallacy of typology B89
falling accidents A29,B157,B197,B211,B227
falls A29,A41,B4,B83,B86,B211
falls in buildings B198
falls, slips and B59
false arrest A38,B64
false ideas about youth A27
familial turbulence B45
familiar environmental cues B65
families B7,B54,B147,B209
families in crisis A78
families, black B233
families, immigrant B7
families, low-income A70
families, middle-income-earning A70
families, non-smoker B107
families, single-parent A36
families, singleparent A36
families, two-parent A36
families, wellbeing of B149
family B37,B39,B58,B66,B71,B86,B95,B119,B147,B153,
B211
family breakdown B112
family conflict A65,B66
family control B66
family discord A46,B173
family earners, principal B47
family environment B211
family health B79
family history B146
family income B24,B101
family independence B66
family law A47,B41
family laws B2
family life B148,B218
family medicine B134
family members B152
family participation A75,B65
family planning A78,B116,B152
family problems 5
family responsibilities A34,A35,A36,B45,B219
family structure B26
family therapy B156
family type 3,A2
family types A36
family violence B101
family, burden of B34
family, context of the B212
family-centered health care B54
familycentered health care A78
fan coil B82
fan coil units A54
fan, continuous low-speed B165
fan, low-speed A7
fans, kitchen exhaust B165
fans, noisy exhaust B165
farm machinery A50,B179
farmers, studies of B192
fat-stored foreign compounds B188
fatalism B112
fatalities B52
father B76
father, gay A47
fatherhood B45
fatherhood revival B45
fathers A35
fathers, gay B66
fatigue A13,B9,B50,B103,B127
fatigue syndrome, allergy tension B160
fatigue, chronic A6
fatigue, nervous A10
fatter A70,B218
favorable attitudes B35
fear B2,B55,B90
fear of aging 13
fear of atomic war B190
fear of crime A66,B147
fears B86
feat, intellectual A27
federal/provincial advisory committee A5
feedback B43
feeling of competence B85
feelings B195
feelings of guilt B33
feelings, unresolved B195
feingold diet B127
female B13,B22,B164
female labour force B183
female managers B50
female patients B136
female/male income disparity 12
females A69,B47,B88
feminine role B143
feminine suburbs B183
fenitrothion B116
fertility B38
fertilizer residue B72
fetal pathology B101
fetus A52,B18,B131
fetus, sensitivity of the B64
fetuses A5,A40,B5,B101
fever, humidifier B82
fiberglass A15
fiberglass, irritation from B177
fibers B141
fibers, synthetic A53,B14,B120,B141
fibers, textile A55
fibreboard B159
fibres, synthetic B211
fibres, textile B14
fibrosing alveolitis B124
fibrosis, cystic A40,B172
fibrous glass exposure B147
field investigations B25
field studies B227
field, electric B15
field, electromagnetic B59,B102,B186
field, low-frequency electric B102
field, magnetic A61,A62,B102
fields of medicine B135
fields, alternating electric B102
fields, electric A62,B28,B132
fields, electro-magnetic B132
fields, electromagnetic A16,B15,B16,B28,B59,B123,
B185,B186,B200
fields, elf B59

- fields, low frequency electromagnetic B153
 fields, low-frequency electromagnetic A61,B15
 fields, magnetic B132
 fields, nonthermal electromagnetic B17
 fields, power frequency A61
 fields, surface B102
 fields, vlf B128
 filling, dental amalgam B94
 filter, fabric A7,B165
 filters A58,B226
 filters, air B21,B113
 filtration B19,B199
 filtration, effectiveness of B21
 financial difficulties A69
 financial resources B33
 financial resources, personal B122
 financial rewards B134
 financial stress 5
 findings, pathological B187
 finishings, interior 15
 fire B103
 fire, electrical A50
 fired cooking stoves, gas B73
 fireplaces A20
 fishing, sport B70
 fitness in the workplace, mental B93
 fitness program, community B76
 fitness promotion, mental B86
 Fitness Survey, Canada B137
 fixed incomes A30,B232
 fixed monitoring stations B174
 fixtures, radiation-shielded fluorescent A61
 flame ionization B227
 flame retardants A55,B14
 flat roofs A9,A54,B76
 flexibility A81,B37
 flexibility, work time B45
 flexible parttime work B93
 flexible work-time A36
 flights, airline B217
 floor construction B106
 floor underlay, particleboard B106
 floor wash, pine-scented B169
 floors B178
 floors, bare B165
 flow rate, nasal inspiratory B186
 flow, peak expiratory B41
 flu symptoms, cold and B103
 fluctuations, temperature B103
 flue-like B50
 fluids, bodily A48
 flunking B45
 fluorescent B154
 fluorescent fixtures, radiation-shielded A61
 fluorescent light ballasts A54,B124
 fluorescent lighting B62,B239
 fluorescent lights A61,B97,B126,B132,B154
 fluorescent tubes A9
 fluorescent tubes, full-spectrum A10
 fluoridator B225
 fluoride B115,B116,B225
 fluorine B115
 fluorocarbons B116
 fluorithyl B172
 flushing B180
 flux density, magnetic B15
 fly ash B200
 foam B25
 foam insulation B14
 foam insulation, urea formaldehyde A6,B14,B38,B51,
 B197
 foam insulation, urea-formaldehyde B20,B31,B79,B209
 foams, urea-formaldehyde B19
 focus, community B63
 focus, technology B224
 fetus A24,B228
 food 5,A56,B90,B105,B149,B168,B240
 food additives A43,A56,A57,A76,B46,B48,B60,B73,B88,
 B113,B136,B138,B187,B190,B200,B229,B240
 food allergies A76,B127,B150,B160,B167,B178,B187
 food allergy B171
 food and drug administration B168
 food containers, polyethylene B105
 food cravings A77,B29
 food diets, junk B18
 Food Guide, Canada's A56
 food hypersensitivity B130
 food ingredients, hidden B138
 food intolerance B50
 food labels, incomplete B138
 food or chemical sensitivities B168
 food poisoning B5
 food production A67,B119
 food selection guidelines B142
 food sensitivities B48,B111
 food sensitivity B150
 food supply B215
 food triggering agents B171
 food, nutritious A2,B161
 food, reactions to B171
 food, restaurant A57,B240
 foods B108,B170
 foods, labeling of processed B168
 foods, processed A57
 foods, refined B189
 footprint A82
 for, human rights issues B2
 force, female labour B183
 force, labour B33,B137,B196
 force, paid labour B33
 force, social B116
 forced air heating system B165
 forced expiratory volume B6,B110,B202
 forced vital capacity B6
 forces B69
 forces, societal B4
 foreign compounds, fat-stored B188
 forensic B130
 forest products B70
 forgotten people B130
 form, physical B69
 formal health care B156
 formal service systems B156
 formaldehyde A6,A8,A13,A15,A19,A25,A32,A53,A55,A74,
 B5,B6,B14,B16,B19,B20,B23,B25,B27,B43,B46,B48,B57,
 B69,B70,B78,B83,B87,B91,B92,B97,B106,B113,B120,B128,
 B136,B141,B142,B146,B151,B152,B154,B157,B159,B162,
 B169,B180,B184,B197,B203,B210,B211,B219,B221,B225,
 B228
 formaldehyde emission B5
 formaldehyde foam insulation, urea A6,B14,B38,B51,
 B197
 formaldehyde in fabrics, urea B63
 formaldehyde, hypersensitized to B83
 formaldehyde, sensitized to B6
 formaldehyde, urea A6,B25,B106
 formaldehyde-based resin binders B106
 formaldehyde-induced cancer B83
 formaldehyde-sensitive A8
 formaldehyde-sensitive individuals B69
 formation, identity B173
 formic acid B162
 fossil fuels B202,B226,B237
 foundations, economic B148
 fraction, particulate B123
 fractures, bone B30
 fragmentation B135
 frail A66,B151
 frail elderly B148
 framework B95

- framework for support B39
framework, organizational B71
framework, policy B34
franchisers B25
freckling, heavy B62
free loans, interest B149
freedom A35,B84
freedom from intergroup conflicts B164
freedom of assembly B139
freedom of choice A65,B163
freedom of press B139
freedom of speech B139
freedom, professional B142
freedoms A47
freedoms, charter of rights and B218
freedoms, democratic A66,B164
freeways, crowded A65
freeze-thaw B25
frenchspeaking quebec B64
freon B120
freons B27
frequencies B16,B28
frequencies of electromagnetic energy B200
frequencies, power B102
frequency electromagnetic fields, low B153
frequency emission, low B128
frequency fields, power A61
frequency range, power A62
frequency response, windows of B200
frequency, extremely low B15
frequency, low exercise B76
frequent conflicts B183
frequently changing shift B28
fresh air A11,B158,B172,B212
fresh air supply B18
friable asbestos B26
frictions, interpersonal A31,B35
friendliness, less B184
friends A72,B76,B95,B216
friendship B68
friendships B36
from aids, mortality 13
from environmental allergens, isolation B124
from fiberglass, irritation B177
from intergroup conflicts, freedom B164
from molds, toxins B52
from numbers, decisions B201
from products, emissions B75
from public attention, deleted B58
from soil, gases A6
from the highway, distance B143
front B213
fruitful stimulus B222
fruitgrowing regions A58,B72
fruits A57
frustration A27,B150,B152,B216
frustrations of hypersensitivity B81
fuel prices B226
fuel, gas cooking B55
fuels B229
fuels, fossil B202,B226,B237
full employment A69,A71,B149,B164
full equality A79
full spectrum A61,B154
full-spectrum fluorescent tubes A10
full-spectrum ultraviolet light B154
full-time home-maker B39
fume, metal oxide B123
fume-free A55,A83,B168
fumehoods B172
fumes, chemical B125
fumes, cooking B55
fumes, cutting B123
fumes, dry-cleaning A32
fumes, metal oxide A13
fumigatus, aspergillus B224
function tests, psychological B83
function tests, respiratory B204
function, brain A8,B108,B170,B198
function, brain synaptic A62,B102
function, immune A64
function, impaired ventilatory B87
function, liver B11
function, lung A40,B42,B124
function, mental B15,B160
function, pulmonary B5,B11,B109,B120,B129,B204,B212
function, reduction in lung B94
function, respiratory B93,B105,B210
functional competence A66
functional requirements, child's B122
functioning women B211
functioning, emotional B119
functioning, humoral immune B54
functioning, immune A10,A64
functioning, optimal B72
functioning, phagocytic immune B53
functioning, psychological B71
functions, cardiovascular B225
functions, defensive B56
functions, psychological B123
functions, pulmonary B121
funding B142
funeral services A47,B55
fungal contamination B184
fungal growth A6,A54,B25
fungal infection A57,B189
fungal infections B58
fungal levels B82
fungal spores B31,B103,B187
fungal volatiles B215
fungi A14,B29,B51,B58,B73,B103,B184,B189,B215
fungi production B103
fungi, epidemiology of B184
fungi, macrocyclic trichothecene-producing B103
fungi, plant B58
fungi, toxigenic A77,B187
fungus A42
fungus growth B31
furnishings 5,B22,B70,B74
furniture A32,B106
furniture, built-in B106
furniture, poorly designed B208
fusarium toxins A57,B189
futility B44
future B236
future of mankind B13
future, commitment to the B63
future, economic B142
future-oriented vision B50
futurist B57
gain, weight A77,B29
gallup poll B218
gamma rays A62
gaps in services B122
garages B155
garages, indoor A16
garbage incinerator A52,B161
garden B165
garden pesticides A26
garments B191
garments, cotton B191
gas analysis B91
gas appliances B73,B86,B219
gas backpacking stove B41
gas chromatographic study B56
gas chromatography B117,B243
gas chromatography, high-resolution A82
gas chromatography-mass spectrometry A82
gas chromatography/mass spectrometry B227
gas cooking appliances B108

- gas cooking fuel B55
gas cooking stoves B155
gas fired cooking stoves B73
gas heater B147
gas space heaters, unvented natural B73
gas stoves A5,B20,B54,B93,B94,B101,B121,B202
gas toxicology B5
gas, natural B169
gas, tracer B155
gas-chromatograph mass-spectrometer B89
gas-fired kitchen appliances B74
gas-fired range B219
gaseous distension B50
gases from soil A6
gases, anaesthetic A32,B209
gases, organic B11
gases, soil B199
gasoline additives B116
gasoline engine A21
gasoline exhaust, catalytic B32
gasoline vapours B124
gasoline, leaded B116
gastric secretions A60,B230
gastro-intestinal problems B160
gastrointestinal difficulties B29
gastrointestinal disorders B5
gastrointestinal system B97
gastrointestinal tract A60,B230
gay A35,A36,A46,A80,B81,B136
gay and lesbian liberation B112
gay couples A47
gay father A47
gay fathers B86
gay liberation B2
gay men A48,B41
gay men, discrimination against lesbians and B41
gay oppression 17,A46,A49,B112
gay people, aging A30
gay people, invisibility of B40
gay people, older B81
gay rights A47,B41,B139
gay sex B4
gay-identified peers A46,B173
gay-supportive housing A48
gays and lesbians aging B81
gays, stereotyped images of B40
gcms B89
gender 3,6,8,A2,A46,A80,B40,B112,B211
gender difference A33,B211
gender differences B40,B88
gender-preference A48
general hospital treatment B216
general physician A74
general public B8
generalist, declining influence of the B135
generalized malaise B103
genetic D74,B221
genetic background B53
genetic disease B37
genetic disorders B108
genetic factors B74
genetic predispositions A68,B37
genetic substrate B157
genetically determined susceptibility B75
genetics B172
genetics, stress B40,B61
genotoxic B80
genuine consideration B231
geographic differences B74
geographic isolation B138
geographic segregation B182
geography B74
geological factors, regional A6
geology, regional B181
geriatric assessment units B56
germs B37
gestation B131
ghandis A1
ghettoization B216
girls B21,B84
glass exposure, fibrous B147
global war A35
glue, urea-formaldehyde A53,B5
glues B6
goals, economic A69
goals, personal B93
good mental health B99
government B34,B90,B112
government administrators B185
government office complex, three-tower B131
government policies B134
government policy B133,B135,B241
government standards B121
governmental lobbying B216
governments B212,B236
grab bars A29
grab sample B141
grade, below B199
graduate students A64,B232
gram negative bacilli B210
graphs B19,B91
grass roots A21
gratification, immediate B90
gratuitous interference B100
great lakes pollution A58,B99
grievances, redress of B139
grieving process B112
ground B89
ground water B19
group activities B234
group needs B1
group social action B213
group, blood sulfhydryl A63
group, dominating B26
group, high-risk A47
group, oppressed B26
group, social B9
groups B163,B182,B196
groups, affinity B182
groups, bottom income B8
groups, citizens B147
groups, dominant B27
groups, ethnic minority B44
groups, ethnic/racial B26
groups, high income B142
groups, high risk A41,B161
groups, high-risk B238
groups, higher income A71
groups, immigrant B44
groups, interest B205
groups, manageable-size B182
groups, minority A38,B64,B233
groups, non-dominant B27
groups, older age B130
groups, risk B124
groups, social A30,B148
groups, socio-economic B213
groups, socioeconomic B88,B137
groups, target B8,B137
groups, vulnerable population B239
growing without schooling B92
growth B101
growth conditions B103
growth, fungal A6,A54,B25
growth, fungus B31
growth, microbial A9,A54,B76
growth, mould A77,B199
growth, population B116
guaranteed upper limit B138
guidance B141,B156

- guide B47
 Guide, Canada's Food A56
 guide, consumer B47
 guidelines B135
 guidelines, customs B4
 guidelines, dietary B142
 guidelines, exposure B43
 guidelines, exposure limit B43
 guidelines, food selection B142
 guidelines, residential exposure 13
 guilt A69,B184
 guilt, feelings of B33
 guns A50,B179
 gynecology B136
 Oxhealth, brain's state of B108
 habitable urban environments B230
 habitat B53
 habits, dangerous A4
 habits, maternal B131
 habits, poorer health B243
 hacking cough A12
 hair analysis B213
 hair dryers A62
 hair spray, aerosol A5,B70
 hairdressers A32,B208
 half-life A6
 hallucinations A43,B103
 halogen A51
 halothane B172
 hand immersion B100
 hand smoke, second B42,B125
 handbook of prevention B59
 handicap 3,A2,A40,B128
 handicapped 6,13,17,A41,A42,A80,B58,B238
 handicapped adults, mentally B153
 handicapped children B122
 handicapped people 6
 handicaps B109,B131
 handicaps, economic B25
 handling, material B152
 handrails A29,A42,B10,B211
 handwriting B127
 happenings, outside B222
 harassment B41
 harassment, sexual A34,B35
 hardship, economic B34
 hardware, add-on exhaust B49
 harm to children B70
 harmful chemical substances B215
 harassment, sexual 12
 hate B41
 hate crimes B67
 hate-motivated violence A46,B67
 hay-fever A9
 hazard B193
 hazard law, environmental B188
 hazard, health B5,B69,B83,B172,B217
 hazardous B209
 hazardous substances B152
 hazardous waste management B223
 hazards B193,B208,B243
 hazards, biological A75,B88,B241
 hazards, chemical B152
 hazards, delayed health A58,B192
 hazards, environmental B23,B53,B71,B188
 hazards, health B3,B6,B12,B16,B18,B19,B21,B22,B24,
 B27,B38,B41,B44,B47,B48,B49,B50,B51,B52,B58,B67,B70,
 B72,B73,B76,B81,B82,B84,B88,B90,B91,B92,B94,B97,B103,
 B104,B105,B106,B108,B109,B110,B120,B123,B124,B125,
 B126,B127,B128,B135,B141,B144,B146,B147,B149,B151,
 B152,B153,B157,B158,B159,B161,B165,B166,B168,B175,
 B176,B178,B179,B180,B184,B186,B187,B188,B189,B190,
 B192,B196,B197,B198,B200,B201,B206,B207,B209,B215,
 B216,B217,B220,B221,B223,B225,B227,B228,B229,B233,
 B240
 hazards, household H53
 hazards, industrial A78,B134
 hazards, industrial health A75,B241
 hazards, industrially generated health B239
 hazards, occupational B78,B102,B135
 hazards, occupational health B134,B186
 hazards, office B126
 hazards, reproductive A15,B38,B193
 hazards, respiratory B123
 hazards, respiratory health B123
 hazards, safety B37
 hazards, unexpected A44
 hazards, vdt radiation B208
 hazards, vulnerability to A81
 hazards, workplace B236
 head-tilt A63
 head-trauma B108
 headache A9,A13,A19,A53,B96,B115,B131,B173,B180
 headaches A6,A9,A16,A56,A77,B29,B74,B103,B153,B160,
 B240
 headaches, vascular B171
 heads, shower A54,A59,B-1
 healing A73
 healing processes B119
 healing, processes of B57
 health admissions, mental 13
 health advocacy, mental A44,B39
 health alliance, environmental B161
 health and built environment B85
 health and safety 3,6,B33,B123
 health and safety committees, joint B185
 health and safety, child H83,H130
 health and safety, occupational A1,A34,B3,H33,B38,
 B44,B73,B123,B127,B128,B134,B135,B157,B165,B180,B185,
 B190,B191,B208,B221,B235,B241
 health and safety, school A1,A12,A81
 health and safety, tenant A1,A81
 health and safety, workplace A15,B239
 health and social policies, Canadian B142
 health and the environment B237
 health behavior B76
 health beliefs B76
 health care B85,B92,B142,B232
 health care delivery B16
 health care delivery systems 3,B156
 health care facilities A75,B192
 health care policies B54
 health care policy B7
 health care professionals B142,B195
 health care professions B134
 health care providers B63,B123
 health care services, managed B142
 health care services, nation's B91
 health care system A74,B63,B86,B123
 health care system, Canadian B224
 health care system, mental B216
 health care systems 3
 health care systems, mental A43
 health care workers B209
 health care workers, native community B96
 health care, adequate B192
 health care, alternatives to B85
 health care, barriers to B192
 health care, Canadian B90
 health care, family-centered B54
 health care, familycentered A78
 health care, formal B156
 health care, holistic B54
 health care, non-judgmental A49,B156
 health care, personal B8
 health care, preventive mental B34
 health care, public expectations for B63
 health claims B142
 health classes, academic B76
 health clinics, independent occupational B241

- health complaints B35
health complaints, employee B131
health cost B27
health data B8
health disturbances in adolescents B151
health education B53,B239
health education program B53
health effects of airborne bacteria B214
health effects of pollution B77
health effects, acute B156
health effects, mental B68
Healthy For All A2,B83
Healthy For All, achieving B133,B134,B204
Healthy For All: implications, achieving B142
health habits, poorer B243
health hazards, delayed A58,B192
health hazards, industrial A75,B241
health hazards, industrially-generated B239
health hazards, occupational B134,B186
health hazards, respiratory B123
health impact statements B57
health in the city B229
health information system B129
health information systems B129
health insurance, universal B8
health issue, preventable B145
health issues, women's B122
health management counselling B204
health monitoring devices 15
health movement, holistic A77,B82
health needs B8,B204
health of Indians B191
health of the elderly B238
health officials B240
health order, new world B117
health path, soft B82
health planning B8,B118
health planning, long term B118
health policies, environmental B237
health policies, public B138
health policy B83,B103,B116,B142
health policy, national B134
health policy, occupational B185
health policy, public B22,B37,B57,B60,B83,B96,B100,
B120,B122,B204,B224,B229
health practices, positive mental B34
health problems B37,B47
health problems, mental B101,B140,B191
health problems, occupational B73
health problems, public B179
health professionals B35,B69,B243
health professionals, mental B141,B216
health promotion A30,A67,A77,A78,B57,B63,B96,B119,
B133,B134,B204,B213
health promotion programs B86,B137
health promotion strategies B204
health protection B201
health publications, mental B51
health research B228
health researchers B109
health risks B70,B220,B221
health science centres B91
health sciences, environmental B229
health services B47,B86
health services, mental A33,B95
health social worker B54
health status B142
health status of children B138
health status, indicator of B8
health status, lower B8
health strategies B8,B133
health strategy, international B239
Healthy Strategy, Premier's Council on B64
health surveillance B129
health survey data B143
Healthy Survey, Canada B46,B137
health survey, national B179
health system, mental B95
health variables B112
health variables, community mental B140
health, behaviours that promote B204
health, child B138
health, child's mental B71
health, children's B183
health, children's mental B183
health, community B36,B109
health, community mental B140
health, community-based public B82
health, complete A1
health, damages to B12
health, data base on environmental B237
health, design for B85
health, diet and B142
health, ecology of B142
health, emotional and physical B101
health, environmental B30,B82,B237
health, experimental environmental 10
health, family B79
health, good mental B99
health, housing and B239
health, human B70,B220,B230
health, maintenance of B57
health, mental A17,A67,A69,A79,B8,B12,B21,B31,B33,
B34,B35,B39,B44,B45,B68,B74,B79,B90,B93,B95,B97,B98,
B108,B112,B119,B133,B138,B140,B151,B152,B156,B162,
B163,B176,B182,B183,B184,B189,B195,B196,B197,B207,
B210,B211,B216,B218,B219,B232,B233
health, occupational B42,B69,B72,B87,B121,B125,B126,
B129,B152,B161,B174,B177,B185,B211,B236,B240
health, occupational safety and B192
health, personal B149
health, poor B80
health, poor mental B88
health, poor physical B44,B88
health, promoting B117
health, psychological B34
health, public B28,B37,B57,B71,B101,B103,B120,B136,
B173,B202,B238
health, respiratory B24,B73,B77,B92,B107,B116,B184
health, social nature of B83
health, socioeconomic causes of ill B133
health, somatic B32
health, unemployment and B46
health, unemployment and physical B112
health, urbanization and human B12
health, women's B35
health, women's mental A34
health-damaging coping efforts B61
health-related problems of women B239
healthful behavior A73,B76
healthfulness of life B8
healthier lifestyle B47
healthsharing B133
healthy city B58
healthy environments B85,B130
healthy lifestyles A78,B96,B229
healthy living B81
healthy populations B232
healthy public policy B83
healthy society B148
healthy working environment B13
hearing A60,B77,B230
hearing impaired child B153
hearing loss, high-frequency A60
hearing losses A29,B56
hearing problems A77
hearing problems, ear and B29
heart attacks, coronary B44
heart disease A58,B22,B100,B112,B137
heart disease, coronary B22,B44

-
- heart disease, susceptibility to B76
 - heart rate A53,B10,B146
 - heart trouble A69,B47
 - heat A63
 - heat, exposure to B100
 - heat, release of B222
 - heater, gas B147
 - heater, unvented space B148
 - heaters B91
 - heaters, kerosene A5
 - heaters, space B91,B219
 - heaters, unvented kerosene space B73
 - heaters, unvented natural gas space B73
 - heating B21,B48,B220
 - heating system, forced air B165
 - heating systems B147
 - heating systems, home A6
 - heating systems, household A20
 - heating, home A51
 - heating, space B194
 - heavy B122
 - heavy drinking B192
 - heavy freckling B62
 - heavy industry B241
 - heavy metal A20,A25
 - heavy metals A13,A40,A43,B126,B157,B161,B200,B208,
B229
 - heavy workload B209
 - height A52
 - helplessness B97
 - helplessness, human B138
 - hematological B17
 - hemoglobin B154
 - hemoglobin concentration B83
 - hemorrhage B38
 - herbicides A20,B53
 - herbicides, phenoxy B192,B193
 - hermetically sealed B206
 - hermetically sealed office complex B18
 - heroin addicts A19,B231
 - heterosexual A48
 - heterosexually-oriented A46
 - hexachloride, benzene A58
 - hexachloro-1,3-butadiene B136
 - hidden A49,A84
 - hidden food ingredients B138
 - hidden program B69
 - hidden sensitivities B125
 - hiding A47,B2
 - hiding differences A79
 - high blood pressure A69,B47,B189
 - high cholesterol levels B44
 - high density B182
 - high density apartments B183
 - high density environments B182
 - high income groups B142
 - high injury rates B114
 - high morbidity B191
 - high protein-low carbohydrate diet B39
 - high rise living B183
 - high risk groups A41,B161
 - high school students B105
 - high tension B44
 - high voltage line B153
 - high-cost technology B91
 - high-crime A65,B213
 - high-density experiences B182
 - high-energy places A21,A64,B95
 - high-frequency hearing loss A60
 - high-frequency radiation A62
 - high-resolution gas chromatography A82
 - high-rise buildings A22,B21,B183
 - high-risk A26,A29
 - high-risk group A47
 - high-risk groups B238
 - high-risk subsample B243
 - high-school students B24
 - high-voltage substation B15
 - high-voltage transmission lines A62
 - higher admission rates B152
 - higher density B183
 - higher income groups A71
 - higher income levels B8
 - higher-exposure homes B59
 - highly specialized personnel B213
 - highrise A41
 - highway A26
 - highway, distance from the B143
 - highway, main B107
 - highway-accident B197
 - hippocampal dysfunction B159
 - hippocampus A25
 - hispanics B67
 - histories, clinical B170
 - histories, residential B122
 - history, family B146
 - history, medical B128
 - history, occupational A75
 - hoarseness A12
 - hockey B7
 - holistic approach B86
 - holistic health care B54
 - holistic health movement A77,B82
 - holistic view B82
 - home A62,B4,B33,B53,B122
 - home accidents B37,B161,B228
 - home appliances A62
 - home care services B7
 - home delivery B131
 - home environment B36,B37,B111,B167,B175,B182
 - home environments B4,B9,B113,B167,B168,B213,B230
 - home heating A51
 - home heating systems A6
 - home office B37
 - home responsibilities, burden of B209
 - home safety B53,B114
 - home safetyproofing B53
 - home, accident in the A4
 - home, adapting the B122
 - home, independence in the B81
 - home, past B111
 - home, pesticide-free B171
 - home, sense of B32
 - home, women's work in the B33
 - home, work outside the B183
 - home-based work B37
 - home-based work, computer-mediated B39
 - home-based workers A18,B39
 - home-maker, full-time B39
 - home-owners B20
 - home-schooling B50
 - homeless A7,H78
 - homeless, plight of the B78
 - homelessness B78
 - homelike B112
 - homeostatic mechanism B169
 - homeostatic mechanisms B170
 - homeowners B217
 - homes 3,5,B74,B89,B94,B121,B146,B148,B219
 - homes for the elderly A66
 - homes, airtight B27
 - homes, Canadian B132
 - homes, higher-exposure B59
 - homes, lower-exposure B59
 - homes, materials for B215
 - homes, mobile A6,B27,B128
 - homes, nursing A33,B122,B166
 - homes, private B220
 - homes, sealing of B215
 - homes, temporal aspects of B230

-
- homes, light A6
 - homogeneity, social class B233
 - homogenization A84,B27
 - homophobia B40,B55
 - homophobic attitudes A47
 - homophobic prejudice B41
 - homosexual A47,B201
 - homosexual adolescent A46
 - homosexual adolescents A49,B156
 - homosexual lifestyles B16
 - homosexual youths B173
 - homosexuality B2,B4,B40,B41,B55,B67,B112,B156,B173
 - homosexuality, adolescent B173
 - homosexuality, overt prejudice against A49
 - homosexuality, prejudices against B2
 - homosexuality, societal biases towards B16
 - homosexuality, stereotypes of B56
 - homosexuals A46,B86
 - honesty in communication A68
 - hood, unvented B165
 - hopelessness 7,10,17,A45,B81,B152
 - hormonal B18
 - hormonal regulatory mechanisms B2
 - hospital B65,B79,B157,B233
 - hospital admission B15,B192
 - hospital admission rates A69
 - hospital admission rates, mental B112
 - hospital admissions 12,A38
 - hospital buildings A16,B207
 - hospital environments, design of B233
 - hospital social relationships B233
 - hospital treatment, general B216
 - hospital, children's B65
 - hospitals A45,A59,A75,B54,B95,B222,B227,B237
 - hospitals, mental B152
 - hospitals, psychiatric B22
 - hospitals, teaching B91
 - host susceptibility B104
 - hostile environment B18
 - hot and dry environment B100
 - hot and wet environment B100
 - hot water system B165
 - hot water systems A54,B21
 - hotels A59,B237
 - hour reductions, annual job B149
 - hours, inconvenient A17,B44
 - house dust B5
 - house dust mite B186
 - house dust mites B186
 - house paint B89
 - house painters B83
 - house, idealized A42,B122
 - house, stigmatized B122
 - house, stigmatizing the B122
 - household B33
 - household accidents A4
 - household comforts B33
 - household hazards B53
 - household heating systems A20
 - household incomes, below-average B68
 - household injuries B52
 - household injuries, childhood B52,B53
 - household products B220
 - household responsibilities B34
 - household technologies B165
 - household type A37
 - household work loads B45
 - households A36,B51
 - households, mother-led A36,B194
 - households, non-smoking A26
 - households, single-parent B194
 - households, two-parent B39
 - houses B28,B43,B74,B86,B88,B101,B140
 - houses, energy-efficient B220
 - houses, stone B11
 - housewife A36
 - housewife, myth of the B183
 - housewives A34,B93,B148
 - housewives, symptomatic B93
 - housework A32,B13,B196
 - housing 17,B12,B32,B35,B62,B66,B71,B78,B122,B177,B227,B238,B243
 - housing alternatives B42
 - housing and health B239
 - housing costs A71,B149
 - housing crisis, rental B62
 - housing design B42,B165,B199,B206,B226
 - housing estates B21
 - housing for elderly, congregate B111
 - housing for the elderly B32
 - housing services B4
 - housing shortage B78
 - housing shortages A36
 - housing, affordable B78,B232
 - housing, appropriate B95
 - housing, congregate A31,B111
 - housing, gay-supportive A48
 - housing, improvements in B36
 - housing, inadequate B216
 - housing, intergenerational B232
 - housing, low-income A7
 - housing, low-income B78
 - housing, new B35
 - housing, permanent B78
 - housing, poor B229
 - housing, public B232
 - housing, rental B233
 - housing, specialized B199
 - housing, subsidized A36,B63
 - housing, supply of B78
 - hum B239
 - human B92,B117,B172
 - human behavior B71,B113,B167,B213,B230
 - human cancer B174
 - human characteristics A1
 - human community B37
 - human context B93
 - human costs B112
 - human costs of unemployment B112
 - human development B131
 - human dignity B163
 - human ecology B17,B166
 - human environment B12
 - human epidemiology B201
 - human exposure B83
 - human health B70,B220,B230
 - human health, urbanization and B12
 - human helplessness B138
 - human immune system B28
 - human metabolism 5
 - human microorganism B220
 - human need B9,B99
 - human needs B163
 - human needs, basic B97
 - human properties B163
 - human reactions B11
 - human relationship, pathology of B25
 - human rights 17,A22,B97,B218
 - Human Rights Code, Ontario B41
 - human rights codes A47,B112
 - human rights issues for B2
 - human rights legislation A79,A80
 - human rights policy B97
 - human rights, basic B222
 - human scale B119,B229
 - human senses A55
 - human sensory irritation B227
 - human service needs, changing B224
 - human smoking B101
 - human subjects B144

-
- humidifier fever B82
 - humidifier A14,B73
 - humidifiers A6,A54,B21,B214
 - humidifiers, cool-mist B224
 - humidifiers, spray B28
 - humidity A16,A17,B77,B131,B206,B210,B220,B238
 - humidity distribution B165
 - humidity, relative A63
 - humiliated A27,B50
 - humiliations A26,B34
 - humoral A64
 - humoral immune functioning B54
 - hungry B25
 - husbandry, animal B239
 - husbands A36,B45
 - hydrocarbon pesticides, chlorinated A82,B117
 - hydrocarbon solvents, volatile A82
 - hydrocarbon, chlorinated B136
 - hydrocarbon, polycyclic aromatic A14
 - hydrocarbons B50,B141,B191
 - hydrocarbons, aromatic B137
 - hydrocarbons, chlorinated B136,B170,B232
 - hydrocarbons, environmental B186
 - hydrocarbons, isoparaffinic B227
 - hydrocarbons, polycyclic B211
 - hydrocarbons, polycyclic aromatic A83,B52,B117
 - hydrogen cyanide B121
 - hydrogen sulfide A8,B69
 - hydrol, paratoluene sulfinic acid of michler's A16
 - hydrolysis A6,B25
 - hydroxide, sodium A8,B69
 - hygiene products, perfumed personal A11
 - hygiene, industrial B137,B242
 - hygiene, personal B173
 - hygienic B152
 - hyoergy B232
 - hyper-sensitivity, delayed-type B14
 - hyperactive behaviour A61
 - hyperactive children A9
 - hyperactivity A43,A52,B127,B132,B167,B239
 - hyperirritability B229
 - hyperkinesia B154
 - hyperlipidemia B22
 - hyperreactivity A51,B196
 - hyperresponsiveness, airway B234
 - hypersensitive 9,B175,B239
 - hypersensitive individuals A44
 - hypersensitive, chemically 6
 - hypersensitivity B50,B67,B113,B125,B170,B175,B216,B238
 - hypersensitivity diseases A45
 - hypersensitivity pneumonitis B177
 - hypersensitivity to non-physiologic chemicals B193
 - hypersensitivity, chemical A55,A82,B81
 - hypersensitivity, food B130
 - hypersensitivity, frustrations of B81
 - hypersensitized to formaldehyde B83
 - hypersusceptibility A41,A76,B45,B113,B138,B161,B208
 - hypersusceptibility, chemical A17,B172
 - hypersusceptibility, nutritional B173
 - hypertension A69,B22,B47,B146,B174,B189
 - hypertensive B146
 - hypocalcemia B30
 - hypochondriacs B138
 - hypochondriasis A74,B170
 - hyoergy B232
 - hypotension B38
 - hypotension, orthostatic B22
 - hypothalamus, cat B186
 - hypoxia A63,B100
 - hysterectomies B136
 - hysteria, mass A17,B64,B138
 - hysterics B138
 - iatrogenic diseases B136
 - iatrogenic disorders, awareness of B157
 - iatrogenic drug therapy B3
 - ice arenas A19
 - ice resurfacing machines B7
 - ice rink H7
 - ice-resurfacing machines A19
 - idealized house A42,B122
 - ideas about youth, false A27
 - ideas, polarized B183
 - identification, pollution source B128
 - identification, public A47
 - identity A30,A67,B9,B42
 - identity formation B173
 - ideology B7
 - idiosyncrasy B45
 - idiosyncratic nature of individual responses B232
 - ignorance A45,A49,A80,B40,B185
 - ill child, chronically B7
 - ill health, socioeconomic causes of B133
 - ill persons, mentally B51
 - ill, chronically mentally B207
 - ill, mentally 17
 - ill-treatment of a child B34
 - illicit drug A24
 - illicit drug abuse B192
 - illicit drug use B131
 - illicit drugs A66,B135
 - illness B37,B143,B233
 - illness among women B143
 - illness experience B7
 - illness prevention B204
 - illness, adaptation to B147
 - illness, building B178,B206,B207
 - illness, building-related A9,B172
 - illness, chronic B122
 - illness, customs for managing B7
 - illness, ecological B95,B125,B149,B160
 - illness, environmental B18,B29,B54,B180,B188,B189,B190
 - illness, environmentally induced B180
 - illness, mass psychogenic B138
 - illness, mental 13,A43,B51,B98,B103,B129,B143,B195
 - illness, physical B47
 - illness, portrayal of mental B51
 - illness, prevention of mental B98
 - illness, psychiatric B190
 - illness, psychosomatic B81
 - illness, respiratory B86,B108
 - illness, sex differences in B143
 - illness, stigma of mental A43,A70,B112
 - illness, vulnerabilities to B149
 - illnesses B37
 - illnesses, mimicry of common B5
 - illnesses, smoking-related B145
 - illnesses, stress-related B34
 - illumination B239
 - image of psychiatry, public B129
 - image, male A35
 - image, superman A35
 - images of gays, stereotyped B40
 - images, normative B27
 - images, stereotyped A24,A46
 - images, stereotypical A49
 - imaging, brain B108
 - imaging, computerized B108
 - imbalance in work loads B45
 - imbalance, metabolic B39
 - imbalanced workload A36
 - immediate gratification B90
 - immersion, hand B100
 - immigrant A39
 - immigrant chinese B7
 - immigrant families B7
 - immigrant groups B44
 - immigrants B26,B151
 - immune abnormalities B171

- immune alteration B52
- immune deficiency syndrome, acquired A47
- immune function A64
- immune functioning A10,A64
- immune functioning, humoral B54
- immune functioning, phagocytic B53
- immune parameters B170
- immune response A64,B40
- immune suppression A83
- immune system A52,A56,A58,A77,B3,B17,B18,B28,B40,B53,
B54,B94,B95,U130,B171,B187
- immune system suppression B58
- immune system, human B28
- immune systems, suppressed A42
- immunity indicators A25,B160
- immunity, cell-mediated B232
- immunity, local B159
- immunity, mammalian B28
- immunization H96,B177,B178
- immunized B100
- immuno-toxic B47
- immuno-toxicity U215
- immunodeficiency B3
- immunogen B14
- immunoglobulin levels A64,B54
- immunoglobulins B68
- immunologic responses B18
- immunological responsiveness, decreased B232
- immunology 9,A83
- immunopathology B3
- immunopathology, environmentally-triggered B3
- immunosuppressive A54,B52
- immunosuppressive agents B103
- immunotoxicity B52
- immunotoxicology B52
- impact assessment B16,B23,B221
- impact assessment, environmental B141
- impact statements, health B57
- impact, environmental B28,B223
- impacts, social B34
- impaired ability to concentrate B227
- impaired child, hearing B153
- impaired children, learning B110
- impaired coordination B87
- impaired motor coordination B23
- impaired ventilatory function B87
- impairment, cognitive B87
- impairment, intellectual B10
- impairment, long-term B64
- impairment, memory B9,B11,B72
- impairments, motor A25
- impasse B111
- impedance, nasal B30
- implications, achieving health for all B142
- imported A57
- imposed setting B167
- impoverished B61
- improvements in housing B36
- improvements, environmental B69
- inability to concentrate B50
- inability to cope B167
- inability to learn B22
- inability to participate B206
- inability to work B58
- inactivity A70
- inactivity, physical B137
- inadequate housing B216
- inadequate information B44,B98
- inadequate ventilation B90
- incapacitating B125
- incapacity of children B144
- incense A26,B165
- incentive, economic B134
- incentives, work B112
- incest A26,B34
- incidence on stairs, accident B114
- incidence, cancer B74
- incineration emissions, waste A21
- incineration, municipal A51
- incineration, waste B126
- incinerator, garbage B161
- incinerators A52,B83,B241
- incinerators, garbage A52
- incinerators, municipal A20
- incinerators, sewage sludge B126
- incitant challenges A83,B169
- income A71,B76,B80,B95,B158
- income alternatives 17
- income children, low B183
- income credits B149
- income disparity, female/male 12
- income groups, bottom B8
- income groups, high B142
- income groups, higher A71
- income levels, higher B8
- income security B85
- income, employment B62
- income, family B24,B101
- income, maintenance B149
- incomes, below-average household B68
- incomes, fixed A30,B232
- incomes, lowest B78
- incomes, poverty-level A36,B194
- incomplete food labels B138
- inconvenient hours A17,B44
- increased blood pressure B44
- increased risk of death B80
- increased status A65
- incubation B236
- independence A30,A31,A43,A65,A66,A75,A79,B95,B106,
B233
- independence in the home B81
- independence, emotional B35
- independence, family B66
- independent contractors A18,B39
- independent learning B162
- independent occupational health clinics B241
- independent reflection B162
- Indian reserve B180
- Indian reserve, serpent river B180,B208
- Indians, health of B191
- Indians, native B191
- Indians, Ontario B180
- Indians, Union of Ontario B180
- indicator B223
- indicator of health status B8
- indicators 12
- indicators of susceptibility B45
- indicators, behaviour 13
- indicators, immunity A25,B160
- indicators, performance 13
- indicators, social H36,B97
- indices of stress B206
- indices, respiratory B41
- indirect responsibility B204
- individual B37
- individual choice B142
- individual control 3,B172
- individual differences B61,B209
- individual needs B106
- individual resistance B37
- individual responses, idiosyncratic nature of B232
- individual responsibility B133
- individual self-care A68
- individual sensitivities B237
- individual social environment, proximal B176
- individual susceptibility B53,B149
- individuality B1,B212
- individualized curative approach B213
- individuals with a disability B95

- individuals, alienated B68
 individuals, chemically susceptible A11,B212
 individuals, direct contact with 12
 individuals, environmentally susceptible B58
 individuals, formaldehyde-sensitive B69
 individuals, hypersensitive A44
 individuals, mold sensitive B29
 individuals, needs of B196
 individuals, older unemployed B47
 individuals, sensitive 10,13
 individuals, susceptible A76,B47,B48,B64
 individuals, unemployed B47
 indoor B197
 indoor air A53,B215
 indoor air pollution 5,B5,B6,B11,B13,B14,B15,B16,B17,
 B19,B21,B22,B27,B28,B29,B35,B38,B42,B43,B46,B47,B48,
 B53,B54,B55,B57,B62,B65,B67,B69,B73,B74,B78,B79,B82,
 B88,B89,B90,B91,B92,B97,B101,B108,B109,B113,B115,
 B120,B123,B124,B126,B127,B128,B135,B136,B137,B139,
 B140,B141,B142,B145,B147,B149,B150,B152,B155,B158,
 B164,B165,B166,B167,B173,B174,B175,B176,B177,B178,
 B181,B184,B186,B188,B189,B190,B194,B196,B197,B198,
 B202,B203,B206,B207,B211,B212,B217,B219,B221,B222,
 B225,B226,B230,B236,B242
 indoor air quality 3,B6,B18,B21,B51,B56,B66,B73,
 B76,B79,B92,B96,B103,B104,B110,B123,B125,B131,B146,
 B153,B159,B165,B166,B172,B180,B184,B187,B189,B197,
 B199,B201,B205,B206,B207,B209,B214,B215,B220,B226,
 B227,B236,B237,B238
 indoor air, bacteria in B214
 indoor air, of B19
 indoor air, polluted 3
 indoor airborne viruses B238
 indoor climate B238
 indoor climate complaints B76
 indoor combustion appliances B73,B219
 indoor emission sources 5
 indoor environment B4,B62,B227,B238
 indoor environments B182,B222,B239
 indoor environments, polluted A74
 indoor garages A16
 indoor microflora B215
 indoor paneling B152
 indoor parking B155
 indoor pollution B210
 indoor pollution, air B86
 indoor surfaces B202
 indoor-outdoor relationships B54
 induced diseases, environmentally B243
 induced illness, environmentally B180
 industrial B231
 industrial activity A51
 industrial chemicals B3,B15,B45
 industrial competition A85
 industrial compounds B157
 industrial disease A15,B185
 industrial emissions A8,B199
 industrial environments A13
 industrial hazards A78,B134
 industrial health hazards A75,B241
 industrial hygiene B137,B242
 industrial irritants A40,B172
 industrial laminates B14
 industrial life, speed of B190
 industrial medicine B46,B53
 industrial nations A57,B136
 industrial processes 5,A52,B60,B241
 industrial settings A13
 industrial sources A20
 industrial workers A13,B15,B93
 industrialization B239
 industrialized nations B4
 industrially-generated health hazards B239
 industries, pharmaceutical B240
 industries, polluting B27
 industry B185,B240
 industry, construction B134
 industry, heavy B241
 industry, margarine B142
 industry, pharmaceutical B240
 industry, textile B67
 ineffective treatment B7
 inequalities B83,B151
 inequalities, reducing B196
 inequitable social order B2
 inequities B115
 inequities, reducing B142
 inequities, social class A78,B134
 inert dust B5
 infancy B158
 infant A52,B18,B66,B131
 infant death A38,B192
 infant death syndrome, sudden B5
 infant death, sudden A40
 infant development B131
 infantile behaviour, regression to B90
 infantile jaundice A61,B84
 infants A40,A53,B5,B131,B166
 infants, young B198
 infection A9,A48,A52,B21,B58,B73,B177,B178,B236
 infection of tissues B215
 infection, airborne B178
 infection, childhood B172
 infection, enhanced susceptibility to bacterial B32
 infection, fungal A57,B189
 infection, nitrogen dioxide B60
 infection, opportunistic B187
 infection, recurrent respiratory B143
 infection, resistance to B17,B60
 infection, respiratory B60
 infection, susceptibility to bacterial A14
 infections B51
 infections, fungal B58
 infections, opportunistic A77
 infections, respiratory B104
 infections, respiratory tract B212
 infections, respiratory virus B178
 infections, viral B104
 infectious aerosol B32
 infectious agents A24,B75
 infectious disease patients B119
 infectious diseases B112
 infectiousness B177
 infectivity B104
 inferior A24,B50
 inferiorization B2
 inferiorized 6,A79
 infiltration A6,B124,B219,B225
 infiltration rates B20
 infiltration, air B91
 inflammatory bronchitis A13,B14
 inflammatory cardiovascular diseases A56,B171
 inflammatory disease B168,B170
 inflammatory diseases A75,B169
 inflammatory type diseases A55
 inflexible B224
 influence B117
 influence of the generalist, declining B135
 influences B74
 influences, behavioral B229
 influences, cognitive B76
 influences, environmental B3
 influences, positive A64,B95
 influenza B5
 information A81
 information about women B35
 information overload A61,A65,B183,B222
 information system, health B129
 information systems, health B129
 information, access to B144

- information, inadequate B44,B98
information, media B51
information, reproductive B129
information, restricted 13
informed choice B63
infrared analyzing B7
infrasound B123
infringement, personal B230
ingenuity 10
ingested lead A25
ingrained attitudes B50
ingredients, hidden food B138
inhalant B149
inhalants, particulate B125
inhalants, volatile B172
inhaled acids B187
inhaled air pollutants B119
inhaled microbes B187
inhaled pollutants B160
inhibition processes, cortical B24
initiatives B91
injection therapy B170
injection, intradermal B180
injuries A4,A25,A41,B52,B83,B130,B192,B211
injuries, accidental B70,B114
injuries, childhood household B52,B53
injuries, household B52
injuries, slip and fall B211
injuries, wound and B164
injurious products B179
injury A50,B34,B73,B173,B179,B185
injury consequences B135
injury control B179
injury prevention B179
injury rates, high B114
injury, personal B192
injury, physical B134,B152
injury, psychological B79
injustices A24,B50
inks, commercial B47
inner ear dysfunction A13
innovations, side-effects of A65
innovations, spread of B122
inorganic A52
inorganic compounds B42
inorganic material B187
inorganic materials B241
input, public A21
inquest A41,B211
insecticides A5,B72
insecticides, organophosphorus A58,B72
insecurity A33,B211
insecurity, job B206
insensitive and elitist design B1
inside/outside boundaries B113
insignificant A75
inspectors, provincial B212
inspiratory flow rate, nasal B186
instability B195,B200
institution, managed B133
institution, policy-making B188
institutional care B219
institutional changes A85
institutional design B56
institutional environments B65
institutional practices B7
institutional problem solving B133
institutional qualities B235
institutional setting B81
institutional settings B235
institutionalized oppression A32
institutions B95,B212,B222,B235
institutions, educational B50,B91
institutions, mental B216
institutions, psychiatric A75,B233
institutions, social B163
institutions, structure of B133
instrumentation B218
instruments, calibrated B3
instruments, measuring B142
insufficiency, vertebrobasilar B38
insulation B25,B210,B215
insulation, foam B14
insulation, thermal A54,B25,B189,B215
insulation, urea formaldehyde foam A6,B14,B38,B51,
B197
insulation, urea-formaldehyde foam B20,B31,B79,B209
insulted A27,B50
insurance A44
insurance, universal health B8
intake, alcohol A15
intake, lead A52
intake, maximum permissible dietary B72
intakes, school air A11
integrated caring neighbourhoods B119
integration B33,B233
integration, community B219
integrity A66
integrity in public life B164
intellectual B119,B183
intellectual feat A27
intellectual impairment B10
intellectual learning B92
intelligence A52,B145,B187
intelligence tests B213
intense lighting sources B62
intensities B232
interaction B183
interaction effects B123
interaction, body/mind B54
interaction, interpersonal B182
interaction, social B87,B234
interaction, uffi-fungal B51
interactions, environment-behaviour B21
interactions, nutrient-toxicant B126
interactions, personal B147
interactive behaviours, socially B234
interactive effects B43,B232
intercourse, anal A48
interdependent factors B235
interdisciplinary B91
interest free loans B149
interest groups B205
interest in reading, decreased B243
interest, lack of B89
interference, gratuitous B100
intergenerational housing B232
intergroup conflicts A66
intergroup conflicts, freedom from B164
interieur, milieu B16
interior construction B199
interior finishings 15
interiors, automobile B243
interiors, low-pollution A5
intermittent exercise B105
internal currents B102
internal pollution B16
international cooperation A22,B222
international health strategy B239
interpersonal events B230
interpersonal frictions A31,B35
interpersonal interaction B182
interprofessional cooperation B179
interrogated A27,B50
interrupting prejudice 11
intersectoral action B239
intersectoral policies B83
intervention research B36
intervention, educational B53
interventions, behavioral B22

- intimate A72
 intimidated A75,B65
 intimidation B67
 intimidation, verbal A46,B67
 intolerance, chemical B105
 intolerance, food B50
 intolerant B169
 intoxication B235
 intradermal injection B180
 invasion, privacy B118
 inventory, danish national B6
 inventory, research B136
 investigations, field B25
 investment B188
 investment, community A44
 invisibility 13,A47,A84,B27
 invisibility of gay people B40
 invisible 7,A49,A80
 invisible minorities B139
 involvement A67,A84
 involvement, community B148
 involvement, crash B130
 involving children in design B235
 involving children, research B144
 ionic motion B102
 ionization, flame B227
 ionizing B60
 ionizing radiation A62,B3,B20
 ions, air B123
 ions, negative air A8,A63,B110
 iron A52,B218
 irregularities, menstrual B152
 irreversible lung damage B73
 irritability A10,B50,B80,B127,B131,B154,B187,B221
 irritable A8,B138
 irritancy threshold A53
 irritant B97
 irritant responses B221
 irritant, ecological B130
 irritant, respiratory B14
 irritants, chemical B130,B190
 irritants, environmental B171
 irritants, industrial A40,B172
 irritants, sensory B62
 irritation A13,A53,A54,B6,B14,B18,B31,B62,B79,B131,
 B140,B173
 irritation from fiberglass B177
 irritation, eye A8,A19,B131
 irritation, human sensory B227
 irritation, mucous membrane B227
 irritation, respiratory A16,B90,B115
 irritation, threshold for B83
 irritation, throat B129
 irritation, upper airway B5
 irritation, upper respiratory B30
 iso-butane B27
 isoamyl nitrate B172
 isocyanic acid B53
 isodecane A16,B82
 isolated A33,A44,A46,A75,B65
 isolated specialization B9
 isolation 5,7,A10,A45,A49,A67,B1,B86,B134,B152,B156,
 B162
 isolation from environmental allergens B124
 isolation, geographic B138
 isolation, social A64,A78,B80,B134,B210
 isoparaffinic hydrocarbons B227
 issue, diversity as an environment B26
 issue, preventable health B145
 issues for, human rights B2
 issues, postnatal prevention B131
 issues, public policy B36,B144,B236
 issues, social B90,B163
 issues, women's health B122
 itching A12
 janitorial chemicals B167
 janitorial supplies B22
 japan B114,B216
 japanese children B216
 jaundice, infantile A61,B84
 Jewish Latina lesbians B27
 Jews A46,B2,B67
 job B34
 job availability B112
 job clarity B66
 job control, little B209
 job demands B44
 job design B185
 job dissatisfaction B44
 job environments A65,B66
 job environments, modern B193
 job hour reductions, annual B149
 job insecurity B206
 job satisfaction B240
 job satisfaction, low B44
 job sharing A18,A36,B45,B93
 job specification B44
 job stress 13,B78
 job, losing a B112
 job, loss of a B78
 job-seeking B112
 jobless B152
 jobs B149,B197
 jobs, access to decent A71
 jobs, monotonous B208
 jobs, service A32
 jobs, unstable B196
 joint health and safety committees B185
 judged A27
 judged constantly B50
 judgement B162
 junction triangle B61,B203
 junk food diets B187
 jurisdiction B243
 justice system, criminal B67,B147
 juvenile delinquency B73
 kepone B49
 kepone and mirex B116
 kerosene B219
 kerosene heaters A5,B110
 kerosene space heaters, unvented B73
 ketone B49
 ketone, methyl ethyl A26,B165
 ketones B137
 kidney B116,B231
 kidney damage A13
 kids A27,B12
 kindergarten B124
 kindergarten children A63,B77
 kindergartens A9,B76
 kitchen B82,B106
 kitchen appliances, gas-fired B74
 kitchen exhaust fans B165
 kitcheners B216
 know, right to A81
 knowledge development B144
 knowledge, children's B243
 knowledge, core body of B176
 labelling of processed foods B168
 labelling A56,A57
 labelling of drugs B86
 labelling, product B138
 labels, incomplete food B138
 labels, price B157
 labels, psychiatric A43,B95
 lability, emotional B64,B160
 laboratories, chemical B172
 laboratories, science A12,B13,B177
 laboratories, testing B75
 laboratory protocols 10

-
- life events, stressful B195
 - life expectancy A53,B8,B61,B88
 - life expectancy, loss of B174
 - life options, reduced B134
 - life setting, new B35
 - life skills A44,B216
 - life stressors B101
 - life style B125
 - life, animal A22
 - life, community B95
 - life, control over one's B
 - life, cornerstones of B57
 - life, family B148,B218
 - life, healthfulness of B8
 - life, integrity in public B164
 - life, pace of A85,B183
 - life, quality of B57,B95,B122,B203
 - life, quality of urban B9,B163
 - life, responsibility for domestic B183
 - life, social B35
 - life, speed of industrial B190
 - life, street A46,B173
 - life, stressors of city B12
 - life, suburban B12
 - life, urban A66,A72,B68,B163,B183
 - life, wheel of B25
 - life-and-death A56
 - life-threatening A54,B58,B215
 - lifelong lovers B81
 - lifestyle 3,6,A30,A46,A76,A80,B76,B137,B147,B157,
B204
 - lifestyle change, efficacy of B76
 - lifestyle changes B151
 - lifestyle modification B8
 - lifestyle, destructive B152
 - lifestyle, healthier B47
 - lifestyle, loss of customary B18
 - lifestyles A36,A39,A71,A84,B8,B46,B85
 - lifestyles, alternative 16
 - lifestyles, collective determinants of B8
 - lifestyles, healthy A78,B96,B229
 - lifestyles, homosexual B16
 - lifestyles, traditional B46
 - lifting A32,B208
 - light A60,B84,B154
 - light ballasts, fluorescent A54,B124
 - light, diffused B62
 - light, full-spectrum ultraviolet B154
 - light, ultraviolet A16,A61,B62,B82
 - light, undiffused B62
 - lighting A17,A32,B123,B239
 - lighting for education B239
 - lighting in offices, windows and B239
 - lighting sources, intense B62
 - lighting sources, unusual occupational B62
 - lighting, artificial A16,B69
 - lighting, fluorescent B62,B239
 - lighting, office B240
 - lighting, poor B206,B208
 - lights B84,B171
 - lights, fluorescent A61,B97,B126,B132,B154
 - limit guidelines, exposure B43
 - limit value, occupational threshold B227
 - limit values, threshold B3,B7
 - limit, guaranteed upper B138
 - limitations, mobility A30,B32
 - limitations, personal B193
 - limited resources B142
 - limited way of thinking B26
 - limits, safety B10
 - line, high voltage B153
 - line, poverty A32
 - linemen, power and telephone B186
 - linemen, telephone B186
 - lines, high-voltage transmission A62
 - lines, paced assembly B44
 - lines, transmission B102,B132
 - lip B231
 - liquid eraser B149
 - liquid process photocopiers B227
 - liquid scintillation B88
 - liquid wastes B60
 - liquors A58
 - literature, psychiatric B211
 - little job control B209
 - livable cities B162
 - liver function B11
 - living arrangements B86
 - living conditions B182
 - living environments B227
 - living quarters B42
 - living units B43
 - living will B189
 - living, healthy B81
 - living, high rise B183
 - living, lower standard of B112
 - living, standard of B97
 - living, suburban B233
 - load factors, external B29
 - load, stress 9
 - load, total B17,B29
 - load, total stress B18
 - loading tests, mental B63
 - loads, household work B45
 - loads, imbalance in work B45
 - loans, interest free B149
 - lobbying, governmental B216
 - local action B83
 - local immunity B159
 - local policies B243
 - local unemployment rates B140
 - locations, outdoor B213
 - locomotor activity, spontaneous B118
 - long distances B21
 - long term health planning B118
 - long-term disability A69
 - long-term exposure B158
 - long-term impairment B64
 - long-wavelength ultraviolet A9,B154
 - longevity B109
 - longitudinal study of aging, Ontario B89
 - longrange transport of airborne pollutants B70
 - losing a job B112
 - loss of a job B78
 - loss of benefits B78
 - loss of customary lifestyle B18
 - loss of life expectancy B174
 - loss of self-esteem 13,B152
 - loss, high-frequency hearing A60
 - loss, memory B23
 - losses, hearing A29,B56
 - lovers, lifelong B81
 - lovers, same-gender A30
 - low back pain, chronic A65,B66
 - low dose B14
 - low doses of toxic substances B15
 - low exercise frequency B76
 - low frequency electromagnetic fields B153
 - low frequency emission B128
 - low frequency, extremely B15
 - low income children B183
 - low job satisfaction B44
 - low levels B158
 - low levels of pollution B233
 - low nitrogen content B103
 - low pay B209
 - low pollution design B184
 - low self confidence B44
 - low self esteem B44
 - low self-esteem A33,B211

- low self-worth B95
low socio-economic status B24
low socioeconomic status B192
low-frequency electric field B102
low-frequency electromagnetic fields A61,B15
low-frequency, extremely B102
low-income A30,A65,B68
low-income families A70
low-income housing A7
low-income street B213
low-income unemployed A69,B47
low-level carbon monoxide exposure B7
low-level contaminants B27
low-level exposures, chronic B49
low-level radiation A16,B69
low-pollution building design A7,B199
low-pollution buildings B198
low-pollution classroom B216
low-pollution interiors A5
low-speed fan A7
low-speed fan, continuous B165
lower density environments B182
lower health status B8
lower standard of living B112
lower-class blacks B9
lower-exposure homes B59
lowest incomes B78
lowincome housing B78
lubricants A55,B14
lung B172,B231
lung biopsy B184
lung cancer A6,A14,B11,B42,B120,B121,B132,B146,B174,
B223
lung damage B205
lung damage, irreversible B73
lung disease B197
lung diseases B233
lung function A40,B42,B124
lung function, reduction in B94
lung volume measurements B204
lungs B188
lupus B171
lymphatic cancers B132
lymphocytes B171
machine man B163
machine-paced A68
machinery, dangerous A15,B73
machinery, farm A50,B179
machines, duplicating B61
machines, ice resurfacing B7
machines, ice-resurfacing A19
machines, photocopying A16
machines, wet-process A16
macho A35
macho stereotype A35
macrocytic trichothecene toxins A54
macrocytic trichothecene-producing fungi B103
macroenvironment B140
macroenvironmental stressors B141
magazines A6,B69
magnesium A56
magnetic field A61,A62,B102
magnetic fields B132
magnetic flux density B15
main highway B107
mainstream B120
maintenance income B149
maintenance materials 3,B74,B125
maintenance of buildings B215
maintenance of health B57
maintenance products 5,A15,B137
make-up air B21
makers, policy B83
makeup air U175
maladaptation B166
maladaptation, environmental B160
maladies H103
maladjustments B71
malaise, generalized B103
male B26
male image A35
male physicians, chauvinistic B136
male stereotyping A35,B99
male workers B45
males B88,B89
males, liberation of B99
malformations, congenital B193
malfunction, brain A76,B178
malignant melanoma B62
mammalian immunity B28
mammals B28,B36
man, machine B163
man-made chemicals B16
man-made electromagnetic energy B17
man-made environments B222
man-made pollutants B37
manageable-size groups B182
managed health care services B142
managed institution B133
management counselling, health B204
management program B243
management, air quality B136
management, ecologic B125
management, energy B239
management, hazardous waste B223
management, obstetrical B131
management, waste A67,B119
manager B37
managerial B240
managers B185
managers, building A74
managers, female B50
managers, middle A15,B185
managing illness, customs for B7
mandatory A8
mandelic acid B191
manifestations of stress B190
manifestations, clinical B240
manipulation B114
manipulation therapy, dietary B49,B50
mankind, future of B13
manly A35,B114
manual carpet sweeper B165
manual labourers B34
manual workers B28
manufacture, carpet B46
manufacturers A11
manufacturers, wine B240
manufacturing B11
manufacturing facilities, chemical B193
margarine industry B142
margin of safety B228
margin, safety B228
marital status B80
market B47
market, consumer B213
market, worldwide 18
markets, export A85
markets, world 3,19
marriage B33,B101
marriage breakup A47,B86
marriage of economy and ecology B236
masculine cities B183
masking phenomenon B169
mass destruction A22,B222
mass hysteria A17,B64,B138
mass psychogenic illness B138
mass respirable particulate B65
mass spectrometry U117
mass spectroscopy B243

- mass transit B229
 mass, particle B155
 mass, respirable particulate B218
 mass-spectrometer, gas-chromatograph B89
 massive changes B134
 massive chemical exposure B125
 mastectomies, radical B136
 material handling B152
 material standards B25
 material, building B92,B137,B181
 material, construction B5
 material, inorganic B187
 material, radioactive A39,B208
 material, suspended microbiological B185
 material, synthetic building B137
 materials for homes B215
 materials, art B84,B225
 materials, art and craft A5
 materials, binding B137
 materials, building 5,15,A6,A15,A25,A54,A77,B6,B20,
 B22,B31,B62,B79,B83,B89,B97,B103,B104,B106,B137,B139,
 B140,B203,B220,B221
 materials, cleaner 3
 materials, construction B6
 materials, craft B84
 materials, inorganic B241
 materials, less-polluting A81
 materials, maintenance 3,B74,B125
 materials, packing B53
 materials, radioactive A62
 materials, science A8,B199
 materials, synthetic B113
 maternal characteristics B131
 maternal compliance B131
 maternal disease B131
 maternal habits B131
 maternal smoking B178
 mathematical models B5
 matrix A1
 matter, particulate B43,B165,B196
 matter, suspended particulate B105
 maturation B66
 maximum permissible dietary intake B72
 mbk neuropathy B127
 meaning, social B9
 measles B177,B178
 measles vaccine B177
 measurement B37
 measurement devices 15
 measurement of pollution B79
 measurement systems B142
 measurement techniques B136
 measurement technology, brainwave A82
 measurement, density B228
 measurements, exposure B129
 measurements, lung volume B204
 measurements, performance A83
 measures, avoidance B18
 measures, cooperative B34
 measures, educational B193
 measures, neurobehavioural B199
 measures, performance B75
 measures, preventive B123
 measures, remedial B210
 measures, specific therapeutic B142
 measuring instruments B142
 mechanical ventilation B210
 mechanically ventilated B206
 mechanically-ventilated A74
 mechanism, homeostatic B169
 mechanism, protective B144
 mechanisms, adaptive B2,B18
 mechanisms, biologic B75
 mechanisms, body defense B119
 mechanisms, coping B61,B68
 mechanisms, defence A78,B237
 mechanisms, homeostatic B170
 mechanisms, hormonal regulatory B2
 mechanisms, non-immunological B68
 mechanisms, pathogenic B74
 mechanisms, pathological defense B90
 mechanisms, respiratory tract defense B187
 media A43,B51,B71,B129,B141
 media information B51
 media, popular print B51
 mediators B232
 mediators of stress B52
 medical and law students B40
 medical assistance B23
 medical attention B52
 medical care facilities B113
 medical care, health insurance B80
 medical community B46
 medical devices B86
 medical education A78,B204
 medical facilities A19,B210
 medical history B128
 medical model, current B204
 medical personnel, disregard by B122
 medical procedures B136
 medical research B90,B91
 medical research programs B91
 medical school curricula A49,B16
 medical schools B204
 medical schools, Canadian B91
 medical science B90
 medical support A74
 medical training A78,B237
 medication, antiasthmatic B58
 medication, therapeutic B178
 medications A30,A70,B112,B204
 medicinal drug 12
 medicine B96,B136,B172
 medicine, Canadian faculties of B204
 medicine, environmental B27,B50,B67,B72,B94,B127,
 B171
 medicine, faculties of B192
 medicine, family B134
 medicine, fields of B135
 medicine, industrial B46,B53
 medicine, occupational B173,B208
 medicine, preventive A78,B204
 medicine, role of B142
 medium efficiency B165
 medium, culture B28
 melanoma, malignant B62
 mellitus, diabetes B137
 members of society, weakest B213
 members, family B152
 membrane excitability B102
 membrane irritation, mucous B227
 membrane, cell B28
 membranes, dry mucous B140
 membranes, mucous A74,B170
 memory A8,A13,A29,A45,A58,B42,B110,B160
 memory deficits B150
 memory impairment B9,B11,B72
 memory loss B23
 memory problems B15
 memory skills, auditory B150
 memory, auditory B110
 memory, changes in B56
 memory, visual A63,B110
 men B37,B38,B40,B72,B86,B99,B183,B185
 men and the elephant, blind A1
 men and women, equality between B13
 men, dehumanization of B99
 men, discrimination against lesbians and gay B41
 men, elderly B87
 men, gay A48,B41

- men, oppression of working-class B114
men, stereotyping of B114
men, suburban B182
men, urban B182
meniere's disease A75,B169
menstrual irregularities B152
mental ability A80
mental and physical disability B234
mental capacity B63
mental confusion A8
mental disabilities B39,B133,B156
mental disabilities, people with B39
mental disabilities, severe B219
mental disability B78
mental disease B94
mental disorder B61
mental disorders B22,B152,B229
mental disorders, chronic B216
mental fitness in the workplace B93
mental fitness promotion B66
mental function B15,B160
mental health A17,A67,A69,A79,B8,B12,B21,B31,B33,B34,
B35,B39,B44,B45,B68,B74,B79,B90,B93,B95,B97,B98,B108,
B112,B119,B133,B138,B140,B151,B152,B156,B162,B163,
B176,B182,B183,B184,B189,B195,B196,B197,B207,B210,
B211,B216,B218,B219,B232,B233
mental health admissions 13
mental health advocacy A44,B39
mental health care system B216
mental health care systems A43
mental health care, preventive B34
mental health effects B68
mental health practices, positive B34
mental health problems B101,B140,B191
mental health professionals B141,B216
mental health publications B51
mental health services A33,B95
mental health system B95
mental health variables, community B140
mental health, child's B71
mental health, children's B183
mental health, community B140
mental health, good B99
mental health, poor B88
mental health, women's A34
mental hospital admission rates B112
mental hospitals B152
mental illness 13,A43,B51,B98,B103,B129,B143,B195
mental illness, portrayal of B51
mental illness, prevention of B98
mental illness, stigma of A43,A70,B112
mental institutions B216
mental loading tests B63
mental performance A54,B140
mental retardation A52,B187,B243
mental stress B146
mental stress, cardiovascular reactivity to B146
mental well-being B240
mentality, sheltered-workshop B112
mentally disabled A70
mentally handicapped adults B153
mentally ill 17
mentally ill persons B51
mentally ill, chronically B207
mentally retarded B110
mercury A13,B15,B38,B49,B89,B94,B116,B157,B164,B211,
B232
mercury adsorption B202
mercury vapor B69
mercury vapour B177
message promotion B144
metabolic effects B158
metabolic imbalance B39
metabolic rate A24
metabolism B27,B48,B80,B221
metabolism, human 5
metal oxide fume B123
metal oxide fumes A13
metal pollution B202
metal salts A55
metal, heavy A20,A25
metals A52,B241
metals, atmospheric emission of B241
metals, heavy A13,A40,A43,B126,B157,B161,B200,B208,
B229
metals, trace A51
methanol A15,B36,B81,B87,B127,B164
methanol poisoning B87
methods, modern engineering B231
methods, treatment A75
methyl alcohol A9,B164
methyl cellulose acetate A14,B226
methyl ethyl ketone A26,B165
methylene chloride A41,B27
methylmercury B84
mice A14
Michler's hydrol, paratoluene sulfinate of A16
micro-hazards B12
microbes B220
microbes, inhaled B187
microbial aerosols B82
microbial contaminants B74
microbial growth A9,A54,B76
microbiological material, suspended B165
microcommunity B111
microelectronics B93
microflora, indoor B215
microorganism B21
microorganism, human B220
microorganisms A53,B28,B43,B76,B203,B215,B238
microorganisms, airborne B82
microwave power B28
microwaves B60,B132
middle age B22
middle class B181
middle class, white B120
middle managers A15,B185
middle-aged B88
middle-class whites, upper B9
middle-income-earning families A70
migraines A56,B171
mildew odours B199
milieu interieur B16
military A66
military resources A35
milk A57
milk samples, breast B136
milk, breast A57,B136
mill effluent, pulp and paper B208
mimicry of common illnesses B5
mimicry, disease B5
mindlessness B183
mine environments B49
mineral dusts A14
miners B114
miniature, adults in B213
minimal brain damage B167
minimal brain disfunction B167
minimal brain dysfunction B33,B167,B229
minimal brain dysfunction syndrome B113
minimal dependence B156
mining, phosphate B141
minor discomfort B47
minorities A85,B2,B139
minorities, cultural A24,B50
minorities, invisible B139
minority A79
minority characteristics 16
minority groups A38,B64,B233
minority groups, ethnic B44

-
- minority rights A80
 - minority views B139
 - minority, sensitive B140
 - mirex B136
 - mirex, kepone and B116
 - misdiagnosed A15
 - misdiagnosis B103
 - misfits, oppression of B81
 - misfits, social B151
 - mistakes B35
 - mistakes, correcting B92
 - mistreatment A38
 - misunderstood A45
 - misuse B163
 - mite, dust B48
 - mite, house dust B186
 - mites, house dust B186
 - mixed-use residential environments B183
 - mmpi, bender-gestalt, wais-r B170
 - mobile homes A6,B27,B128
 - mobility 5,B122
 - mobility limitations A30,B32
 - mobility problems B122
 - mobilization and excretion of xenobiotics B188
 - model, current medical B204
 - model, multiple-effects B64
 - modeled by peers, unhealthful behavior B76
 - modeling, peer B76
 - models, mathematical B5
 - modern engineering methods B231
 - modern job environments B193
 - modern technology A42,B229
 - modification, lifestyle B8
 - modification, process B134
 - modulation of pain A65
 - modulations, pulse B28
 - moist A54,B103
 - moist conditions B215
 - moisture B43
 - mold B51
 - mold sensitive individuals B29
 - molds B29
 - molds, toxins from B52
 - molecular biology B67
 - moles B62
 - moles, raised B62
 - molestation B34
 - molestation, sexual A26
 - money B91
 - money, bonus B39
 - monitoring A12,B91,B127,B136,B194
 - monitoring devices, health 15
 - monitoring stations, fixed B174
 - monitoring, biological B129
 - monitoring, personal B218
 - monitoring, personal exposure B155
 - monitoring, pollution B136
 - monitors B142
 - monomers B67,B243
 - monosodium salicylate B164
 - monotonous A68
 - monotonous jobs B208
 - monotonous work A32,B209
 - monoxide exposure, low-level carbon B7
 - monoxide poisoning, carbon B221
 - monoxide, carbon A5,A8,A15,A16,A19,A24,A52,A53,A76,
B5,B7,B19,B27,B29,B33,B41,B42,B43,B49,B57,B63,B80,
B88,B91,B92,B101,B107,B118,B121,B136,B141,B143,B155,
B184,B191,B195,B203,B211,B219,B228,B232
 - mood, subjective B235
 - mop, damp B165
 - moral underdevelopment B185
 - morbidity B22,B89,B129,B131
 - morbidity, high B191
 - morbidity, mortality and B117
 - morbidity/mortality data B237
 - mortality 12,A51,A52,A53,B22,B80,B88,B89,B129,B132,
B137,B143,B174,B197
 - mortality and morbidity B117
 - mortality data B132
 - mortality from aids 13
 - mortality rates B191
 - mortality studies B155
 - mortality, socioeconomic status and excess B80
 - mother B39,B76,B153
 - mother, single B62
 - mother-led households A36,B194
 - mothering A69,B33
 - mothers, smoking A5,B101
 - motion, ionic B102
 - motivation A65,B32
 - motivations B209
 - motives, ulterior B190
 - motor ability B198
 - motor coordination, impaired B23
 - motor dysfunctions, perceptual A52
 - motor impairments A25
 - motor performance A15,B235
 - motor sensory skill B235
 - motor skill, reductions in B56
 - motor vehicles A20,B218
 - motorization B143
 - motors A62
 - mould 15,A74,B48,B82,B123
 - mould avoidance B48
 - mould growth A77,B199
 - mould spores A54
 - mould, viable B82
 - moulds, sensitivities to A77
 - move, ability to B167
 - movement, air B123,B131
 - movement, environmental B61
 - movement, holistic health A77,B82
 - movement, self-care A77
 - movement, women's liberation B13
 - mucociliary transport, tracheal B70
 - mucosa, nasal B48
 - mucous membrane irritation B227
 - mucous membranes A74,B170
 - mucous membranes, dry B140
 - mucous, nasal B5
 - mucus, pharyngeal B210
 - multi-department program B70
 - multi-disciplinary programs A74
 - multicultural A38
 - multiculturalism 11,B120
 - multiorgan response B125
 - multiple deaths B185
 - multiple exposures B29
 - multiple sclerosis A9,B87
 - multiple-effects model B64
 - municipal building code B78
 - municipal incineration A51
 - municipal incinerators A20
 - municipal officials B63
 - municipal waste A52,B241
 - murder B41
 - muscle sensitization, smooth A83,B169
 - muscle strength A61
 - muscles, smooth A74,B170
 - music rooms A12
 - mutagenic B16,B80,B88,B159,B221
 - mutagenicity A53,B146,B158
 - mutagens A22,B168
 - mutual support A35,A37
 - my backyard phenomenon, not in B78
 - my backyard, not in A7
 - mycotoxins 15,A54,A57,A77,B187,B189,B215
 - mycotoxins, trichothecene B187
 - myelogenous leukemia, acute B186

- myeloid leukemia, acute B132
myrothecium B189
myth of the housewife B183
myths B193
names, trade B33
naphthalenes, chlorinated B136
narcotic B149
nasal congestion A16,B30,B31
nasal enzymes B48
nasal impedance B30
nasal inspiratory flow rate B186
nasal mucosa B48
nasal mucous B5
nasal provocation B186
nasal resistance B5
nation's health care services B91
national centres of research B90
national council, Swedish B123
national health policy B134
national health survey B179
National Inventory, Danish B6
national priorities B228
nations, developing B4
nations, industrial A57,B136
nations, industrialized B4
Nations, United B222,B236,B238
native B180
native community B96
native community health care workers B96
native Indians B191
native organizations B192
native people A38,B96,B191,B192
natural B47
natural allergens B3
natural daylight A10,B154
natural environment 3,B21
natural environments 5,A22,B156
natural gas B169
natural gas space heaters, unvented B73
natural radioactivity B221
natural setting B37
natural settings A22
naturally-based B47
nature of health, social B83
nature of individual responses, idiosyncratic B232
nature of relationships B44
nature, closeness to B183
nausea A6,A19,A33,A57,B7,B50,B80,B128,B193,B240
near-home territories B213
need for affiliation B12
need for control A67,B99
need for equality of opportunity B212
need, human B9,B99
needle-felt carpets A9,B76
needs of adults B25
needs of children B212
needs of individuals B196
needs of the young B213
needs of women B196
needs, basic B97
needs, basic human B97
needs, changing human service B224
needs, diversity of A78
needs, group B1
needs, health B8,B204
needs, human B163
needs, individual B106
needs, nutritional A56
needs, personal B107,B229
needs, physical B229
needs, psychological B78
needs, special or unique B35
needs, supportive of special B232
negative air ions A8,A63,B110
negative attitudes A44
negative bacilli, gram B210
negative life events B195
neglect A26,B34
neighborhood B4
neighborhood, residential B36
neighborhoods, dense urban A67,B12
neighborhoods, white B233
neighborliness B231
neighbors B122
neighbour-oriented B9
neighbourhood A21,B95,B162
neighbourhoods A67,B23
neighbourhoods, integrated caring B119
neighbourhoods, racially-integrated suburban B233
neonatal B101
nerve conduction B157
nerve endings, trigeminal B11
nerves, trigeminal A54,B62
nervous B17
nervous fatigue A10
nervous system B8,B49,B102,B103,B116,B193,B215
nervous system damage, central A15,A57,B215
nervous system depressants, central B172
nervous system dysfunction B51
nervous system effects B7
nervous system reactions B161
nervous system, autonomic B191
nervous system, central A8,A76,B23,B80,B119,B125,
B157,B164,B223,B225,B235
nervous system, peripheral B94,B235
network, social support 7,A67,B119
network, supportive social A29,A45,B81
networks, personal B68
networks, social A73,B68
networks, social, support A72
neuro-behavioural effects B179
neuro-physiological B102
neurobehavioral tests B164
neurobehavioral toxicity B235
neurobehavioural measures B199
neurobehavioural toxicity A8,A15,B172
neurobehavioral toxicity B164
neurologic changes A58
neurologic sequelae, subtle B192
neurological deficit B223
neurological deficits A25
neurological disorders B23
neurological disturbances B159
neurological effects B59
neurometrics B214
neuropathies, solvent B157
neuropathy, mbk B127
neuropathy, peripheral A13
neuropsychiatric disorders B10
neuropsychiatric disorders, nonspecific B10
neuropsychiatric symptoms A77,B29
neuropsychologic deficits B145
neuropsychologic effects B145
neurotoxic A56,B9
neurotoxicity B7,B145,B164,B170,B223
neurotoxicology B179
neurotoxin A25,B145
neurotransmitter A63,B110
new housing B35
new life setting B35
new perspective B8
new technologies 7,A81,A82
new world health order B117
New York B95
new-born B206
newborn A52,B18
newspapers A8,B51,B69,B130
niche A85
nitrate, isoamyl B172
nitrates, nitrites and B116

- nitrites and nitrates B116
 nitrogen compounds B91,B92
 nitrogen content A54
 nitrogen content, low B103
 nitrogen dioxide A5,A19,A41,A52,A53,B5,B7,B18,B19,
 B20,B41,B43,B57,B60,B65,B73,B74,B92,B93,B105,B109,
 B121,B135,B142,B147,B191,B196,B197,B202,B203,B211,
 B219
 nitrogen dioxide, asthmatics B153
 nitrogen dioxide B155
 nitrogen oxides B91,B116,B136,B141,B143,B156,B184,
 B187
 nitrogen, oxides of B123
 nitropyrenes A15
 nitrosamines A24,B75
 no-noise code, rigid B231
 noise A11,A16,A21,A32,B69,B109,B123,B140,B152,B208
 noise exposure B230
 noise on sleep B230
 noise pollution B17,B77,B126,B139,B232
 noise reduction B231
 noise, abatement of B231
 noise, aircraft A60,B77,B139
 noise, community A65,B32
 noise, outside B1
 noise, sociological response to B230
 noise, unpredictable B231
 noise, urban B12
 noisy A65,B183
 noisy exhaust fans B165
 non smokers B206
 non-adaptive classroom behavior B145
 non-ambulant A42
 non-ambulant disabled persons B103
 non-compliance B7
 non-discrimination in employment B139
 non-dominant groups B27
 non-immunological mechanisms B68
 non-industrial B76
 non-industrial setting B3
 non-ionizing B102
 non-ionizing radiation B102
 non-judgmental health care A49,B156
 non-odorous products A7
 non-physiologic chemicals, hypersensitivity to B193
 non-polluting product A85
 non-polluting products 18
 non-renewable resources A22,B222
 non-smoker B231
 non-smoker families B107
 non-smokers B42,B86,B89,B144,B207,B231
 non-smokers, rights of B144
 non-smoking households A26
 non-toxic B47
 non-toxic substances B47
 non-visual effects B84
 nonmanagerial personnel B176
 nonpoverty areas B80
 nonsmokers A5,A14,B173,B174,B176
 nonsmokers, risks to B175
 nonspecific clearance system B188
 nonspecific neuropsychiatric disorders B10
 nonsusceptible workers B240
 nonthermal electromagnetic fields B17
 nonthreshold effects B168
 nontoxic B47
 nontraumatic phlebitis B171
 noradrenaline B152
 nordic countries B83
 normal, desire to appear as B122
 normative images B27
 normotensives B146
 norms B62
 northern B216
 nose B169,B170
 nose, dryness in the B6
 nose, running A6
 not in my backyard A7
 not in my backyard phenomenon B78
 notification system B6
 noxious agent B157
 noxious stimulation A65,B32
 nuclear accident A67,B61
 nuclear accidents A68
 nuclear age B74
 nuclear arms A35,B13
 nuclear energy B90,B237
 nuclear power A66,B60
 nuclear technology A66
 nuclear war A66,B74
 nuclear weapon B61
 nuclear weapons A22,A67,B60,B222
 nuclei, droplet B178
 nuclides, radon daughter A53
 number of close ties B87
 number of people in a space B234
 numbers, decisions from B201
 nursery B124
 nurses A32,A78,B116,B134,B209
 nurses' associations B116
 nursing A26,B116,B165
 nursing homes A33,B122,B166
 nutrient-toxicant interactions B126
 nutrients B30
 nutrition A30,B30,B51,B126,B142,B178,B187,B192,B204,
 B214
 nutrition, poor A70,A76,B201
 nutritional deficiencies A56,B23,B73
 nutritional deficiency A41,B126
 nutritional hypersusceptibility B173
 nutritional levels, poor B113
 nutritional needs A56
 nutritional programmes B152
 nutritional status A40
 nutritional training A26
 nutritionists B142
 nutritious food A2,B161
 nylon A55,B14
 oasis A11
 obesity A70,B22,B137
 obey, compelled to A27,B50
 obsessive-compulsive trends B54
 obsessive-compulsive trends A64
 obstacles B233
 obstetrical management B131
 obstetrics B136
 obstruction, upper airway A16
 occupancy B183
 occupants' satisfaction B75
 occupation B28,B46
 occupational B15,B49,B79,B111,B172,B184,B185,B186
 occupational asthma B82
 occupational categories B208
 occupational choices A76
 occupational disease A75,B241
 occupational diseases A13,B28,B36,B135
 occupational exposure B87,B185,B186
 occupational exposure, chronic B69
 occupational hazards B78,B102,B135
 occupational health B42,B69,B72,B87,B121,B125,B126,
 B129,B152,B161,B174,B177,B185,B211,B236,B240
 occupational health and safety A1,A34,B3,B33,B38,B44,
 B73,B123,B127,B128,B134,B135,B157,B165,B180,B185,
 B190,B191,B208,B221,B235,B241
 occupational health clinics, independent B241
 occupational health hazards B134,B186
 occupational health policy B185
 occupational health problems B73
 occupational history A75
 occupational lighting sources, unusual B62

- occupational medicine B173,B208
 occupational regulations B15
 occupational respiratory allergic disorders B159
 occupational safety B24
 occupational safety and health B192
 occupational standards B226
 occupational stress B44
 occupational threshold limit value B227
 octachlorostyrene B136
 occupational safety B57
 odors B62
 odour A54,B29
 odour dispersal B165
 odour patterns B19
 odour, body B62
 odours A82,B62,B108
 odours, mildew B199
 odours, sensitivity to A74,B169
 of indoor air B19
 of, health B37
 off-gassing B14
 off-task behaviours B97
 offenders B178
 office A62,A74,B35,B69,B82,B113,B129,B137,B177,B227
 office accidents A16,B35
 office building B115,B149
 office buildings B131,B206,B217,B220
 office buildings, airtight B220
 office buildings, sealed A16,B35,B207
 office complex, hermetically sealed B18
 office complex, three-tower government B131
 office environment B69
 office environment, typical B127
 office environments B90
 office equipment B74
 office hazards B126
 office lighting B240
 office politics B44
 office products B149
 office work environments B207
 office workers B77,B148,B149,B207,B239,B240
 office workers, women B78
 office, home B37
 offices 6,B76,B78,B127,B128,B165,B206,B211,B239
 offices, academic A11,B212
 offices, aldehydes in B137
 offices, closed B131
 offices, windows and lighting in B239
 officials, elected B188
 officials, health B240
 officials, law enforcement A38
 officials, municipal B63
 offspring B87
 oil combustion A22,B217
 oil, cutting A26,B165
 oil, road B223
 oil, waste B223
 oilway A38,B180
 old B25
 old age B86
 old people B35,B36,B151
 older adults B106
 older age groups B130
 older gay people B81
 older people B111,B148
 older persons B85
 older unemployed individuals B47
 older women A33,B122
 older workers B112
 olfactory B19
 one's life, control over 5
 Ontario B63,B89,B134,B152
 Ontario Human Rights Code B41
 Ontario Indians B180
 Ontario Indians, Union of B180
 Ontario longitudinal study of aging B89
 open plan B131
 open spaces B21
 open windows B172
 operating practices A10
 operators, radar B186
 operators, radar B186
 opportunistic infection B187
 opportunistic infections A77
 opportunities B84
 opportunities, employment B85
 opportunities, work B95
 opportunity, equal A32
 opportunity, need for equality of B212
 oppressed group B26
 oppression 5,6,11,A79,B2,B27,B41
 oppression of misfits B81
 oppression of working-class men B114
 oppression of young people B220
 oppression, gay 17,A46,A49,B112
 oppression, institutionalized A32
 oppression, umbrella of B26
 oppressive A35
 oppressive behaviour A80
 oppressive environment A46
 oppressive relationships A34
 oppressor role B99
 optic tract B186
 optimal functioning B72
 optimism B112
 optimum environment B98
 options, reduced life B134
 options, work B93,B151
 oral contraceptives A70,B137
 order, inequitable social B2
 order, new world health B117
 organ systems A84
 organ systems, vital B119
 organ, target B193
 organic brain disorders B160
 organic chemicals, volatile 15,B5
 organic compound B227
 organic compound emissions B137
 organic compounds B140,B184
 organic compounds, total volatile B227
 organic compounds, volatile A54,B57,B140,B142,B243
 organic contaminants, volatile A77
 organic dusts B73
 organic gases B11
 organic solvents A24,B9,B75
 organic solvents, volatile A13
 organic vapours A53
 organics, volatile B227
 organisms, airborne B177
 organisms, biological B37
 organisms, viable B120
 organization of worktime B196
 organization, social B7
 organizational framework B71
 organizational settings A72,B105
 organizational structure B44
 organizations, consumer B83
 organizations, native B192
 organo-phosphates B49
 organochlorine compounds A58
 organochlorine pesticides B193
 organohalides B180
 organophosphorous compounds B232
 organophosphorus insecticides A58,B72
 orientation 3,A2,A46
 orientation, psychosocial B183
 orientation, sexual 6,13,17,A46,A47,A80,B26,B41,B112,
 B156
 origin, ethnic A38,A80,B218
 orthomolecular therapy B73

- orthostatic hypotension B22
 osteomalacia B30
 ostracism A47
 ostracism, social B55
 others, relationships with B114
 otitis A74,B169
 out, coming A47,B86
 outbreak B177
 outcasts A79
 outcomes, deleterious reproductive B192
 outdated perception B218
 outdoor air 5
 outdoor air pollution B158,B166,B174
 outdoor environment B21,B84
 outdoor exposures B227
 outdoor locations B213
 outdoor space B84
 outdoor spaces A20,B84
 outdoor/indoor exposure B147
 outlet for expression B9
 outside environment, eyes to the B222
 outside happenings B222
 outside noise B1
 outside the home, work B183
 outsiders B195
 overcrowding A21,B229
 overexposure, chemical B168
 overload A64
 overload of synthetic chemicals B170
 overload, cognitive B181
 overload, decision A65
 overload, decisional B183
 overload, environmental B169
 overload, information A61,A65,B183,B222
 overload, staff B216
 overload, work A68,B44
 overpowered A75,B65
 oversupply of physicians B224
 overt discrimination B55
 overt prejudice against homosexuality A49
 overtime B28
 overuse B163
 overworked A71,B151
 owner B37
 oxazepam B172
 oxidant B105
 oxidants A52,B156
 oxidants, photochemical B191
 oxidase deficiency, sulfite B88
 oxide fume, metal B123
 oxide fumes, metal A13
 oxides B241
 oxides of nitrogen B123
 oxides, nitrogen B91,B116,B136,B141,B143,B156,B184,
 B187
 oxides, sulfur B91,B211
 oxides, sulphur A51
 oxygen transport B49
 ozone A5,A13,A16,A40,A51,A52,B23,B30,B43,B60,B82,B91,
 B101,B105,B109,B123,B142,B156,B162,B196,B211
 pace of life A65,B183
 paced assembly lines B44
 packing materials B53
 PAHs A83
 paid labour force B33
 paid work A34
 paid work, dignifying B219
 paid workforce B208
 pain B66
 pain, abdominal A57,B240
 pain, chest A83,B169
 pain, chronic low back A65,B66
 pain, modulation of A65
 pains, leg B103
 paint remover A41,B207,B208
 paint, house B89
 paint, spray A26,B165
 painters A13,B9,B10
 painters, house B83
 painters, spray A13,B127
 painting A5,A8,B84,B127,B190,B199
 painting trades B190
 paints 15,B6,B200
 paintwork, deteriorating B137
 pancreas B231
 pandemic B173
 paneling, indoor B152
 panelling, wall B128
 panels, wood-chip B159
 paper A8
 paper dermatitis B69
 paper mill effluent, pulp and B208
 paper products A5,B97
 paper, art A8,B69
 paper, carbon B69
 paper, carbonless A8,B69
 paper, carbonless copy B30,B129
 paper, copier B69
 paper, typing B69
 papers, photocopier A8
 paradigm, ecological B37
 parameters, immune B170
 paranoia A43,B103
 paraquat A57,B215
 parasuicides B152
 parathion A58,B99
 paratoluene sulfinate B129
 paratoluene sulfinate of michler's hydrol A16
 parent B45,B75
 parent's workplace exposure B165
 parental B76
 parental permission B84,B144
 parental smoking A10,A26,B24,B65,B76,B212
 parentally-defined range restrictions B84
 parenting A32,A35,B13
 parenting relationships B45
 parents B37,B71,B74,B76,B122,B146,B149
 parents' desire B122
 parents' exposures B165
 parents, property of their B213
 parking A16
 parking, indoor B155
 parks B243
 part-time work A18,A36,B45
 participants B35
 participate, inability to B206
 participating B95
 participating in caretaking B7
 participating in design A67
 participation 3,11,18,19,A21,A64,A78,A79,A84,B33,B44,
 B95,B148
 participation in design A84
 participation, citizen A67,B9,B23
 participation, community B36,B95,B239
 participation, family A75,B65
 participation, public B9,B119,B120
 participation, user 7
 participation, worker A15,A68,B185
 participatory planning B99
 participatory roles of children B25
 particle B42
 particle board A53,B5,B79,B152
 particle boards A6
 particle mass B155
 particle size B184,B233
 particle-board A19
 particle-free environment A83,B168
 particleboard B20,B70,B106
 particleboard floor underlay B106
 particles B54,B100,B136,B142

- particles, beta A62,B20
particles, personal exposures to B156
particles, respirable B42
particles, resuspension of precipitated B165
particulate B15,B158,B173
particulate air pollution A52,B155
particulate emissions B73
particulate fraction B123
particulate inhalants B125
particulate levels, elevated B94
particulate mass, respirable B218
particulate matter B43,B165,B196
particulate matter, suspended B105
particulate pollution B15,B233
particulate, mass respirable B65
particulate, suspended B29
particulates A15,B43
particulates, respirable A53
particulates, respirable suspended B93,B173
particulates, suspended B205
partitioned B131
partners, difficulties with B40
partners, sexual B201
parttime work, flexible B93
passer-by B37
passive A27
passive exposure B200
passive sampler B148
passive sampling B190
passive smoking B10,B24,B107,B120,B132,B158,B174,
B175,B203,B207,B217,B231
past home B111
paternalism A43,B95,B100
paternalistic B216
path, soft health B82
pathogen, viral B32
pathogenic mechanisms B74
pathogens, climatic B177
pathological defense mechanisms B90
pathological findings B187
pathology B42,B174
pathology of human relationship B25,
pathology, fetal B101
pathways, ecological B60
patient, de-institutionalized B216
patient, psychiatric A43,B112
patients in acute care B233
patients, chemically sensitive B171
patients, female B136
patients, infectious disease B119
patients, psychiatric B34
patterns of behaviour B182
patterns, chemical B19
patterns, employment A69,B34
patterns, land use B43
patterns, odour B19
patterns, response B54
patterns, sleep B80
patterns, work B196
pay, equal A32
pay, low B209
payoff, clinical B31
pbbs B188
pcb A68
pcb spill B200
pcbs A15,A50,A54,A57,B64,B116,B124,B136,B168,B180,
B188,B242
pcdds B242
pcdfs B242
peaceful purposes B90
penches A42
penk expiratory flow B41
peckham experiment B157
pectoris, angina B10
pediatric cancer B74
pediatric disorders B52
pediatricians B243
peer behaviors B76
peer cohesion A65,B66
peer modeling B76
peer support A73,B76
peers A46
peers, gay-identified A46,B173
peers, unhealthful behavior modeled by B76
pejorative B51
penicillium A57
penicillium toxins B189
pension entitlements, enriched B149
pensions B33
pentobarbital A9,B172
pentobarbital-like discriminative effects B172
people A80
people acting through consent B133
people in a space, number of B234
people of value A29
people with mental disabilities B39
people's liberation, young B220
people's reactions to burglary B113
people's well-being B222
people, able-bodied A4
people, aboriginal A39
people, aging gay A30
people, children as B235
people, control of B133
people, environmentally-sensitive A22
people, exiled B133
people, forgotten B130
people, handicapped 6
people, invisibility of gay B40
people, native A38,B96,B191,B192
people, old B35,B36,B151
people, older B111,B148
people, older gay B81
people, oppression of young B220
people, variability among A24
people, young 12,A24,A27,B152
people/ environmental relationships B235
peoples, dominated B2
perceived climate B98
perceived control A85,B32
perception B9,B74
perception, outdated B218
perceptions B176,B184,B193,B238
perceptions, child's A50
perceptual B198,B202
perceptual capabilities A50,B209
perceptual disorders A52
perceptual motor dysfunctions A52
perceptual problems B150
perennial rhinitis B186
performance 7,A76,B63,B93,B235,B239
performance decrement B118
performance deficits A76
performance effects 15
performance in traffic, children's B202
performance indicators 13
performance measurements A83
performance measures B75
performance tests B11,B223
performance, classroom B145
performance, mental A54,B140
performance, motor A15,B235
performance, scholastic A10,B24,B166
performance, school B64
performance, total building A10,B75
performance, ventilation B207
performance, ventilatory B194
performance, work B17
perfume A74,B169
perfumed personal hygiene products A11

- perfumes A44,B81
 periods of exposure B232
 periods of unemployment B112
 peripheral nervous system B94,B235
 peripheral neuropathy A13
 permanence of the disability B122
 permanent disability A29,A50,B89
 permanent housing B78
 permeability of the placenta B64
 permissible dietary intake, maximum B72
 permission, parental B84,B144
 permissive B231
 persistent symptoms A15
 person-oriented B57
 person-to-person transmission B214
 personal B33
 personal attitudes A79,A80
 personal behaviour B8
 personal biases B173
 personal characteristics B148
 personal control 7,A30,B106,B233
 personal exposure B147
 personal exposure monitoring B155
 personal exposures to particles B156
 personal financial resources B122
 personal goals B93
 personal health B149
 personal health care B8
 personal hygiene B173
 personal hygiene products, perfumed A11
 personal infringement B230
 personal injury B192
 personal interactions B147
 personal limitations B193
 personal monitoring B218
 personal needs B107,B229
 personal networks B68
 personal power A28
 personal relationships A26,B40
 personal relationships, strained B40
 personal resources B106
 personal responsibility B8
 personal samplers B196
 personal uncertainty B80
 personality A60
 personality changes A13
 personality disorders, chemically-induced B9
 personality factors B54,B230
 personality traits A65,A72,B32,B55
 personalization A31,B111
 personnel, disregard by medical B122
 personnel, highly specialized B213
 personnel, nonmanagerial B176
 persons, children as B212
 persons, mentally ill B51
 persons, non-ambulant disabled B103
 persons, older B85
 persons, respect for B144
 perspective, child's B193
 perspective, consumer B95
 perspective, ecological B106
 perspective, new B8
 perspective, social-ecological B98
 perspectives, environmental B144
 pessimism B112
 pest control 5
 pesticide A57,A58,B171
 pesticide dumping B215
 pesticide exposure B192
 pesticide exposures A62
 pesticide residues A82
 pesticide toxicity B193
 pesticide use B48
 pesticide users B192
 pesticide-free home B171
 pesticides A20,A40,A43,A51,B15,B50,B53,B60,B106,B117,
 B126,B161,B165,B169,B170,B179,B192,B193,B229,B242
 pesticides, banned B215
 pesticides, chlorinated B188
 pesticides, chlorinated hydrocarbon A82,B117
 pesticides, garden A26
 pesticides, organochlorine B193
 petrochemicals B50
 pets, exotic B213
 pH B25
 phagocytic immune functioning B53
 pharmaceutical A57
 pharmaceutical industries B240
 pharmaceutical industry B240
 pharmaceuticals B60,B240
 pharyngeal mucus B210
 pharynx B6,B231
 phenol A74,B169
 phenomena, spastic vascular A56,B171
 phenomenon, masking B169
 phenomenon, not in my backyard B78
 phenomenon, spreading B169,B170
 phenoxy herbicides B192,B193
 philosophic beliefs B100
 philosophical concepts B142
 phlebitis B171
 phlebitis, nontraumatic B171
 phlegm B87
 phosphate mining B141
 phosphate, tricresyl B195
 phosphates B116
 phosvel A57,B215
 photochemical air pollution B104
 photochemical oxidants B191
 photochemical smog B83
 photocopier papers A8
 photocopiers B82,B125,B126,B149
 photocopiers, liquid process B227
 photocopiers, wet-process A15,B137
 photocopying machines A16
 photography, time-lapse B155
 photooxidation reactions B97
 phthalate, butyl benzyl B178
 physical B135
 physical abuse B34,B134
 physical activity B76,B80
 physical agents B75,B241
 physical ailments, serious B47
 physical characteristics B158
 physical comfort B66
 physical design B107
 physical disabilities, coping with B148
 physical disability A79
 physical disability, mental and B234
 physical diversification B233
 physical effort B44
 physical environment B9,B36,B58,B84,B107,B140,B148,
 B233
 physical environment, control in the B65
 physical environments, urban B164
 physical examination B128
 physical exploitation B114
 physical factors B206
 physical form B69
 physical health, emotional and B101
 physical health, poor B44,B88
 physical health, unemployment and B112
 physical illness B47
 physical inactivity B137
 physical injury B134,B152
 physical needs B229
 physical planners B9
 physical planning B183
 physical planning decisions B9
 physical setting B1,B164

- physical settings, decay of B163
physical violence B220
physically disabled A80
physician B79
physician visits B47
physician, general A74
physicians B134, B142, B177, B204, B240, B241
physicians, chauvinistic male B136
physicians, oversupply of B224
physiological B17, B230
physiological change B230
physiological effects B149, B232
physiological responses B230
physiological rhythms A61
physiological strain B44
physiological stress B150
picture, chronic unemployment B33
pigmentation B62
pigments B165
pine-scented floor wash B169
pipes, water A58, B189
pitfalls to avoid B225
placebo A76
placebo response B50
placenta, permeability of the B64
places B163
places of work B76
places, high-energy A21, A64, B95
places, public J, A19, B220
plaintiff B79
plan of action B123
plan, open B131
planet B236
planners, physical B9
planning decisions, physical B9
planning processes B98
planning, community B231
planning, development B222
planning, family A78, B116, B152
planning, health B8, B118
planning, long term health B118
planning, participatory B99
planning, physical B183
planning, urban A21, B8, B162
plant closings B140
plant fungi B58
planting trees B9
plants A56
plastic A58, B25, B105, B195
plastic chairs B195
plastics A5, B53
play B65, B231
play facilities B65
play spaces A22, B21
play, prohibition of A65
playground B243
playgrounds B21, B83
plight of the homeless B78
plumbing A22
plumbing system B70
plutonium A67, B20
plutonium, toxicity of B61
plywood B70, B79
pneumonia A58, B236
pneumonia, streptococcal B60
pneumonitis, hypersensitivity B177
pneumophila, legionella A59, B237
points system, priority B63
poisoning B72, B114
poisoning of air B90
poisoning, carbon monoxide B221
poisoning, food B5
poisoning, lead B64, B111
poisoning, methanol B87
polarized ideas B183
police racism B64
police/community relations B41
policies B57, B182
policies for leaves of absence B93
policies, canadian health and social B142
policies, environmental health B237
policies, equity-oriented educational B239
policies, government B134
policies, health care B54
policies, intersectoral B83
policies, local B243
policies, public health B138
policies, shelter B149
policy B66, B151, B205
policy alternatives, public B203
policy brutality B64
policy coordination B144
policy framework B34
policy issues, public B36, B144, B236
policy makers B83
policy responses B185
policy, compensation B241
policy, government B133, B135, B241
policy, health B83, B103, B116, B142
policy, health care B7
policy, healthy public B83
policy, human rights B97
policy, national health B134
policy, occupational health B185
policy, public B8, B9, B33, B34, B35, B39, B42, B45, B49, B57, B58, B63, B82, B85, B87, B90, B93, B94, B95, B96, B97, B98, B103, B112, B115, B118, B119, B120, B121, B127, B129, B133, B134, B137, B140, B144, B148, B151, B156, B176, B179, B183, B188, B191, B196, B197, B200, B201, B204, B206, B209, B212, B216, B217, B221, B222, B228, B232, B237, B239
policy, public health B22, B37, B57, B60, B83, B96, B100, B120, B122, B204, B224, B229
policy, social B90, B218, B233
policy-making B57
policy-making institution B188
political B37, B235
political factors B119
political pressures B4
political rights A49, B139
political vision 19
politicians 17
politics B188
politics, office B44
poll, gallup B218
pollens B211
pollutant exposure standards B30
pollutant, atmospheric B133
pollutants B30, B37, B188, B189
pollutants, chemical B18, B229
pollutants, exhaust of B199
pollutants, inhaled B160
pollutants, inhaled air B119
pollutants, longrange transport of airborne B70
pollutants, man-made B37
pollutants, range of vulnerability to B199
pollutants, toxicity of environmental B126, B192
pollutants, volatile B16
polluted indoor air J
polluted indoor environments A74
polluted society B133
polluting industries B27
pollution B12, B18, B23, B30, B45, B60, B72, B73, B94, B115, B118, B126, B127, B149, B160, B161, B162, B164, B166, B187, B188, B200, B225, B230, B233, B236
pollution control B27
pollution control, air B184, B223
pollution design, low B184
pollution detection, air B91
pollution monitoring B136
pollution relief centres A11

- pollution sampler, domestic air B93
 pollution source identification B128
 pollution sources B184
 pollution, air B5,B15,B18,B19,B23,B25,B28,B36,B38,
 B41,B43,B46,B49,B50,B52,B53,B60,B61,B63,B67,B79,B81,
 B87,B88,B89,B90,B91,B92,B93,B97,B100,B104,B105,B107,
 B108,B109,B111,B115,B116,B117,B118,B119,B124,B133,
 B137,B141,B143,B145,B152,B156,B160,B172,B184,B187,
 B189,B190,B191,B192,B194,B196,B197,B200,B203,B204,
 B210,B213,B214,B223,B224,B229,B237,B241,B242
 pollution, air indoor B86
 pollution, chemical B33,B127,B172
 pollution, domestic air B93
 pollution, environmental B126,B131,B134,B136,B161,
 B190,B204,B232
 pollution, great lakes A58,B99
 pollution, health effects of B77
 pollution, indoor B210
 pollution, indoor air 5,B5,B6,B11,B13,B14,B15,B16,
 B17,B19,B21,B22,B27,B28,B29,B35,B38,B42,B43,B46,B47,
 B48,B53,B54,B55,B57,B62,B65,B67,B69,B73,B74,B78,B79,
 B82,B88,B89,B90,B91,B92,B97,B101,B108,B109,B113,B115,
 B120,B123,B124,B126,B127,B128,B135,B136,B137,B139,
 B140,B141,B142,B145,B147,B149,B150,B152,B155,B158,
 B164,B165,B166,B167,B173,B174,B175,B176,B177,B178,
 B181,B184,B186,B188,B189,B190,B194,B196,B197,B198,
 B202,B203,B206,B207,B211,B212,B217,B219,B221,B222,
 B225,B226,B231,B234,B236,B242
 pollution, internal B16
 pollution, low levels of B233
 pollution, measurement of B79
 pollution, metal B202
 pollution, noise B17,B77,B126,B139,B232
 pollution, outdoor air B158,B166,B174
 pollution, particulate B15,B233
 pollution, particulate air A52,B155
 pollution, photochemical air B104
 pollution, radon B88,B181
 pollution, soil B23
 pollution, thermal B17
 pollution, tolerance for A84
 pollution, visual B17
 pollution, water B12,B116,B189
 pollution-free air B119
 polybrominated biphenyls B188
 polychlorinated B188
 polychlorinated biphenyls A54,B180
 polycyclic aromatic hydrocarbon A14
 polycyclic aromatic hydrocarbons A83,B52,B117
 polycyclic hydrocarbons B211
 polyester A55,B14,B123
 polyethylene food containers B105
 polymerization B53
 polymers A55,B14
 polyurethane B6
 poor B13,B25,B201
 poor health B80
 poor housing B229
 poor lighting B206,B208
 poor mental health B88
 poor nutrition A70,A76,B201
 poor nutritional levels B113
 poor physical health B44,B88
 poor ventilation B63,B206,B208
 poor, children of the A70
 poor, elderly B194
 poorer health habits B243
 poorly designed furniture B208
 popular print media B51
 population B37
 population groups, vulnerable B239
 population growth B116
 population, adult B149
 population, aging A74,B224
 population, allergic B198
 population, chemically susceptible B198
 population, sensitivity of the B124
 population, stigmatized B201
 population, stressed A60
 population, susceptible B198
 population-at-risk records B129
 populations, comparison 9
 populations, healthy B232
 populations, sensitive B5
 populations, susceptible B215
 porches B213
 pork A57
 pornography A26,B34
 portable classrooms B210
 portland cement B162
 portrayal of mental illness B51
 ports B222
 positive influences A64,B95
 positive mental health practices B34
 possessions B111
 possessions, decorative B112
 post-industrial development A69
 post-industrial society A74
 postindustrial society B224
 postnatal prevention issues B131
 potential, visual evoked B24
 potentials, electrical B230
 poverty 3,5,13,A36,A43,A64,A66,A70,A78,A79,B24,B62,
 B71,B80,B95,B113,B115,B116,B137,B138,B183,B194,B217
 poverty among women 12
 poverty and education B46
 poverty area B80
 poverty line A32
 poverty, child B149
 poverty, cycle of A33,B134
 poverty-level incomes A36,B194
 power A27,A35,A66,B9,B12,B26,B117,B186
 power and telephone linemen B186
 power frequencies B102
 power frequency fields A61
 power frequency range A62
 power frequency, electromagnetic sensitivity B171
 power transmission, electrical A61,B28
 power, centres of B25
 power, microwave B28
 power, nuclear A66,B60
 power, personal A28
 power, sense of A67
 power, token A79
 powerful B183
 powerless A27
 powerlessness A43,B12,B95
 practical difficulties B122
 practice, social B212
 practices, educational B235
 practices, engineering B238
 practices, institutional B7
 practices, operating A10
 practices, positive mental health B34
 practices, professional B46
 practices, same-gender sexual A46
 pre-school children B205
 precipitated particles, resuspension of B165
 precipitation, acid A52,B33,B156
 precipitation, acidic A22
 predelinquents B147
 predictability A60,A66,B107,B231
 predisposed, biochemically B147
 predisposing factors A40
 predispositions, genetic A68,B37
 prefabricated woodchip board B160
 preference B167
 pregnancy A24,A26,B75,B165,B228
 pregnancy problems in women B206
 pregnancy, unwanted 13

- pregnant rats B59
 pregnant women A40,B5,B198
 prejudice 5,6,13,17,19,A7,A23,A38,A46,A47,A79,A80,
 A84,H40,R41,B167
 prejudice against children B50
 prejudice against homosexuality, overt A49
 prejudice, culture of B211
 prejudice, eliminating 15
 prejudice, homophobic B41
 prejudice, interrupting 11
 prejudices 3
 prejudices against homosexuality B2
 prejudices, social A28
 prejudices, structural 3
 prejudicial treatment A32
 premature ventricular contractions A55,B191
 Premier's Council on Healthy Strategy B64
 premises, unexamined B50
 prenatal A24,B131
 prenatal care B131
 prenatal exposure B64,B75
 preparation programs, work B216
 preschool B19
 preschool age B52
 preschools A12
 prescription drugs A15
 preservatives A57,B168,B240
 preservatives, sulfites as B168
 press coverage B129
 press treatments, durable B14
 press, freedom of B139
 pressure, blood A53,B10,B22,B146
 pressure, diastolic blood A70,B146
 pressure, elevated blood B152
 pressure, elevated diastolic blood B137
 pressure, high blood A69,B47,B189
 pressure, increased blood B44
 pressure, work A65,B66
 pressures, political B4
 pressures, work 5
 pressures, work-associated B34
 preventable health issue B145
 prevention B22,B46,B51,B59,B71,B98,B123,B131,B147,
 B241
 prevention issues, postnatal B131
 prevention of cancer B46
 prevention of environmental problems B71
 prevention of mental illness B98
 prevention programs B76
 prevention programs, primary B71
 prevention techniques B113
 prevention, accident B70,B179,B211
 prevention, cancer B46
 prevention, disease A67,A77,B46,B119,B213
 prevention, handbook of B59
 prevention, illness B204
 prevention, injury B179
 prevention, strategies for B144
 preventive behavior B59
 preventive measures B123
 preventive medicine A78,B204
 preventive mental health care B34
 preventive services B131
 preventive strategies B233
 price labels B157
 prices, fuel B226
 pride B220
 primary blocks B192
 primary care B63
 primary prevention programs B71
 principal family earners B47
 principles, ethical B144
 principles, toxicological B6
 print media, popular U51
 printing A61,B62
 printmaking A5,B84
 priorities B130
 priorities, national B228
 priorities, research B3,B31,B35,B37,B51,B52,B55,B61,
 B65,B69,B82,B85,B101,B111,B114,B117,B120,B123,B143,
 B153,B155,B157,B163,B179,B187,B191,B199,B203,B206,
 B209,B214,B221,B229,B232,B237
 priority B117
 priority points system B63
 privacy A29,A65,A75,B56,B118,B163
 privacy invasion B118
 privacy problems B233
 privacy, acoustic A60,B230,B231
 privacy, lack of B235
 private homes B220
 private sector B91
 private sector, discrimination in the B41
 private sectors B34
 private spaces B111
 privilege A35
 privileges A47
 problem solving B84
 problem solving, institutional B133
 problem, endemic social B193
 problem, safety B130
 problem-solvers B59
 problem-solving A35
 problems during childhood, behaviour B134
 problems in women, pregnancy B206
 problems of women, health-related B239
 problems, access-oriented B211
 problems, behavior B71
 problems, behaviour B101,B167
 problems, behavioural B167,B243
 problems, bone-joint A69,B47
 problems, cardiac B198
 problems, ear and hearing B29
 problems, family 5
 problems, gastro-intestinal B160
 problems, health B37,B47
 problems, hearing A77
 problems, learning A45,B150,B167
 problems, memory B15
 problems, mental health B101,B140,B191
 problems, mobility B122
 problems, occupational health B73
 problems, perceptual B150
 problems, prevention of environmental B71
 problems, privacy B233
 problems, psychosocial B140
 problems, public health B179
 problems, research B91
 problems, respiratory B29,B209
 problems, school-related B112
 problems, social B142,B179,B233
 problems, structural A68
 procedures, abatement B125,B165
 procedures, drug discrimination B172
 procedures, medical B136
 procedures, research B36
 procedures, safety B103
 procedures, surgical A33
 proceedings, conference B147
 process modification B134
 process of consent A28
 process photocopiers, liquid B227
 process, decision-making B43,B44
 process, design B234
 process, detective 12
 process, grieving B112
 processed foods A57
 processed foods, labeling of B168
 processes of healing U57
 processes, age-related B86
 processes, cortical inhibition B24

- processes, decision H209
 processes, healing B119
 processes, industrial 5,A52,B60,B241
 processes, planning B98
 processes, reproduction A8
 processes, risk assessment B237
 processes, social B37
 processing, speech B145
 product design A50
 product disclosure B33
 product labelling B138
 product review B75
 product, non-polluting A85
 production, food A67,B119
 production, fungi B103
 productive work B183
 productivity B35,B49,B61,B199
 productivity, work B240
 products B6
 products in daily use B47
 products, building A6
 products, by B37
 products, combustion B91,B166,B220
 products, consumer A50,B47,B97,B179
 products, daughter A6
 products, electronic B63
 products, emissions from B75
 products, forest B70
 products, household B220
 products, injurious B179
 products, laundry A5
 products, less-polluting 15,A85
 products, maintenance 5,A15,B137
 products, non-odorous A7
 products, non-polluting 18
 products, office B149
 products, paper A5,B97
 products, perfumed personal hygiene A11
 products, radon daughter A54
 products, radon decay B89
 products, scented cleaning B166
 products, wood B14
 professional awareness A28
 professional background B47
 professional dominance B97
 professional educators B222
 professional freedom B142
 professional level B216
 professional practices B46
 professional training B40
 professional woman B39
 professionals B1,B9,B93,B219
 professionals, caregiving B54
 professionals, health B35,B69,B243
 professionals, health care B142,B195
 professionals, mental health B141,B216
 professions B96
 professions, disabling B96
 professions, health care B134
 profile, distressed B39
 profit-oriented corporation B188
 profitability B188
 profound distrust of adults B74
 program compliance B76
 program, community fitness B76
 program, exercise B76
 program, health education B53
 program, hidden B69
 program, management B243
 program, multi-department B70
 programmable ambulatory electrocardiograph recordings B191
 programmes, nutritional B152
 programming B234
 programs, abatement B70
 programs, bridging A38,B44
 programs, clinical B204
 programs, design of B8
 programs, educational B204
 programs, environmental B199
 programs, health promotion B86,B137
 programs, medical research B91
 programs, multi-disciplinary A74
 programs, prevention B76
 programs, primary prevention B71
 programs, safety B185
 programs, universal leave B149
 programs, volunteer B216
 programs, work preparation B216
 prohibiting smoking B207
 prohibition of play A65
 projects, research B152
 promote health, behaviours that B204
 promoters, weak cancer B132
 promoting health B117
 promotion B152
 promotion programs, health B86,B137
 promotion strategies, health B204
 promotion, health A30,A67,A77,A78,B57,B63,B96,B119,
 B133,B134,B204,B213
 promotion, mental fitness B86
 promotion, message B144
 propaganda B41
 propane B27,B162
 propellants, aerosol B27
 properties, human B163
 properties, rental A36,B63
 property B139
 properly damage B192
 property of their parents B213
 property rights B188
 property, distribution of A47,B2
 prosecution, criminal B185
 prostitution A26,A46,B34,B173
 protected victims B147
 protection act, environmental B94
 protection of children B179
 protection technologies B188
 protection, environmental B71
 protection, health B201
 protection, legal B25
 protective equipment B152
 protective mechanism B144
 protein-low carbohydrate diet, high B39
 proteins B187
 proteus, amoeba B224
 protocols, laboratory 10
 protoporphyrin, zinc B223
 provide, failure to B34
 provider B45
 providers, health care B63,B123
 provincial inspectors B212
 provocation tests B159
 provocation, nasal B186
 proximal individual social environment B176
 pseudo-retardation, allergic B127
 psychiatric B119
 psychiatric disorders A72,B45,B52,B72
 psychiatric experts B130
 psychiatric facilities B233
 psychiatric facility B234
 psychiatric hospitals B22
 psychiatric illness B190
 psychiatric institutions A75,B233
 psychiatric labels A43,B95
 psychiatric literature B211
 psychiatric patient A43,B112
 psychiatric patients B34
 psychiatric sequelae B72
 psychiatric services B86

- psychiatrists B79
 psychiatry B147
 psychiatry, community B79
 psychiatry, public image of B129
 psycho-social B100
 psychoanalysis B211
 psychoanalytic B211
 psychogenic illness, mass B138
 psychological B98,B230,B233
 psychological adaptations B147
 psychological aspects of the arms race B74
 psychological association B60
 psychological benefits B39
 psychological development B21,B183
 psychological difficulties B122
 psychological distress A69,B47,B66
 psychological disturbances A66
 psychological effects B42,B74,B123
 psychological function tests B83
 psychological functioning B71
 psychological functions B123
 psychological health B34
 psychological injury B79
 psychological needs B78
 psychological reactions B65
 psychological response A60,B230
 psychological stress A10,A64,B53,B54
 psychological suffocation A47,B2
 psychological support B112
 psychological test B170
 psychological tests B54
 psychological variables A65
 psychological well-being B176
 psychology B8,B163,B229
 psychology, behavioural B162
 psychology, developmental B118
 psychology, environmental B65,B107,B233
 psychometric techniques B223
 psychosis B160
 psychosocial orientation B183
 psychosocial problems B140
 psychosocial situations, stressful B122
 psychosocial stress B241
 psychosocial stressors B17
 psychosomatic diseases B152
 psychosomatic disorders, somatic and B152
 psychosomatic illness B81
 psychotic reactions B23
 psychotropic drugs B22
 public access buildings B73
 public actions B139
 public attention, deleted from B58
 public attitudes B51
 public awareness B83
 public buildings A19,B43,B54,B209
 public education 19,B216
 public environments 5
 public expectations for health care B63
 public health B28,B37,B57,B71,B101,B103,B120,B136,
 B173,B202,B238
 public health policies B138
 public health policy B22,B37,B57,B60,B83,B96,B100,
 B120,B122,B204,B224,B229
 public health problems B179
 public health, community-based B82
 public housing B232
 public identification A47
 public image of psychiatry B129
 public input A21
 public life, integrity in B164
 public participation B9,B119,B120
 public places 3,A19,B220
 public policy B8,B9,B33,B34,B35,B39,B42,B45,B49,B57,
 B58,B63,B82,B85,B87,B90,B93,B94,B95,B96,B97,B98,B103,
 B112,B115,B118,B119,B120,B121,B127,B129,B133,B134,
 B137,B140,B144,B148,B151,B156,B176,B179,B183,B188,
 B191,B196,B197,B200,B201,B204,B206,B209,B212,B216,
 B217,B221,B222,B228,B232,B237,B239
 public policy alternatives B203
 public policy issues B36,B144,B236
 public policy, healthy B83
 public spaces 6
 public spaces, affection in A49
 public, general B8
 publications, mental health B51
 publicness B148
 pulmonary B174
 pulmonary disease A53,B174,B187,B188
 pulmonary effects B191
 pulmonary function B5,B11,B109,B120,B129,B204,B212
 pulmonary functions B121
 pulmonary tuberculosis B178
 pulp and paper mill effluent B208
 pulse modulations B28
 pulse rate B44
 pungent chemicals B209
 punishment A26
 punishment of a child B34
 pupils of the eyes A60,B230
 purposes, peaceful B90
 put-down B81
 PVCs B191
 pyrite B181
 qualities, aesthetic B162
 qualities, institutional B235
 quality B63
 quality complaints, air B212
 quality management, air B136
 quality of care B156
 quality of life B57,B95,B122,B203
 quality of urban life B9,B163
 quality, water B70
 quantitative analysis B225,B226
 quantitative electrophysiology B108
 quarters, living B42
 Quebec, frenchspeaking B64
 quo, status B26
 race 3,A2,A38,A80,B80,B218
 race relations B64
 race, arms A66,B74
 race, psychological aspects of the arms B74
 racial B50,B243
 racial discrimination B44,B109,B233
 racial exclusiveness B183
 racial stereotypes, reading B243
 racially-integrated suburban neighbourhoods B233
 racism 17,A38,A46,B64,B79
 racism scale 13
 racism, police B64
 racist conduct B79
 radar operators B186
 radiation A24,A32,B60,B75,B121,B209
 radiation exposure levels B20
 radiation exposures B63
 radiation hazards, vdt B208
 radiation, electromagnetic B16,B28,B97,B102,B132
 radiation, high-frequency A62
 radiation, ionizing A62,B3,B20
 radiation, low-level A16,B69
 radiation, non-ionizing B102
 radiation, ultraviolet B30,B178
 radiation-shielded fluorescent fixtures A61
 raditions B128
 radical mastectomies B136
 radio B186
 radioactive clouds B61
 radioactive material A39,B208

- radioactive materials A62
radioactive waste B61
radioactive wastes A67,B61
radioactivity, natural B221
radon A6,A15,A54,B11,B19,B20,B28,B43,B46,B57,B92,
B104,B121,B123,B132,B136,B141,B142,B146,B104,B203,
B210,B211,B217,B221
radon barrier B46
radon daughter nuclides A53
radon daughter products A54
radon daughters B89
radon decay products B89
radon pollution B88,B181
radon variability B181
radon, airborne B88
radon-222, waterborne B88
radon operators B186
railroad workers B223
railway workers B15
rain, acid A22,A56,B70,B116,B155,B216,B217
raised moles B62
raising, consciousness B85
ramp A42,B211
range of competencies B202
range of vulnerability 5,19,A17
range of vulnerability to pollutants B199
range restrictions, parentally-defined B84
range, gas-fired B219
range, power frequency A62
rape crisis centres A34,B35
rape, victims of B13
rashes, skin B80
rate, emission B128
rate, heart A53,B10,B146
rate, metabolic A24
rate, nasal inspiratory flow B186
rate, pulse B44
rate, ventilation B73,B158,B174
rates, cancer A58
rates, exposure B174
rates, high injury B114
rates, higher admission B152
rates, hospital admission A69
rates, infiltration B20
rates, local unemployment B140
rates, mental hospital admission B112
rates, mortality B191
rates, socioeconomic differences in smoking B145
rates, special tax B149
rates, ventilation B3,B210,B227
ratings, aggregate co-worker B176
rats A14
rats, pregnant B59
raynaud's disease A56,B171
rays, gamma A62
re-organizing worktime B196
reaction time A15,B7,B235
reaction, adverse B169
reaction, aero-allergenic B187
reactions B170,B195
reactions to burglary, people's B113
reactions to food B171
reactions, adverse drug B86
reactions, allergic 12,B3,B16,B154
reactions, allergic skin A55
reactions, asthmatic A13
reactions, biological chain B90
reactions, cognitive B168
reactions, depressive A58,B72
reactions, emotional B90
reactions, human B11
reactions, nervous system B161
reactions, photoxidation B97
reactions, psychological B65
reactions, psychotic B23
reactions, skin B172
reactions, sulfite-related adverse B240
reactions, urticarial A13,B14
reactive chemistry B155
reactivity 5,8
reactivity to mental stress, cardiovascular B146
reactivity, airway B38
reading A26,A52,A60
reading comprehension A8,B22,B150
reading levels B77
reading, decreased interest in B243
real estate brokers B233
real-life experiences A10
realities, contradictory B183
rearrangement of worktime B196
reasonable claim B79
reasoning A52,B187
rebates, tuition B149
recirculated B177
recirculation of return air B19
recirculation system A11,B212
recognition A27,A32,A67,B9,B219
recognition, lack of B209
recordings, programmable ambulatory
electrocardiograph B191
records, population-at-risk B129
recreation B243
recreational areas B65
recreational waters B21
recurrent bronchitis B169
recurrent respiratory infection B143
recycled air B165
red blood synthesis B116
redress of grievances B139
reduced cleaning budgets B76
reduced life options B134
reduced ventilation B19,B172
reduced ventilation, residence B19
reducing airborne contaminants B145
reducing inequalities B196
reducing inequities B142
reducing risks to children B209
reduction in lung function B94
reduction of stress B37
reduction, exposure B6
reduction, noise B231
reduction, source B27
reductions in motor skill B56
reductions, annual job hour B149
reentry B172
refined foods B189
reflection, independent B162
reflexes, abdominal A9
refrigerants A5,B166
refrigerator B171
refuge A43,B95
regenerating community B133
region, york B113
regional geological factors A6
regional geology B181
regions, fruitgrowing A58,B72
regions, temperate B232
regression analysis B204
regression to infantile behaviour B90
regulation B116
regulations, occupational B15
regulations, safety B24
regulatory B33
regulatory agencies B185,B201,B206
regulatory mechanisms, hormonal B2
rehabilitation A43,B95,B178,B181
rehabilitation systems B156
rehabilitation, community-based A44
reinvestment, community B156
rejection A46,B211

- related accidents, building B114
relation, dose-response B140
relations with supervisors B206
relations, police/community B41
relations, race B64
relations, social B182
relationship, causal B243
relationship, dose-response B10
relationship, pathology of human B25
relationships B40,B57
relationships with others B114
relationships, close A35,B99
relationships, dose-response B65,B120,B124,B174
relationships, equal A35
relationships, exposure-response B210
relationships, hospital social B233
relationships, indoor-outdoor B54
relationships, nature of B44
relationships, oppressive A34
relationships, parenting B45
relationships, people/ environmental B235
relationships, personal A26,B40
relationships, social A72,B87,B230
relationships, spatial B69
relationships, strained personal B40
relative humidity A63
relatives B216
release of heat B222
relevant disciplines B91
reliance, self B119
relief centres, pollution A11
religion 17,A38,A80
religious B100
remedial classes A70,B45
remedial environmental B128
remedial measures B210
remote areas B216
removal B25
removal efficiency B21
remover, paint B207
removers, paint A41
rendering, colour A10
renewable energy B237
renovated A9
renovating buildings B9
renovation B222
renovations B96
rental housing B233
rental housing crisis B62
rental properties A36,B63
renter B37
repeal of undesirable legislation B139
repertoires, as A72
repertoires, coping B55
repetitive B44
report, the epp B133
repose B183
representation B35
repression B90
reproduction processes A8
reproductive hazards A15,B38,B193
reproductive information B129
reproductive outcomes, deleterious B192
requirements, admission A38,B44
requirements, child's functional B122
requirements, special environmental B239
research B85,B98,B112,B144,B197
research and development, scientific B222
research inventory B136
research involving children B144
research priorities B3,B31,B35,B37,B51,B52,B55,B61,
B65,B69,B82,B85,B101,B111,B114,B117,B120,B123,B143,
B153,B155,B157,B163,B179,B187,B191,B199,B203,B206,
B209,B214,B221,B229,B232,B237
research problems B91
research procedures B36
research programs, medical B91
research projects B152
research study B203
research subjects A28,B144
research, behavioral B144
research, behavioural B234
research, brain B108
research, canadian centres of B91
research, danish B139
research, environmental B26
research, health B228
research, intervention B36
research, medical B90,B91
research, national centres of B90
researchers, design B26
researchers, health B109
reserve, indian B180
reserve, serpent river indian B180,B208
reserves A38,B191
residence B141,B227
residence reduced ventilation B19
residences B74,B198,B219
residential A5,B65,B79,B132,B163,B177,B231
residential access B211
residential accidental death B4
residential areas B23
residential buildings A74,B19,B57,B184,B242
residential crowding B183
residential density B183
residential design B32,B114
residential environment B231
residential environments, mixed-use B183
residential exposure guidelines 13
residential histories B122
residential neighborhood B36
residential safety B114,B161,B228
residential settings B106
residential treatment settings B234
residents B94
residents, urban B203,B220
residue, detergent A19
residue, dried detergent B115
residue, drug A57,B161
residue, fertilizer B72
residues of chemicals B215
residues, pesticide A82
resin A13
resin binders, formaldehyde-based B106
resin compounds B14
resin-bonded A6
resins B6
resins, acrylic B67
resins, vinylidene chloride B67
resistance A46,A68,B2,B60,B143
resistance of the skin A60,B230
resistance to infection B17,B60
resistance, airway B5,B6,B153
resistance, community A7,B78
resistance, individual B37
resistance, nasal B5
resistance, skin B44
resistance, thermal B25
resistance, tissue B119
resistant, unusually B45
resources, community B127
resources, coping A65,B32
resources, emotional B33
resources, environmental A1
resources, financial B33
resources, lack of B122,B167
resources, limited B142
resources, military A35
resources, non-renewable A22,B222
resources, personal B106

-
- radioactive materials A62
 - radioactive waste B61
 - radioactive wastes A67,B61
 - radioactivity, natural B221
 - radon A6,A15,A54,B11,B'9,B20,B28,B43,B46,B57,B92,
B104,B121,B123,B132,B136,B141,B142,B146,B184,B203,
B210,B211,B217,B221
 - radon barrier B46
 - radon daughter nuclides A53
 - radon daughter products A54
 - radon daughters B89
 - radon decay products B89
 - radon pollution B88,B181
 - radon variability B181
 - radon, airborne B88
 - radon-222, waterborne B88
 - radon operators B186
 - railroad workers B223
 - railway workers B15
 - rain, acid A22,A56,B70,B116,B155,B216,B217
 - raised moles B62
 - raising, consciousness B85
 - ramp A42,B211
 - range of competencies B202
 - range of vulnerability S,19,A17
 - range of vulnerability to pollutants B199
 - range restrictions, parentally-defined B84
 - range, gas-fired B219
 - range, power frequency A62
 - rape crisis centres A34,B35
 - rape, victims of B13
 - rashes, skin B80
 - rate, emission B128
 - rate, heart A53,B10,B146
 - rate, metabolic A24
 - rate, nasal inspiratory flow B186
 - rate, pulse B44
 - rate, ventilation B73,B156,B174
 - rates, cancer A58
 - rates, exposure B174
 - rates, high injury B114
 - rates, higher admission B152
 - rates, hospital admission A69
 - rates, infiltration B20
 - rates, local unemployment B140
 - rates, mental hospital admission B112
 - rates, mortality B191
 - rates, socioeconomic differences in smoking B145
 - rates, special tax B149
 - rates, ventilation B3,B210,B227
 - ratings, aggregate co-worker B176
 - rats A14
 - rats, pregnant B59
 - raynaud's disease A56,B171
 - rays, gamma A62
 - re-organizing worktime B196
 - reaction time A15,B7,B235
 - reaction, adverse B169
 - reaction, aero-allergenic B187
 - reactions B170,B195
 - reactions to burglary, people's B113
 - reactions to food B171
 - reactions, adverse drug B86
 - reactions, allergic 12,B3,B16,B154
 - reactions, allergic skin A55
 - reactions, asthmatic A13
 - reactions, biological chain B90
 - reactions, cognitive B168
 - reactions, depressive A58,B72
 - reactions, emotional B90
 - reactions, human B11
 - reactions, nervous system B161
 - reactions, photoxidation B97
 - reactions, psychological B65
 - reactions, psychotic B23
 - reactions, skin B172
 - reactions, sulfite-related adverse B240
 - reactions, urticarial A13,B14
 - reactive chemistry B155
 - reactivity 5,8
 - reactivity to mental stress, cardiovascular B146
 - reactivity, airway B38
 - reading A26,A52,A60
 - reading comprehension A8,B22,B150
 - reading levels B77
 - reading, decreased interest in B243
 - real estate brokers B233
 - real-life experiences A10
 - realities, contradictory B183
 - rearrangement of worktime B196
 - reasonable claim B79
 - reasoning A52,B187
 - rebates, tuition B149
 - recirculated B177
 - recirculation of return air B19
 - recirculation system A11,B212
 - recognition A27,A32,A67,B9,B219
 - recognition, lack of B209
 - recordings, programmable ambulatory
electrocardiograph B191
 - records, population-at-risk B129
 - recreation B243
 - recreational areas B65
 - recreational waters B21
 - recurrent bronchitis B169
 - recurrent respiratory infection B143
 - recycled air B165
 - red blood synthesis B116
 - redress of grievances B139
 - reduced cleaning budgets B76
 - reduced life options B134
 - reduced ventilation B19,B172
 - reduced ventilation, residence B19
 - reducing airborne contaminants B145
 - reducing inequalities B196
 - reducing inequities B142
 - reducing risks to children B209
 - reduction in lung function B94
 - reduction of stress B37
 - reduction, exposure B6
 - reduction, noise B231
 - reduction, source B27
 - reductions in motor skill B56
 - reductions, annual job hour B149
 - reentry B172
 - refined foods B189
 - reflection, independent B162
 - reflexes, abdominal A9
 - refrigerants A5,B166
 - refrigerator B171
 - refuge A43,B95
 - regenerating community B133
 - region, york B113
 - regional geological factors A6
 - regional geology B181
 - regions, fruitgrowing A58,B72
 - regions, temperate B232
 - regression analysis B204
 - regression to infantile behaviour B90
 - regulation B116
 - regulations, occupational B15
 - regulations, safety B24
 - regulatory B33
 - regulatory agencies B185,B201,B206
 - regulatory mechanisms, hormonal B2
 - rehabilitation A43,B95,B178,B181
 - rehabilitation systems B156
 - rehabilitation, community-based A44
 - reinvestment, community B156
 - rejection A46,B211

-
- related accidents, building B114
 - relation, dose-response B140
 - relations with supervisors B208
 - relations, police/community B41
 - relations, race B64
 - relations, social B182
 - relationship, causal B243
 - relationship, dose-response B10
 - relationship, pathology of human B25
 - relationships B40,B57
 - relationships with others B114
 - relationships, close A35,B99
 - relationships, dose-response B65,B120,B124,B174
 - relationships, equal A35
 - relationships, exposure-response B210
 - relationships, hospital social B233
 - relationships, indoor-outdoor B34
 - relationships, nature of B44
 - relationships, oppressive A34
 - relationships, parenting B45
 - relationships, people/ environmental B235
 - relationships, personal A28,B40
 - relationships, social A72,B87,B230
 - relationships, spatial B69
 - relationships, strained personal B40
 - relative humidity A63
 - relatives B216
 - release of heat B222
 - relevant disciplines B91
 - reliance, self B119
 - relief centres, pollution A11
 - religion 17,A38,A80
 - religious B100
 - remedial classes A70,B45
 - remedial environmental B128
 - remedial measures B210
 - remote areas B216
 - removal B25
 - removal efficiency B21
 - remover, paint B207
 - removers, paint A41
 - rendering, colour A10
 - renewable energy B237
 - renovated A9
 - renovating buildings B9
 - renovation B222
 - renovations B96
 - rental housing B233
 - rental housing crisis B62
 - rental properties A36,B63
 - renter B37
 - repeal of undesirable legislation B139
 - repertoires, as A72
 - repertoires, coping B55
 - repetitive B44
 - report, the epp B133
 - repose H183
 - representation B35
 - repression B90
 - reproduction processes A8
 - reproductive hazards A15,B38,B193
 - reproductive information B129
 - reproductive outcomes, deleterious B192
 - requirements, admission A38,B44
 - requirements, child's functional B122
 - requirements, special environmental B239
 - research B85,B98,B112,B144,B197
 - research and development, scientific B222
 - research inventory B136
 - research involving children B144
 - research priorities B3,B31,B35,B37,B51,B52,B55,B61,
B65,B69,B82,B85,B101,B111,B114,B117,B120,B123,B143,
B153,B155,B157,B163,B179,B187,B191,B199,B203,B206,
B209,B214,B221,B229,B232,B237
 - research problems B91
 - research procedures B36
 - research programs, medical B91
 - research projects B152
 - research study B203
 - research subjects A28,B144
 - research, behavioral B144
 - research, behavioural B234
 - research, brain B108
 - research, canadian centres of B91
 - research, danish B139
 - research, environmental B26
 - research, health B228
 - research, intervention B36
 - research, medical B90,B91
 - research, national centres of B90
 - researchers, design B26
 - researchers, health B109
 - reserve, indian B180
 - reserve, serpent river indian B180,B208
 - reserves A38,B191
 - residence B141,B227
 - residence reduced ventilation B19
 - residences B74,B198,B219
 - residential A5,B65,B79,B132,B165,B177,B231
 - residential access B211
 - residential accidental death B4
 - residential areas B23
 - residential buildings A74,B19,B57,B184,B242
 - residential crowding B183
 - residential density B183
 - residential design B32,B114
 - residential environment B231
 - residential environments, mixed-use B183
 - residential exposure guidelines 13
 - residential histories B122
 - residential neighborhood B36
 - residential safety B114,B161,B228
 - residential settings B106
 - residential treatment settings B234
 - residents B94
 - residents, urban B203,B220
 - residue, detergent A19
 - residue, dried detergent B115
 - residue, drug A57,B161
 - residue, fertilizer B72
 - residues of chemicals B215
 - residues, pesticide A82
 - resin A13
 - resin binders, formaldehyde-based B106
 - resin compounds B14
 - resin-bonded A6
 - resins B6
 - resins, acrylic B67
 - resins, vinylidene chloride B67
 - resistance A46,A68,B2,B60,B143
 - resistance of the skin A60,B230
 - resistance to infection H17,B60
 - resistance, airway U5,U6,U153
 - resistance, community A7,B78
 - resistance, individual B37
 - resistance, nasal B5
 - resistance, skin B44
 - resistance, thermal B25
 - resistance, tissue B119
 - resistant, unusual B45
 - resources, community B127
 - resources, coping A65,B22
 - resources, emotional B33
 - resources, environmental A1
 - resources, financial H33
 - resources, lack of B122,B167
 - resources, limited B142
 - resources, military A35
 - resources, non-renewable A22,B222
 - resources, personal B106

- resources, personnel financial B122
 respect A27,A49,A71,B220
 respect for persons B144
 respect, lack of A28,B162
 respirable particles B42
 respirable particulate mass B218
 respirable particulate, mass B65
 respirable particulates A53
 respirable sulfates B54
 respirable suspended particulates B93,B173
 respiratory B154,B158
 respiratory allergic disorders A13
 respiratory allergic disorders, occupational B159
 respiratory conditions, acute B24
 respiratory disease B51,B67,B109,B156,B159,B173,B231,
 B241
 respiratory diseases A20,A63,B41,B80,B90,B119,B185
 respiratory disorders A51,B143
 respiratory effects B197
 respiratory function B93,B105,B210
 respiratory function tests B204
 respiratory hazards B123
 respiratory health B24,B73,B77,B92,B107,B116,B184
 respiratory health hazards B123
 respiratory illness B86,B108
 respiratory indices B41
 respiratory infection B60
 respiratory infection, recurrent B143
 respiratory infections B104
 respiratory irritant B14
 respiratory irritation A16,B90,B115
 respiratory irritation, upper B30
 respiratory problems B29,B209
 respiratory response B147
 respiratory sensitization B31
 respiratory symptoms A8,B86,B108
 respiratory symptoms in children B108
 respiratory system A75,B36,B228
 respiratory tract A52,B5,B14,B16,B160,B187
 respiratory tract defense mechanisms B187
 respiratory tract diseases B202
 respiratory tract infections B212
 respiratory tract, upper B131
 respiratory virus infections B178
 response patterns B54
 response to noise, sociological B230
 response, antibody A83,B52
 response, community B23
 response, immune A64,B40
 response, multiorgan B125
 response, placebo B50
 response, psychological A60,B230
 response, respiratory B147
 response, sensitivity B200
 response, variability of A84
 response, windows of frequency B200
 responses to stressors, delayed B40
 responses, affective B55
 responses, allergic B51,B90,B215
 responses, behavioural B160
 responses, cell-mediated B232
 responses, delayed A64
 responses, idiosyncratic nature of individual B232
 responses, immunologic B18
 responses, irritant B221
 responses, physiological B230
 responses, policy B185
 responsibilities B143
 responsibilities, burden of home B209
 responsibilities, family A34,A35,A36,B45,B219
 responsibilities, household B34
 responsibility B23,B58,B202,B236
 responsibility for domestic life B183
 responsibility, community A78
 responsibility, consumer B8
 responsibility, equitable B222
 responsibility, indirect B204
 responsibility, individual B133
 responsibility, personal B8
 responsible action B162
 responsiveness, decreased immunological B232
 responsiveness, levels of B232
 restaurant food A57,B240
 restricted information 13
 restrictions on behaviour B44
 restrictions, parentally-defined range B84
 resumes B112
 resurfacing machines, ice B7
 resuspension of precipitated particles B165
 retardants, flame A55,B14
 retardation B64
 retardation, mental A52,B187,B243
 retarded B81
 retarded, mentally B110
 retirement A30,B204
 retrofitting B91
 return air, recirculation of B19
 revealing diversity B26
 reversals, economic B138
 review B52
 review, product B75
 revival, fatherhood B45
 reward structure B182
 rewards, financial B134
 rezoning A11
 rheumatoid arthritis A76,B49,B50,B171
 rhinitis A53,A74,B31,B141,B169,B173
 rhinitis, allergic B186
 rhinitis, perennial B186
 rhythms B84
 rhythms, physiological A61
 rickets A61,B30,B84
 right to shelter B78
 right-to-know A81
 right-to-know legislation B104,B221
 rights and freedoms, charter of B218
 rights code, ontario human B41
 rights codes, human A47,B112
 rights issues for, human B2
 rights legislation, human A79,A80
 rights of non-smokers B144
 rights of women A35
 rights policy, human B97
 rights statutes, civil A79
 rights, basic human B222
 rights, children's B25,B212
 rights, civic A49,B139
 rights, civil A49,B139
 rights, custody B86
 rights, employee A18
 rights, equality of B218
 rights, gay A47,B41,B139
 rights, human 17,A22,B97,B218
 rights, minority A80
 rights, political A49,B139
 rights, property B188
 rights, spousal A47,B2
 rights, violations of 11
 rights, women's B13
 rigid B36,B224
 rigid environmental controls B170
 rigid no-noise code B231
 rigidities, destructive B149
 rink, ice B7
 rise living, high B183
 risk B4,B30,B231
 risk assessment B21,B123,B168,B201
 risk assessment processes B237
 risk factors B30,B71,B137,B233
 risk groups B124

- risk groups, high A41,B161
risk of accidents B198
risk of death A70
risk of death, increased B80
risk, assessment of B200
risk, cancer B59
risk, carcinogenic B173
risk, childhood B209
risk, children at B202
risk, children's B165
risk, leukemia B186
risk-taking B85
risks B237
risks to children, reducing B209
risks to nonsmokers B175
risks, health B70,B220,B221
risky driving B61
river indian reserve, serpent B180,B208
road crashes B130
road oil B223
road safety education B202
road vehicles A50,B179
road-crash A25,B130
roadways A20
role ambiguity B44
role at work B44
role conflict B44
role of language, the B26
role of medicine B142
role of women, economic B218
role, feminine B143
role, oppressor B99
role, sick B143
roles B183
roles of children, participatory B25
roles of citizen and community B133
roles, decision-making B35
roles, sex B84
roles, social B143
roller coaster B138
roofs, flat A9,A54,B76
room, dust-free B242
rooms, changing A12
rooms, music A12
rooms, windowless A61
roots of ageism B25
roots, grass A21
route, airborne B104
routine B235
rubber, sponge A5
runner, distance B49
runners, distance A20
running B49
running nose A6
rural areas B148
rural communities B86
saccharin B116
safe environments B83
safeguards A78,B239
safer environments B147
safer sex A47,A48,B2,B4
safer stair design B114
safety A50,A66,B10,B52,B144,B183,B185,B192,B193,B198,
B202
safety and health, occupational B192
safety awareness B86
safety campaigns B193
safety committees, joint health and B185
safety conditions B177
safety education A50
safety education, road B202
safety hazards B37
safety limits B10
safety margin B228
safety problem B130
safety procedures B103
safety programs B185
safety regulations B24
safety standards B92
safety underground B49
safety, child health and B63,B130
safety, health and 3,6,B33,B123
safety, home B53,B114
safety, margin of B228
safety, occupational B24
safety, occupational health and A1,A34,B3,B33,B38,
B44,B73,B123,B127,B128,B134,B135,B157,B165,B180,B185,
B190,B191,B208,B221,B235,B241
safety, occupational B57
safety, residential B114,B161,B228
safety, school health and A1,A12,A81
safety, stair B157
safety, tenant health and A1,A81
safety, traffic A27
safety, workplace health and A15,B239
safety-related behaviour B192
safetyproofing, home B53
safety B209
salicylate, monosodium B164
saliva A60,B230
salts A52,B241
salts, metal A55
same-gender lovers A30
same-gender sexual practices A46
sample, grab B141
sampler, domestic air pollution B93
sampler, passive B148
samplers, personal B196
samples, breast milk B136
sampling B15
sampling devices, air B184
sampling, passive B190
sanction, legal B185
sanctions, legal B220
sanitation A21,B12
sanitation services B229
satisfaction, job B240
satisfaction, low job B44
satisfaction, occupants B75
satisfactions B107
sauna B180
saving, energy B11
scale A75
scale, anti-gay 13
scale, human B119,B229
scale, racism 13
scale, sexism 13
scarce accommodation B63
scarcity of work B151
scented A5
scented cleaning products B166
scented cosmetics A44
schedule, time B182
schedules, work B151
schizophrenia B160
schizophrenic A58,B72
scholastic performance A10,B24,B166
scholastic supplies B22
school A8,B50,B93,B105,B162,B167,B209,B213,B229
school achievement B213
school activities B243
school air B56
school air intakes A11
school boards A76
school building, community B43
school buildings A11
school buildings, allergens in B113
school buses A9,B22,B177
school children B92,B107,B109,B137,B158,B214
school children around B41

- school children, elementary B222
 school curricula A26
 school curricula, medical A49,B16
 school environment B84
 school environments B149
 school health and safety A1,A12,A81
 school performance B64
 school sites B75
 school students, elementary A60
 school students, high B105
 school, community B42
 school, elementary B158,B177
 school, secondary B177
 school, skipping B50
 school-safe A11
 school-safe standard B226
 schoolchildren B77,B159,B225
 schooling, compulsory A10,B162
 schooling, growing without B92
 school-related problems B112
 schools 3,5,6,19,B13,B22,B24,B25,B43,B54,B71,B76,B84,
 B96,B113,B150,B154,B158,B164,B177,B188,B189,B198,
 B205,B227,B235
 schools, canadian medical B91
 schools, elementary B242
 schools, medical B204
 science B93
 science centres, health B91
 science classrooms B24
 science laboratories A12,B13,B177
 science materials A8,B199
 science, aerosol B242
 science, behavioral B71
 science, contemporary B91
 science, medical B90
 sciences, behavior B59
 sciences, environmental health B229
 scientific base B91
 scientific research and development B222
 scientific understanding B163
 scientists B91,B177
 scientists, behavioral B163
 scientists, social B9
 scintillation, liquid B88
 sclerosis, multiple A9,B87
 screening A82,B20,B187
 screening strategies, lead B20
 sculpting A5
 sealants B6
 sealed buildings B171,B206,B217
 sealed office buildings A16,B35,B207
 sealed office complex, hermetically B18
 sealed, hermetically B206
 sealing of homes B215
 search for shelter, the B78
 seating, vinyl B195
 second hand smoke B42,B125
 second-class corporate citizens B39
 second-hand smoke B132
 second-hand tobacco smoke B72
 secondary school B177
 secreted A49
 secretions, gastric A60,B230
 sector, discrimination in the private B41
 sector, private B91
 sectors, excluded B25
 sectors, private B34
 secure A7
 security A30,A37,A43,A66,B25,B42,B95,B148
 security, environmental B184
 security, income B85
 sedentary activity B17
 segregate B156
 segregated A44
 segregation A32,B208
 segregation of smokers H207
 segregation, age A10,B162
 segregation, geographic B182
 seizures A57,B23,B240
 selection guidelines, food B142
 selective vulnerability B157
 selenium A56
 self confidence, low B44
 self esteem, low B44
 self reliance B119
 self-care B82,B133
 self-care movement A77
 self-care, individual A68
 self-contained dwelling B32
 self-defeating A79
 self-directed change B109
 self-ego B118
 self-esteem 7,17,A2,A26,A46,A66,A67,A79,B99,B173,
 B184
 self-esteem, loss of 13,B152
 self-esteem, low A33,B211
 self-generated B162
 self-help development B39
 self-help systems B85
 self-identity A66,B111,B184
 self-interested approaches B34
 self-leveling cement B82
 self-reliance A67,B162
 self-sufficiency B45
 self-tailored environments A1
 self-worth A43
 self-worth, low B95
 self-identity A31
 sellers B233
 semi-ambulant A42
 semirural communities A72,B68
 senior citizens B90,B204
 senior citizens apartment B106
 senior citizens apartments A6
 seniors A28
 sense of control B111
 sense of home B32
 sense of power A67
 senses, human A55
 sensibilities, acoustic B231
 sensitive areas B70
 sensitive children, chemically B150,B188,B189
 sensitive individuals 10,13
 sensitive individuals, mold B29
 sensitive minority B140
 sensitive patients, chemically B171
 sensitive populations B5
 sensitive students A11
 sensitive sub-population B121
 sensitive subjects B143
 sensitive, chemically A62
 sensitive, electromagnetically A62,B171
 sensitivities 5,B150
 sensitivities to moulds A77
 sensitivities, chemical B47,B167
 sensitivities, food B48,B111
 sensitivities, food or chemical B168
 sensitivities, hidden B125
 sensitivities, individual B237
 sensitivity B30,B34,B63,B169,B197,B203
 sensitivity of the fetus B64
 sensitivity of the population B124
 sensitivity response B200
 sensitivity to odours A74,B169
 sensitivity, bronchial B153
 sensitivity, chemical B50,B53,B111,B138,B149,B150,
 B160,B167
 sensitivity, environmental A44,B27,B117,B170
 sensitivity, food B150
 sensitization A17,A53,B14,B146,B154,B169

- sensitization, allergic B97
sensitization, chemical B97
sensitization, respiratory B31
sensitization, smooth muscle A83,B169
sensitized A6,A55
sensitized to formaldehyde B6
sensitizer B80
sensitizers B190
sensitizing agent B228
sensitizing chemicals B3
sensory irritants B62
sensory irritation, human B227
sensory skill, motor B235
sensorymotor disabilities B108
sequelae, psychiatric B72
sequelae, subtle neurologic B192
serious physical ailments B47
serotonin A63,B110
serpent river indian reserve B180,B208
serum cholesterol A70,B152
serum cholesterol level, elevated B137
serve B211
service jobs A32
service level B216
service needs, changing human B224
service systems, formal B156
service workers B208
service, social B86
services, access to B86
services, availability of B144
services, delivery of B86
services, equitable access to B8
services, funeral A47,B55
services, gaps in B122
services, health B47,B86
services, home care B7
services, housing B4
services, managed health care B142
services, mental health A33,B95
services, nation's health care B91
services, preventive B131
services, psychiatric B86
services, sanitation B229
services, social B86,B120
services, support B71,B119
setting A4
setting, imposed B167
setting, institutional B81
setting, natural B37
setting, new life B35
setting, non-industrial B3
setting, physical B1,B164
settings A1
settings, commercial A16
settings, community B148
settings, decay of physical B163
settings, environmental B163
settings, industrial A13
settings, institutional B235
settings, natural A22
settings, organizational A72,B105
settings, residential B106
settings, residential treatment B234
settings, small town B148
settings, social B163
settings, tertiary care B233
settlements A19
severe mental disabilities B219
sewage sludge incinerators B126
sewage treatment facilities B21
sex and law in Canada B112
sex differences B84,B143
sex differences in illness B143
sex discrimination A33
sex education B100
sex roles B84,B182
sex, gay B4
sex, safer A47,A48,B2,B4
sexism 12,17,19,A32,A33,B40,B99,B134,B136,B211
sexism scale 13
sexism, cause of B13
sexism, elimination of B13
sexist actions A35
sexist behaviour A35
sexrole stereotypes A26,B243
sexual abuse B13,B34,B134
sexual assault A33,B34
sexual decision-making B173
sexual harassment A34,B35
sexual harassment 12
sexual molestation A26
sexual orientation 6,13,17,A46,A47,A80,B26,B41,B112,
B156
sexual partners B201
sexual practices, same-gender A46
sexually compulsive A35,B99
sexually transmitted disease B131
sexually transmitted diseases 13,A47,B2,B173
shampoos, carpet B115
sharing, job A18,A36,B45,B93
shavings, wood A53,B5
shell, building B222
shelter A43,A67,B78,B95,B119
shelter policies B149
shelter, dignified B78
shelter, right to B78
shelter, the search for B78
shelter, transitional B78
sheltered A80
sheltered-workshop mentality B112
shelters A34,B101,B134
shelters for battered women B35
shelving B106
shielding A62,B128,B154
shift, frequently changing B28
shifts, affective B108
shock B112
shock, anaphylactic A57,B240
shoes B38
shops, automotive A12
shops, woodworking A12,B13
short attention span B167
shortage, housing B78
shortages, housing A36
shorter A70,B218
shortness of breath A6
shortness of breath B50
shower A6
shower heads A54,A59,B21
showering B165
shrinkage B25
sick B151
sick building syndrome A9,A17,A82,B3,B35,B73,B76,
B140,B153,B177,B237,B238
sick role B143
side effects B186
side effects of innovations A65
side effects, unhealthy B220
sidestream cigarette smoke B73
sidewalks B213
sign, early warning A74
signals, electromagnetic B17
similarities B26
single B122
single mother B62
single-parent families A38
single-parent households B194
sinusitis A74,B169
sinusoidal vibration B100
sites, construction B134

- sites, school B75
 situations, stressful psychosocial B122
 size B1
 size, particle B184,B233
 sizing agents A55,B14
 skating B7
 skeletal-muscle tension A60,B230
 skill deficits B150
 skill, motor sensory B235
 skill, reductions in motor B56
 skilled B34
 skills B44
 skills development A26
 skills, auditory memory B150
 skills, coping A64,B54,B71
 skills, environmental B202
 skills, life A44,B216
 skills, social A45
 skin allergies B215
 skin cancer A61
 skin colour I3
 skin conditions B209
 skin dryness B131
 skin rashes B80
 skin reactions B172
 skin reactions, allergic A55
 skin resistance B44
 skin, resistance of the A60,B230
 skipping school B50
 sleep A6,A60,B231
 sleep patterns B80
 sleep, noise on B230
 sleeplessness B80,B131
 sleepy B138
 slip and fall injuries B211
 slip-resistant surfaces B10
 slips and falls B59
 slow learners A8,A70,B45
 sludge incinerators, sewage B126
 sluggishness A9
 small town environments B68
 small town settings B148
 small towns B148
 smelter, lead B190,B223
 smog, photochemical B83
 smoke B41,B184,B228
 smoke, cigarette A74,B83,B169,B217,B220
 smoke, second hand B42,B125,B132
 smoke, second hand tobacco B72
 smoke, sidestream cigarette B73
 smoke, tobacco A8,A14,A15,A16,A53,B29,B42,B43,B55,
 B62,B74,B86,B125,B137,B173,B174,B175,B176,B199,B203,
 B207,B225,B231
 smoke-free B144
 smoke-free air travel B217
 smokers A14,B24,B89,B198,B206,B207
 smokers, children of A5
 smokers, non B206
 smokers, segregation of B207
 smokes B184
 smokeslack, factory B176
 smoking B11,B22,B24,B29,B61,B74,B76,B80,B92,B107,
 B132,B189,B207,B210,B212,B217,B228,B231
 smoking areas B207
 smoking mothers A5,B101
 smoking rates, socioeconomic differences in B145
 smoking surveys B137
 smoking, cigarette A70,B22,B44,B54,B74,B87,B89,B93,
 B94,B97,B132,B135,B137,B144,B240
 smoking, human B101
 smoking, maternal B178
 smoking, parental A10,A26,B24,B65,B76,B212
 smoking, passive B10,B24,B107,B120,B132,B158,B174,
 B175,B203,B207,B217,B231
 smoking, prohibiting B207
 smoking, tobacco A5,A19,A40,A76,B100,B158,B172
 smoking, viewpoints on B132
 smoking-related cancers B231
 smoking-related illnesses B145
 smooth muscle sensitization A83,B169
 smooth muscles A74,B170
 sneezing A53,B173,B178
 social action B116
 social action, group B213
 social activity B36
 social behavior B12
 social change B133,B163
 social choices B148
 social class B135,B202
 social class homogeneity B233
 social class inequities A78,B134
 social climate B176
 social community B71
 social conditioning B221
 social conditions A64
 social consequences B182
 social damages B157
 social development B21,B222
 social disadvantages B182
 social discrimination B139
 social drinking B235
 social environment B8,B98,B107,B176
 social environment, common B176
 social environment, proximal individual B176
 social environments B229
 social environments, alternative B157
 social force B116
 social group B9
 social groups A30,B148
 social impacts B34
 social indicators B36,B97
 social institutions B163
 social interaction B87,B234
 social isolation A64,A78,B80,B134,B210
 social issues B90,B163
 social life B35
 social meaning B9
 social misfits B151
 social nature of health B83
 social network, supportive A29,A45,B81
 social networks A73,B68
 social order, inequitable B2
 social organization B7
 social ostracism B55
 social policies, canadian health and B142
 social policy B90,B218,B233
 social practice B212
 social prejudices A28
 social problem, endemic B193
 social problems B142,B179,B233
 social processes B37
 social relations B182
 social relationships A72,B87,B230
 social relationships, hospital B233
 social roles B143
 social scientists B9
 social service B86
 social services B86,B120
 social settings B163
 social skills A45
 social space B176
 social status, women's A34,B35
 social stressors B40
 social structure B2,B218
 social structures B121
 social support B52
 social support A2,B52,B68,B87,B151,B153
 social support network 7,A67,B119
 social support networks A72
 social support systems B52

- social supports B61,B138
social symbol A67,B9
social ties A31,B111
social welfare B116
social work B54
social worker, health B54
social-ecological perspective B98
socialization A30,B32,B76,B176,B235
socializing A64
socially active B204
socially interactive behaviours B234
societal attitude B231
societal biases towards homosexuality B16
societal forces B4
societal level B216
societies, children's aids B179
society, broader B212
society, democratic B163
society, healthy B148
society, polluted B133
society, post-industrial A74,B224
society, weakest members of B213
socio-economic groups B213
socio-economic status A10,B151
socio-economic status, low B24
socioeconomic B87,B137
socioeconomic causes of ill health B133
socioeconomic differences in smoking rates B145
socioeconomic disparities B137
socioeconomic factors B80,B202
socioeconomic groups B88,B137
socioeconomic status B80,B137
socioeconomic status and excess mortality B80
socioeconomic status, low B192
socioeconomic task B142
sociological response to noise B230
sodium dodecyl sulfate B115
sodium hydroxide A2,B69
soft drinks A58
soft health path B82
soil gases B199
soil pollution B23
soil, contaminated A25
soil, gases from A6
soil, wet B21
soils B200
soils, urban A20
soldiers B77
solution, technologic B135
solutions B226
solvent B140
solvent exposure B10
solvent neuropathies B157
solvents A14,B83,B137,B225,B226,B229
solvents, chlorinated A26,B165
solvents, organic A24,B9,B75
solvents, volatile hydrocarbon A82
solvents, volatile organic A13
solving, institutional problem B133
solving, problem B84
somatic and psychosomatic disorders B152
somatic complaints B101
somatic health B32
sons, teenaged A36,B62
sore throats B103
sound A60,B230
sound levels B230
soundproof B139
soundproofed A60
soundproofing B139
source control B146
source identification, pollution B128
source reduction B27
sources of energy, alternative B237
sources of stress B40,B44
sources, combustion A52
sources, energy B237
sources, indoor emission 5
sources, industrial A20
sources, intense lighting B62
sources, pollution B184
sources, unusual occupational lighting B62
southeast asians B67
soviet standards A62
space design B163
space heater, unvented B148
space heaters B91,B219
space heaters, unvented kerosene B73
space heaters, unvented natural gas B73
space heating B194
space, common B43
space, number of people in a B234
space, outdoor B84
space, social B176
space, wall B222
spaces B163
spaces, affection in public A49
spaces, closed B131
spaces, exterior B213
spaces, open B21
spaces, outdoor A20,B84
spaces, play A22,B21
spaces, private B111
spaces, public 6
span test, digit B11
span, attention A25
span, decreased attention B150
span, short attention B167
spark discharges B102
spasm, coronary. A56,B171
spastic vascular phenomena A56,B171
spasticity B103
spatial abilities A35,B84
spatial layout A75,B233
spatial relationships B69
special care B86
special education B110
special environmental requirements B239
special needs, supportive of B232
special or unique needs B35
special tax rates B149
specialist B87
specialists, learning A74
specialization A21
specialization, isolated B9
specialized housing B199
specialized personnel, highly B213
specific adaptation B53
specific therapeutic measures B142
specification, job B44
specifications B134
spectrometry, gas chromatography-mass A82
spectrometry, gas chromatography-mass B227
spectrometry, mass B117
spectroscopy, mass B243
spectrum of symptoms B180
spectrum, electromagnetic A60
spectrum, full A61,B154
speech disturbances A52
speech processing B145
speech, freedom of B138
speed of industrial life B190
spelling B150
spells of faint-dizziness B47
spill, pcb B200
spillage of exhaust A6
spills, accidental B74
spills, toxic B200
spina bifida B131
spinoffs, economic 18

-
- spirit duplicating B181
 - spirit duplicators A9,B164
 - spiritual A64
 - spiritual distress B34
 - spirometry B202
 - sponge rubber A5
 - spontaneous locomotor activity B118
 - spores, fungal B31,B103,B187
 - spores, mould A54
 - sport fishing B70
 - sports equipment A50,B179
 - spot ventilation B199,B219
 - spousal rights A47,B2
 - spray humidifiers B28
 - spray paint A26,B165
 - spray painters A13,B127
 - spray, aerosol hair A5,B70
 - sprays, aerosol A32,A74,B22,B169,B209
 - spread of bacteria B123
 - spread of innovations B122
 - spreading A55
 - spreading phenomenon B169,B170
 - sputum A5
 - stability A37,B25
 - stability, emotional A64,B54
 - stachybotrys B189
 - staff overload B216
 - stair conditions, domestic B114
 - stair design A4,B114
 - stair design, domestic B114
 - stair design, safer B114
 - stair safety B157
 - stairs B10,B114
 - stairs, accident incidence on B114
 - stairway accidents A20,B10
 - stance, advocacy B216
 - standard of living B97
 - standard of living, lower B112
 - standard, ashrae ventilation B227
 - standard, double B176
 - standard, school-safe B226
 - standard-setting B185
 - standards B6,B7,B30,B43,B125,B161,B209,B225
 - standards, air quality B140,B175
 - standards, ashrae B226
 - standards, clean air B30
 - standards, codes and B128
 - standards, comprehensive B35
 - standards, government B121
 - standards, material B25
 - standards, occupational B226
 - standards, pollutant exposure B30
 - standards, safety B92
 - standards, soviet A62
 - standing water B21
 - standing water, warm A59
 - state of health, brain's B108
 - state, emotional B1,B39
 - statements, health impact B57
 - states B151
 - states, emotional B54
 - stations, fixed monitoring B174
 - stations, subway B243
 - stations, train A19
 - statistical 8
 - Statistics Canada B194
 - statistics, accident B36
 - status B9
 - status and excess mortality, socioeconomic B80
 - status for women, equal 3
 - status of children, health B138
 - status of women B35,B134
 - status quo B26
 - status, citizenship B95
 - status, consumer B133
 - status, economic B35,B140
 - status, employment B80
 - status, health B142
 - status, increased A65
 - status, indicator of health B8
 - status, low socioeconomic B24,B192
 - status, lower health B8
 - status, marital B80
 - status, nutritional A40
 - status, socioeconomic A10,B80,B137,B151
 - status, true citizenship B212
 - status, women's economic B33
 - status, women's social A34,B35
 - statutes, civil rights B79
 - steady work B39
 - stereotype A35
 - stereotype, macho A35
 - stereotyped characterization B50
 - stereotyped images A24,A46
 - stereotyped images of gays B40
 - stereotypes 8,13,A64,B86
 - stereotypes of homosexuality B56
 - stereotypes, sexrole A26,B243
 - stereotypical images A49
 - stereotyping B26,B182
 - stereotyping of men B114
 - stereotyping, male A35,B99
 - stigma B216
 - stigma of mental illness A43,A70,B112
 - stigmatization A47
 - stigmatized house B122
 - stigmatized population B201
 - stigmatizing A42
 - stigmatizing the house B122
 - stimulation A30,A67,B12,B42
 - stimulation, environmental A64,A66,A75,B107,B210,B233,B234
 - stimulation, noxious A65,B32
 - stimulus B183
 - stimulus underload A64
 - stimulus, auditory A60
 - stimulus, fruitful B222
 - stone houses B11
 - storage B242
 - stove B171
 - stove, gas B54,B101
 - stove, gas backpacking B41
 - stoves B219
 - stoves, electric B202
 - stoves, gas A5,B20,B93,B94,B101,B121,B202
 - stoves, gas cooking B155
 - stoves, gas fired cooking B73
 - strain B33
 - strain, environmental B17
 - strain, physiological B44
 - strained personal relationships B40
 - strategies for change B35
 - strategies for prevention B144
 - strategies, behavior change B22
 - strategies, conservation B120
 - strategies, control B137
 - strategies, coping A46
 - strategies, health B8,B133
 - strategies, health promotion B204
 - strategies, lead screening B20
 - strategies, preventive B233
 - strategy B165
 - strategy, international health B239
 - Strategy, Premier's Council on Healthy B64
 - street life A46,B173
 - street, low-income B213
 - streets, urban B221
 - strength, bone A29
 - strength, muscle A61
 - streptococcal pneumonia B50

- streptococci B210
streptococcus H32
stress A32,B17,H37,B40,H44,B50,B52,B53,B54,B55,B60,
U61,H66,H69,B78,H81,H100,B103,B105,B107,B108,B121,
B126,B138,B149,B152,B153,B181,B182,B184,B190,B195,
B200,B206,B209,B210,B216
stress at work B44
stress factors B232
stress genetics B40,B61
stress levels A68
stress load 9
stress load, total B18
stress of unemployment B112
stress test B146
stress trials B100
stress, acceleration B100
stress, cardiovascular A41
stress, cardiovascular reactivity to mental B146
stress, consequences of B40
stress, economic 9,B149
stress, emotional B39
stress, examination A10,A64
stress, financial 5
stress, indices of B206
stress, job 13,B78
stress, manifestations of B190
stress, mediators of B52
stress, mental B146
stress, occupational B44
stress, physiological B150
stress, psychological A10,A64,B53,B54
stress, psychosocial B241
stress, reduction of B37
stress, sources of B40,B44
stress, work B121
stress, workplace B34
stress, workplace-related B122
stress-buffering B52
stress-inducing environments B183
stress-reducing A68
stress-reducing environments A68,B183
stress-related behaviour B108
stress-related illnesses B34
stressed population A66
stresses B157
stresses, environmental B45
stressful 5,A64
stressful events B61
stressful life events B195
stressful psychosocial situations B122
stressor B17
stressor, environmental B40
stressors B206
stressors of city life B12
stressors, ambient A65,B31
stressors, delayed responses to B40
stressors, environmental B31,B40,B66,B108,B141,B183,
B210
stressors, life B101
stressors, macroenvironmental B141
stressors, psychosocial B17
stressors, social B40
stroke B108,B112
strokes A58,B100
strong A35
structural changes A71
structural deterioration A6,B38
structural prejudices 3
structural problems A68
structure B235
structure of associations B133
structure of institutions B133
structure, family B26
structure, organizational B44
structure, reward B182
structure, social B2,B218
structures, social H121
student achievement B84
students A8,A30,B17,B24,B149,B198,B204,B209,B232
students, allergic B113
students, doctoral B53
students, elementary school A60
students, graduate A64,B232
students, high school B24,B105
students, medical and law B40
students, sensitive A11
studies of farmers B192
studies, animal B102,B133,B157,B186
studies, case B200
studies, epidemiologic B192
studies, epidemiological B65,B102
studies, field B227
studies, mortality B155
studios, arts A12
study B56
study of aging, ontario longitudinal B89
study on aging B89
study, epidemiologic B203
study, epidemiological B61
study, gas chromatographic B56
study, research B203
studying a trade B44
style, life B125
styrene A13,A14,B191,B226
styrene exposure B11,B123
sub-antarctic B232
sub-clinical effects, clinical and B111
sub-population, sensitive B121
sub-stations B132
subdivisions B243
subfloor B128
subgroup, susceptible B204
subjective mood B235
subjects, human B144
subjects, research A28,B144
subjects, sensitive B143
sublingual allergy testing B111
subordinates B44
subsample, high-risk B243
subsidized housing A36,B63
substance abuse 13,A78,B108,B134,B135,B192
substance exposure, toxic B108
substance use B173
substance, designated B125
substances, addictive B112
substances, allergenic B29
substances, bioaccumulative B180
substances, carcinogenic B12
substances, chemical B47
substances, discharge of toxic B222
substances, harmful chemical B215
substances, hazardous B152
substances, low doses of toxic B15
substances, non-toxic B47
substances, toxic B14,B53,B60,B104,B152,B180,B208,
B209,B237
substances, volatile A11
substation, high-voltage B15
substitution B134
substrate A6,B215
substrate, genetic B157
subtle behavioral alteration B64
subtle neurologic sequelae B192
suburban environments B12
suburban life B12
suburban living B233
suburban men B182
suburban neighbourhoods, racially-integrated B233
suburbanites A19
suburbs A34,B183,B233

- suburbs, feminine B183
 subway stations B243
 sucrose B39
 sudden infant death A40
 sudden infant death syndrome B5
 suffering, emotional B79
 suffering, ensuing B79
 suffocation, psychological A47,B2
 sugar, blood B187
 suicide A33,B132,B134,B189
 suicide, youth 12
 suitable activities B84
 suits, civil A38
 sulfate depositions, wet B70
 sulfate, sodium dodecyl B115
 sulfates B42
 sulfates, respirable B54
 sulfhydryl group, blood A63
 sulfide, hydrogen A8,B69
 sulfinate of michler's hydrol, paratoluene A16
 sulfinate, paratoluene B129
 sulfite B168
 sulfite oxidase deficiency B88
 sulfite-related adverse reactions B240
 sulfites A57,B88,B240
 sulfites as preservatives B168
 sulfites, dietary B88
 sulfonamide A57,B161
 sulfur compounds B91
 sulfur dioxide B18,B46,B65,B88,B93,B105,B121,B133,
 B141,B142,B154,B184,B191,B234
 sulfur dioxide, respirable suspended particles B93
 sulfur oxides B91,B211
 sulfur, acidic B187
 sulfuric acid A8,B69
 sulphydryl, blood B100
 sulphates B54
 sulphur A39,B181
 sulphur dioxide A5,A20,A22,A51,A52,B116,B196,B216
 sulphur dioxide emissions B27
 sulphur oxides A51
 sulphuric acid factory A38,B180,B208
 sunburn, tendency to B62
 sunshine A76
 sunshine, excess B46
 suntan A61
 suntan effect B84
 superfluous A27,B220
 superman image A35
 supersensitivity B99
 supervisors A17
 supervisors, relations with B206
 support, social B52
 supplies, art B47,B167
 supplies, janitorial B22
 supplies, scholastic B22
 supplies, water A6
 supply of housing B78
 supply, air B225
 supply, clean water B116
 supply, food B215
 supply, fresh air B18
 supply, water A58,A78,B181
 support B87,B95,B156
 support for citizen action B144
 support network, social 7,A67,B119
 support networks, social A72
 support services B71,B119
 support systems, social B52
 support, adult A25
 support, emotional A66
 support, framework for B39
 support, medical A74
 support, mutual A35,A37
 support, peer A73,B76
 support, psychological B112
 support, social A2,H52,H68,H87,B151,B153
 supportive of special needs B232
 supportive social network A29,A45,B81
 supportiveness B55
 supports, social B61,B138
 suppressant, dust B223
 suppressed A84
 suppressed immune systems A42
 suppression, immune A83
 suppression, immune system B58
 surface fields B102
 surfaces, indoor B202
 surfaces, slip-resistant B10
 surgical procedures A33
 surveillance A27,A78,A83,B192,B235,B239
 surveillance, health B129
 survey B131
 survey data, health B143
 Survey, Canada Fitness B137
 Survey, Canada Healthy B46,B137
 survey, environmental B131
 survey, national health B179
 survey, work environment B206
 surveys, blood lead B205
 surveys, smoking B137
 survival B58
 survival of domination B2
 susceptibilities, chemical B209
 susceptibility B21,B30,B40,B60,B97,B149,B166,B168,
 B170,B172,B196,B197,B238,B240
 susceptibility to bacterial infection A14
 susceptibility to bacterial infection, enhanced B32
 susceptibility to heart disease B76
 susceptibility to infection, streptococcus B32
 susceptibility to toxins, tissue B119
 susceptibility, chemical A6,B22,B126,B160,B161,B166,
 B188,B189
 susceptibility, disease A64,B40
 susceptibility, genetically determined B75
 susceptibility, host B104
 susceptibility, indicators of B45
 susceptibility, individual B53,B149
 susceptibility, worker disease B129
 susceptible individuals A76,B47,B48,B64
 susceptible individuals, chemically A11,B212
 susceptible individuals, environmentally B58
 susceptible population B198
 susceptible population, chemically B198
 susceptible populations B215
 susceptible subgroup B204
 susceptible teachers B166
 susceptible workers A17
 susceptible, chemically B199
 suspended microbiological material B165
 suspended particulate B29
 suspended particulate matter B105
 suspended particulates B205
 suspended particulates, respirable B93,B173
 swedish national council B123
 sweeper, manual carpet B165
 symbol, social A67,B9
 symbolic associations B183
 symptomatic housewives B93
 symptoms in children, respiratory B108
 symptoms, asthmatic B51
 symptoms, cold and flu B103
 symptoms, neuropsychiatric A77,B29
 symptoms, persistent A15
 symptoms, respiratory A8,B86,B108
 symptoms, spectrum of B180
 synaptic function, brain A62,B102
 synchronization B84
 syndrome, acquired immune deficiency A47
 syndrome, allergy tension fatigue B160

- syndrome, down's B131
 syndrome, minimal brain dysfunction B113
 syndrome, sick building A9,A17,A82,B3,B35,B73,B76,
 B140,B153,B177,B237,B238
 syndrome, sudden infant death B5
 syndrome, tension-fatigue B113
 syndrome, tight building B18,B149
 synergistic A52,B43
 synergistic effect B60,B191
 synergistic effects 7
 synergy B60
 synthesis, red blood B116
 synthetic A55
 synthetic building material B137
 synthetic chemicals, overload of B170
 synthetic clothing A32,A55
 synthetic compounds B168
 synthetic fabric B191
 synthetic fibers A53,B14,B120,B141
 synthetic fibres B211
 synthetic materials B113
 system damage, central nervous A15,A57,B215
 system depressants, central nervous B172
 system dysfunction, nervous B51
 system effects, nervous B7
 system reactions, nervous B161
 system suppression, immune B58
 system, auditory A60,B230
 system, autonomic nervous B191
 system, canadian health care B224
 system, cardiovascular B94,B171,B191
 system, central nervous A8,A76,B23,B80,B119,B125,
 B157,B164,B223,B225,B235
 system, computerized diagnostic A82,B108
 system, criminal justice B67,B147
 system, education A10,B216
 system, educational A27
 system, forced air heating B165
 system, gastrointestinal B97
 system, health care A74,B63,B86,B123
 system, health information B129
 system, hot water B165
 system, human immune B28
 system, immune A52,A56,A58,A77,B3,B17,B18,B28,B40,
 B53,B54,B94,B95,B130,B171,B187
 system, mental health B95
 system, mental health care B216
 system, nervous B8,B49,B102,B103,B116,B193,B215
 system, nonspecific clearance B188
 system, notification B6
 system, peripheral nervous B94,B235
 system, plumbing B70
 system, priority points B63
 system, recirculation A11,B212
 system, respiratory A75,B36,B228
 system, ventilating B177
 systemic toxicity B215
 systems B17
 systems, activity B182
 systems, aircleaning B210
 systems, data B138
 systems, diagnostic B108
 systems, formal service B156
 systems, health care 3
 systems, health care delivery 3,B156
 systems, health information B129
 systems, heating B147
 systems, home heating A6
 systems, hot water A54,B21
 systems, household heating A20
 systems, measurement B142
 systems, mental health care A43
 systems, organ A84
 systems, rehabilitation B156
 systems, self-help B85
 systems, social support B52
 systems, suppressed immune A42
 systems, ventilation B54,B73,B76,B207
 systems, vital organ B119
 systems, water A59
 systems, welfare B95
 systolic B146
 T-lymphocytes B171
 tan B62
 tantrums, temper A52
 taps, domestic water B21
 target groups B8,B137
 target organ B193
 task, socioeconomic B142
 tax rates, special B149
 TCDF B242
 teachers A32,B162,B209,B222,B235
 teachers, susceptible B166
 teaching hospitals B91
 techniques, analytical B136,B238
 techniques, discipline B147
 techniques, measurement B136
 techniques, prevention B113
 techniques, psychometric B223
 techniques, therapeutic A77
 techniques, therapeutic and diagnostic B53
 techniques, treatment A76
 technologic B91
 technologic solution B135
 technological change A82
 technologies A50
 technologies, emerging 3
 technologies, household B165
 technologies, new 7,A81,A82
 technologies, protection B188
 technology A2,A79,B60,B117
 technology focus B224
 technology, appropriate A87,B165
 technology, brainwave measurement A82
 technology, development of B190
 technology, high-cost B91
 technology, modern A42,B229
 technology, nuclear A66
 teenaged sons A36,B62
 teenagers B24,B212
 telecommuting A18,B37,B39
 telegraph B186
 telephone linemen B186
 telephone linemen, power and B186
 television A26,B243
 temper tantrums A52
 temperate regions B232
 temperature A16,A17,B131,B206
 temperature equalization B165
 temperature fluctuations B103
 temporal aspects of homes B230
 temporarily able-bodied A40,A41
 tenant A36
 tenant health and safety A1,A81
 tenants 19,B184
 tendency to sunburn B62
 lenders A11,B75
 tension A64,B33,B54
 tension fatigue syndrome, allergy B160
 tension, high B44
 tension, skeletal-muscle A60,B230
 tension-fatigue syndrome B113
 tennis B41
 teratogenesis B75
 teratogenic A52,B18,B221
 teratogenic effects B70,B238
 teratogens A22,B75,B168
 term health planning, long B118
 terminals, transportation A19,B210
 terminals, video display A16,A32,A62,B126,B128,B206

- territoriality A65,B163
 territories, near-home B213
 tertiary care settings B233
 test, digit span B11
 test, psychological B170
 test, stress B146
 testing laboratories B75
 testing, sublingual allergy B111
 tests, blood B171
 tests, intelligence B213
 tests, laboratory B157
 tests, mental loading B63
 tests, neurobehavioral B164
 tests, performance B11,B223
 tests, provocation B159
 tests, psychological B54
 tests, psychological function B83
 tests, respiratory function B204
 tetrachloride, carbon B27
 tetrachlorobenzenes B136
 tetrachloroethylene B149
 textile fibers A55
 textile fibres B14
 textile industry B67
 textiles A9,B151
 that promote health, behaviours B204
 the epp report B133
 the role of language B26
 the search for shelter B78
 their parents, property of B213
 theory, dominant B142
 therapeutic A51,B134
 therapeutic and diagnostic techniques B53
 therapeutic environment B65,B157
 therapeutic environments B65
 therapeutic measures, specific B142
 therapeutic medication B178
 therapeutic techniques A77
 therapy, attribution B141
 therapy, dietary A76,B50
 therapy, dietary manipulation B49,B50
 therapy, family B156
 therapy, iatrogenic drug B3
 therapy, injection B170
 therapy, orthomolecular B73
 thermal climate B123
 thermal effects in tissues B17
 thermal factors B238
 thermal insulation A54,B25,B189,B215
 thermal pollution B17
 thermal resistance B25
 thermactinomyces vulgaris B224
 thinking, egocentric A50,B209
 thinking, limited way of B26
 threatening A65,B183
 three-tier government office complex B131
 threshold A17,B227
 threshold concentrations B21
 threshold effects B168
 threshold for irritation B83
 threshold limit value, occupational B227
 threshold limit values B3,B7
 threshold, irritancy A53
 throat A74,B169,B170
 throat irritation B129
 throat, dry A12,B115
 throats, sore B103
 through consent, people acting B133
 ties, affectional A72,B87
 ties, number of close B87
 ties, social A31,B111
 tight building syndrome B18,B149
 tight buildings B62
 tight homes A6
 tightly-constructed B177
 tightly-sealed building B16
 tightly-sealed building envelope B199
 tightly-constructed A74
 tightness, chest A9
 tiles, vinyl B178
 time flexibility, work B45
 time schedule B182
 time, allocation of B44
 time, commuting A18,B37
 time, reaction A15,B7,B235
 time-lapse cameras A10
 time-lapse photography B155
 tissue resistance B119
 tissue susceptibility to toxins B119
 tissue, adipose B188
 tissue, brain B232
 tissues B69
 tissues, body A82
 tissues, brain B102
 tissues, infection of B215
 tissues, thermal effects in B17
 tobacco B46,B112,B116,B152,B157,B231
 tobacco smoke A8,A14,A15,A16,A53,B29,B42,B43,B55,B62,
 B74,B86,B125,B137,B173,B174,B175,B176,B199,B203,B207,
 B225,B231
 tobacco smoke, second-hand B72
 tobacco smoking A5,A19,A40,A76,B100,B158,B172
 tobacco use B144
 toiletries B22
 token power A79
 tolerance A68,A79,B2,B37
 tolerance for pollution A84
 tolerance of differences A17
 tolerance of differentness B34
 tolerances, toxic B36
 toll, emotional A42,A69,B147
 toluene A9,A14,A15,B149,B172,B226
 toluene diisocyanate A14
 toluene diisocyanate B38,B151
 tomatoes A42
 toner A16
 tonics A32
 tools, basic design B1
 Toronto B104,B119,B125,B126
 tort actions B79
 total building performance A10,B75
 total load B17,B29
 total stress load B18
 total volatile organic compounds B227
 tourism B70
 towards aging, attitudes B90
 towards homosexuality, societal biases B16
 towers, cooling A54,B21,B236
 town A21
 town environments, small B68
 town settings, small B148
 towns B72
 towns, small B148
 toxic chemical exposure B109
 toxic chemicals B45,B47,B94,B177
 toxic compounds B168
 toxic encephalopathy B9
 toxic spills B200
 toxic substance exposure B108
 toxic substances B14,B53,B60,B104,B152,B180,B208,
 B209,B237
 toxic substances, discharge of B222
 toxic substances, low doses of B15
 toxic tolerances B36
 toxic waste dump A39,B181
 toxicants, energy-related B238
 toxicity A56,B16,B36,B47,B105,B113,B118,B187,B192,
 B228,B235
 toxicity of environmental pollutants B126,B192
 toxicity of plutonium B61

- toxicity, chemical R180
toxicity, environmental B94
toxicity, neurobehavioral B235
toxicity, neurobehavioural A9,A15,B172
toxicity, neurobehavioural B164
toxicity, pesticide R193
toxicity, systemic D215
toxicological principles B6
toxicology B43,B199,B220,B221
toxicology, behavioral B7,B229
toxicology, behaviour B73
toxicology, behavioural A76
toxicology, environmental B60
toxicology, gas B5
toxigenic fungi A77,B187
toxin B37
toxins B47,B157
toxins can A57
toxins from molds B52
toxins, environmental B103,B214
toxins, fusarium A57,B189
toxins, macrocyclic trichothecene A54,B103
toxins, penicillium B189
toxins, tissue susceptibility to B119
trace elements B200
trace metals A51
tracer gas R155
tracheal mucociliary transport B70
track-etch detector B89
tract defense mechanisms, respiratory B187
tract diseases, respiratory B202
tract infections, respiratory B212
tract, gastrointestinal A60,B230
tract, optic B186
tract, respiratory A52,B5,B14,B16,B160,B187
tract, upper respiratory B131
trade B135
trade names B33
trade, studying a B44
trades A13
trades, painting B190
traditional lifestyles B46
traditional window designs B222
traffic A22,B21,B183
traffic accident B4
traffic congestion A21,B109
traffic dust, automobile A22,B217
traffic safety A27
traffic tunnels A20,B221
traffic, children's performance in B202
train stations A19
trained B211
training, medical A78,B237
training, nutritional A26
training, professional B40
trains B210
traits, personality A65,A72,B32,B55
transfusions, blood B100
transfers B93
transit, mass R229
transitional shelter B78
transitions B54
transmission B28
transmission facilities, electricity B132
transmission lines B102,B132
transmission lines, high-voltage A62
transmission, electrical B153
transmission, electrical power A61,B28
transmission, person-to-person B214
transmitted disease, sexually B131
transmitted diseases, sexually 13,B2,B173
transport of airborne pollutants, longrange B70
transport, oxygen B49
transport, tracheal mucociliary B70
transportation A51,A67,B8,B119,B243
transportation terminals A19,B210
trauma B51,B157
traumatic experience B125
travel, smoke-free air B217
treated fabrics A5
treatment R124,B180,B209
treatment facilities, sewage B21
treatment methods A75
treatment of water B21
treatment settings, residential B234
treatment techniques A76
treatment, detoxification B188
treatment, general hospital B216
treatment, ineffective B7
treatment, prejudicial A32
treatments, durable press B14
treatments, durable-press A55
trees, planting B9
tremor B87
trends B135
trends, obsessive-compulsive B54
trends, obsessive-compulsive A64
trials, clinical B22
trials, stress B100
triangle, junction B61,B203
tribe B58
trichlorethane A9,A15
trichloroethane B27,B172
trichloroethylene A15,B63,B149
trichothecene mycotoxins B187
trichothecene toxins, macrocyclic A54,B103
trichothecene-producing fungi, macrocyclic B103
trichothecium B189
tricresyl phosphate B195
trigeminal nerve endings B11
trigeminal nerves A54,B62
triggering agents, food B171
trinitrofluorenone A15
tris B116
trouble, heart A69,B47
troubles, attentional A60,B139
true citizenship status B212
tuberculosis, pulmonary B178
tubes, fluorescent A9
tubes, full-spectrum fluorescent A10
tuition rebates B149
tumors B193
tumours B28
tunnels, traffic A20,B221
turbulence, familial B45
tutorial B53
twentieth century disease B216
two-parent families A36
two-parent households B39
type a behavior B22
type a coronary-prone behavior B50
type, family 3,A2,A36
type, household A37
typical office environment B127
typing paper B69
typology, fallacy of B89
ubiquitous 13
UFFI A6,B31,B51,B52
UFFI-fungal interaction B51
ulcers A58,B112,B152
ulterior motives B190
ultra-trace levels A82
ultraviolet B62
ultraviolet light A16,A61,B62,B82
ultraviolet light, full-spectrum B154
ultraviolet radiation B30,B178
ultraviolet, long-wavelength A9,B154
umbrella of oppression B20
unemployment picture, chronic B33
uncertainty A66,B181

- uncertainty, personal B80
 uncooperativeness B150
 under-nutrition A70
 under-nutrition, chronic B218
 under-representation 13
 underdevelopment, moral B185
 undergraduate curricula B204
 underground environment B49
 underground, safety B49
 underlay A6
 underlay, particleboard floor B106
 underload, stimulus A64
 underprivileged B196
 understanding B144, B184
 understanding, scientific B163
 undervalued 16
 undervaluing 6
 undesirable legislation, repeal of B139
 undiffused light B62
 unemployed 17, A71, B151, B152
 unemployed individuals B47
 unemployed individuals, older B47
 unemployed, blue-collar A69, B47
 unemployed, low-income A69, B47
 unemployment 3, 5, 13, A64, A69, A78, A79, B33, B46, B88, B112, B134, B140, B151, B152, B196
 unemployment and health B46
 unemployment and physical health B112
 unemployment benefits B151
 unemployment rates, local B140
 unemployment, effects of B112
 unemployment, human costs of B112
 unemployment, periods of B112
 unemployment, stress of B112
 unemployment-related B151
 unexamined premises B50
 unexpected hazards A44
 unfamiliar environment B65
 unfeeling A35, B99
 unhealthful behavior modeled by peers B76
 unhealthy side effects B220
 unhealthy working conditions A38
 unimportant A27
 union of Ontario Indians B180
 unionization B33
 unions B34, B69, B152
 unique needs, special or B35
 united nations B222, B236, B238
 units, environmental care B190
 units, fan coil A54
 units, geriatric assessment B56
 units, living B43
 universal employment B148
 universal health insurance B8
 universal leave programs B149
 universities B91
 university conditioning classes B76
 university entrance B46
 unlearning B26
 unpleasant work conditions B44
 unpredictability A60
 unpredictable noise B231
 unprepared communities B156
 unrealistic deadlines B209
 unrealistic expectations B195
 unrecognized allergy B167
 unresolved feelings B195
 unskilled B34
 unskilled adults, young B151
 unstable amplifier B200
 unstable jobs B196
 unusual occupational lighting sources B62
 unusually resistant B45
 unvented combustion appliances A5
 unvented hood B165
 unvented kerosene space heaters B73
 unvented natural gas space heaters B73
 unvented space heater B148
 unwanted pregnancy 13
 upheaval and A79
 upper airway irritation B5
 upper airway obstruction A16
 upper airways B6
 upper limit, guaranteed B138
 upper middle-class whites B9
 upper respiratory irritation B30
 upper respiratory tract B131
 uprooting A65, B122
 urban B183
 urban areas A21, B200, B221
 urban built environment B107
 urban centres B86, B191
 urban density B12
 urban design B230
 urban development B162
 urban dwellers A19, A66, B184
 urban ecosystem B230
 urban environment B12, B36, B58, B109, B119, B230
 urban environments B12, B107, B229
 urban environments, habitable B230
 urban life A66, A72, B68, B163, B183
 urban life, quality of B9, B163
 urban men B182
 urban neighborhoods, dense A67, B12
 urban noise B12
 urban physical environments B164
 urban planning A21, B8, B162
 urban residents B203, B220
 urban soils A20
 urban streets B221
 urban-rural difference B11
 urbanites, crowded A67, B12
 urbanization B12, B239
 urbanization and human health B12
 urea B70
 urea formaldehyde A6, B25, B106
 Ureco Formaldehyde Foam Insulation A6, B14, B20, B31, B38, B51, B79, B197, B209
 urea formaldehyde foams B19
 urea formaldehyde glue A53, B5
 urea formaldehyde in fabrics B83
 urticaria A16, A57, B240
 urticarial reactions A13, B14
 use of the dwelling B113
 user B37
 user complaints, building B3
 user experience B95
 user participation 7
 users, pesticide B192
 utilities A20, A51
 vaccination B177
 vaccine B144
 vaccine, measles B177
 vaccines B86
 vacuum cleaner B165
 value, occupational threshold limit B227
 value, people of A29
 values, adult A66, B74
 values, changing B120
 values, communicating B111
 values, threshold limit B3, B7
 vandalism A46, B67
 vapor, acrolein B32
 vapor, mercury B69
 vapour, mercury B177
 vapour, water A5, B199
 vapours B11
 vapours, gasoline B124
 vapours, organic A53
 variability B30

- variability among people A24
variability of response A84
variability, radon B181
variables, community mental health B140
variables, environmental B98,B140
variables, health B112
variables, psychological A65
varnishers B10
vascular B170
vascular dysfunction B171
vascular headaches B171
vascular phenomena, spastic A56,B171
vasculitides, auto-immune B171
vasculitis B171
vasculitis, collagen B171
vdl radiation hazards B208
vols B90
vegetables A57
vegetation B108
vegetation, devoid of B208
vehicles B101,B210
vehicles, motor A20,B218
vehicles, road A50,B179
ventilated, mechanically B206
ventilating experts A74
ventilating system B177
ventilation A6,A8,A9,A14,A16,A32,B11,B18,B19,B20,B21,
B22,B29,B48,B66,B74,B78,B79,B88,B89,B91,B92,B123,
B124,B131,B134,B146,B149,B152,B172,B173,B174,B177,
B181,B184,B194,B199,B217,B219,B220,B225,B227,B236,
B237,B242
ventilation costs R49
ventilation performance B207
ventilation rate B73,B158,B174
ventilation rates B3,B210,B227
ventilation standard, ashrae B227
ventilation systems B54,B73,B76,B207
ventilation, artificial B177
ventilation, bathroom B165
ventilation, building B211
ventilation, creative A7
ventilation, effective A17
ventilation, inadequate B90
ventilation, mechanical B210
ventilation, poor B63,B206,B208
ventilation, reduced B19,B172
ventilation, residence reduced B19
ventilation, spot B199,B219
ventilation, whole-house B219
ventilatory function, impaired B87
ventilatory performance B194
venting, direct B207
ventricular contractions, premature A55,B191
verbal intimidation A46,B67
vertebrobasilar insufficiency B38
vertigo A75,B169
vessels, empty B162
vestibular dysfunction B9
veterans, vietnam B193
viable mould B82
viable organisms B120
vibration A63,B100,B123
vibration, sinusoidal B100
victimization B25
victims of rape B13
victims of welfare, child B45
victims, abuse A28
victims, protected B147
video display terminals A16,A32,A62,B126,B128,B206
vietnam veterans B193
view, child-centered B25
view, holistic B82
viewpoints on smoking B132
views, minority B139
vinyl chloride A15,B27,B116,B243
vinyl seating B195
vinyl tiles B178
vinylidene chloride resins B67
violations of rights 11
violence A33,A34,A35,A66,B13,B67,B71,B127,B134,B135,
B178
violence, domestic B147
violence, family B101
violence, hate-motivated A46,B67
violence, physical B220
violent B99
violent crime B178
viral A54
viral infections B104
viral pathogen B32
virus infections, respiratory B178
virus, epstein-barr A24,B75
viruses A9,A29,A76,B46,B104
viruses in air B104
viruses, indoor airborne B238
visibility B58
vision disorders B38
vision for community B133
vision, blurred A9
vision, future-oriented B50
vision, political 19
visions, collective B183
visits, physician B47
visual acuity B7
visual changes B56
visual disturbances A9,B87,B106
visual evoked potential B24
visual memory A63,B110
visual pollution B17
vital capacity B202
vital capacity, forced B6
vital organ systems B119
vitamin d B30
vitamins A56
vlf fields B128
voc B140
vocs B227
volatile hydrocarbon solvents A82
volatile inhalants B172
volatile organic chemicals 15,B5
volatile organic compounds A54,B57,B140,B142,B243
volatile organic compounds, total B227
volatile organic contaminants A77
volatile organic solvents A13
volatile organics B227
volatile pollutants B16
volatile substances A11
volatiles B76
volatiles, fungal B215
voltage B102
voltage line, high B153
volume measurements, lung B204
volume, forced expiratory B6,B110,B202
voluntarism A34,B219
voluntary euthanasia B189
volunteer programs B216
volunteers A30,B204
vomiting A6,B128,B225
vulgaris, thermoactinomyces B224
vulnerabilities B109
vulnerabilities to illness B149
vulnerability J,A40,A48,B47,B54,B131
vulnerability to hazards A81
vulnerability to pollutants, range of B199
vulnerability, range of 5,19,A17
vulnerable children A26,B243
vulnerable population groups B239
vulnerability, selective B157
walking A9
wall panelling B128

-
- uncertainty, personal B80
 - uncooperativeness B150
 - under-nutrition A70
 - under-nutrition, chronic B218
 - under-representation 13
 - underdevelopment, moral B185
 - undergraduate curricula B204
 - underground environment B49
 - underground, safety B49
 - underlay A6
 - underlay, particleboard floor B106
 - underload, stimulus A64
 - underprivileged B196
 - understanding B144,B184
 - understanding, scientific B163
 - undervalued 16
 - undervaluing 6
 - undesirable legislation, repeal of B139
 - undiffused light B62
 - unemployed 17,A71,B151,B152
 - unemployed individuals B47
 - unemployed individuals, older B47
 - unemployed, blue-collar A69,B47
 - unemployed, low-income A69,B47
 - unemployment 3,5,13,A64,A69,A78,A79,B33,B46,B88,B112,
B134,B140,B151,B152,B196
 - unemployment and health B46
 - unemployment and physical health B112
 - unemployment benefits B151
 - unemployment rates, local B140
 - unemployment, effects of B112
 - unemployment, human costs of B112
 - unemployment, periods of B112
 - unemployment, stress of B112
 - unemployment-related B151
 - unexamined premises B50
 - unexpected hazards A44
 - unfamiliar environment B65
 - unfeeling A35,B99
 - unhealthful behavior modeled by peers B76
 - unhealthy side effects B220
 - unhealthy working conditions A38
 - unimportant A27
 - union of Ontario Indians B180
 - unionization B33
 - unions B34,B69,B152
 - unique needs, special or B35
 - United Nations B222,B236,B238
 - units, environmental care B190
 - units, fan coil A54
 - units, geriatric assessment B56
 - units, living B43
 - universal employment B148
 - universal health insurance B8
 - universal leave programs B149
 - universities B91
 - university conditioning classes B76
 - university entrance B46
 - unlearning B26
 - unpleasant work conditions B44
 - unpredictability A60
 - unpredictable noise B231
 - unprepared communities B156
 - unrealistic deadlines B209
 - unrealistic expectations B195
 - unrecognized allergy B167
 - unresolved feelings B195
 - unskilled B34
 - unskilled adults, young B151
 - unstable amplifier B200
 - unstable jobs B196
 - unusual occupational lighting sources B62
 - unusually resistant B45
 - unvented combustion appliances A5
 - unvented hood B165
 - unvented kerosene space heaters B73
 - unvented natural gas space heaters B73
 - unvented space heater B148
 - unwanted pregnancy 13
 - upheaval and A79
 - upper airway irritation B5
 - upper airway obstruction A16
 - upper airways B6
 - upper limit, guaranteed B138
 - upper middle-class whites B9
 - upper respiratory irritation B30
 - upper respiratory tract B131
 - uprooting A65,B122
 - urban B183
 - urban areas A21,B200,B221
 - urban built environment B107
 - urban centres B86,B191
 - urban density B12
 - urban design B230
 - urban development B162
 - urban dwellers A19,A66,B184
 - urban ecosystem B230
 - urban environment B12,B36,B58,B109,B119,B230
 - urban environments B12,B107,B229
 - urban environments, habitable B230
 - urban life A66,A72,B68,B163,B183
 - urban life, quality of B9,B163
 - urban men B182
 - urban neighborhoods, dense A67,B12
 - urban noise B12
 - urban physical environments B164
 - urban planning A21,B8,B162
 - urban residents B203,B220
 - urban soils A20
 - urban streets B221
 - urban-rural difference B11
 - urbanites, crowded A67,B12
 - urbanization B12,B239
 - urbanization and human health B12
 - urea B70
 - urea formaldehyde A6,B25,B106
 - Urea Formaldehyde Foam Insulation A6,B14,B20,B31,B38,
B51,B79,B197,B209
 - urea formaldehyde foams B19
 - urea formaldehyde glue A53,B5
 - urea formaldehyde in fabrics B83
 - urticaria A16,A57,B240
 - urticarial reactions A13,B14
 - use of the dwelling B113
 - user B37
 - user complaints, building B3
 - user experience B95
 - user participation 7
 - users, pesticide B192
 - utilities A20,A51
 - vaccination B177
 - vaccine B144
 - vaccine, measles B177
 - vaccines B86
 - vacuum cleaner B165
 - value, occupational threshold limit B227
 - value, people of A29
 - values, adult A66,B74
 - values, changing B120
 - values, communicating B111
 - values, threshold limit B3,B7
 - vandalism A46,B67
 - vapor, acrolein B32
 - vapor, mercury B69
 - vapour, mercury B177
 - vapour, water A5,B199
 - vapours B11
 - vapours, gasoline B124
 - vapours, organic A33
 - variability B30

- variability among people A24
variability of response A84
variability, radon B181
variables, community mental health B140
variables, environmental B98,B140
variables, health B112
variables, psychological A65
varnishers B10
vascular B170
vascular dysfunction B171
vascular headaches B171
vascular phenomena, spastic A56,B171
vasculitides, auto-immune B171
vasculitis B171
vasculitis, collagen B171
vdt radiation hazards B208
vdfs B90
vegetables A57
vegetation B108
vegetation, devoid of B208
vehicles B101,B210
vehicles, motor A20,B218
vehicles, road A50,B179
ventilated, mechanically B206
ventilating experts A74
ventilating system B177
ventilation A6,A8,A9,A14,A16,A32,B11,B18,B19,B20,B21,
B22,B29,B48,B66,B74,B78,B79,B88,B89,B91,B92,B123,
B124,B131,B134,B146,B149,B152,B172,B173,B174,B177,
B181,B184,B194,B199,B217,B219,B220,B225,B227,B236,
B237,B242
ventilation costs B49
ventilation performance B207
ventilation rate B73,B158,B174
ventilation rates B3,B210,B227
ventilation standard, ashrae B227
ventilation systems B54,B73,B76,B207
ventilation, artificial B177
ventilation, bathroom B165
ventilation, building B211
ventilation, creative A7
ventilation, effective A17
ventilation, inadequate B90
ventilation, mechanical B210
ventilation, poor B63,B206,B208
ventilation, reduced B19,B172
ventilation, residence reduced B19
ventilation, spot B199,B219
ventilation, whole-house B219
ventilatory function, impaired B87
ventilatory performance B194
venling, direct B207
ventricular contractions, premature A55,B191
verbal intimidation A46,B67
vertebrobasilar insufficiency B38
vertigo A75,B169
vessels, empty B162
vestibular dysfunction B9
veterans, vietnam B193
viable mould B82
viable organisms B120
vibration A63,B100,B123
vibration, sinusoidal B100
victimization B25
victims of rape B13
victims of welfare, child B45
victims, abuse A28
victims, protected B147
video display terminals A16,A32,A82,B126,B128,B206
vietnam veterans B193
view, child-centered B25
view, holistic B82
viewpoints on smoking B132
views, minority B139
vinyl chloride A15,B27,B116,B243
vinyl seating B195
vinyl tiles B178
vinylidene chloride resins B67
violations of rights 11
violence A33,A34,A35,A66,B13,B67,B71,B127,B134,B135,
B178
violence, domestic B147
violence, family B101
violence, hate-motivated A46,B67
violence, physical B220
violent B99
violent crime B178
viral A54
viral infections B104
viral pathogen B32
virus infections, respiratory B178
virus, epstein-barr A24,B75
viruses A9,A29,A76,B46,B104
viruses in air B104
viruses, indoor airborne B238
visibility B58
vision disorders B38
vision for community B133
vision, blurred A9
vision, future-oriented B50
vision, political 19
visions, collective B183
visits, physician B47
visual acuity B7
visual changes B56
visual disturbances A9,B87,B106
visual evoked potential B24
visual memory A63,B110
visual pollution B17
vital capacity B202
vital capacity, forced B6
vital organ systems B119
vitamin d B30
vitamins A56
vlf fields B128
voc B140
vocs B227
volatile hydrocarbon solvents A82
volatile inhalants B172
volatile organic chemicals 15,B5
volatile organic compounds A54,B57,B140,B142,B243
volatile organic compounds, total B227
volatile organic contaminants A77
volatile organic solvents A13
volatile organics B227
volatile pollutants B16
volatile substances A11
volatiles B76
volatiles, fungal B215
voltage B102
voltage line, high B153
volume measurements, lung B204
volume, forced expiratory B6,B110,B202
voluntarism A34,B219
voluntary euthanasia B189
volunteer programs B216
volunteers A30,B204
vomiting A6,B128,B225
vulgaris, thermoactinomyces B224
vulnerabilities B109
vulnerabilities to illness B149
vulnerability 3,A40,A48,B47,B54,B131
vulnerability to hazards A81
vulnerability to pollutants, range of B199
vulnerability, range of 5,19,A17
vulnerable children A26,B243
vulnerable population groups B239
vulnerability, selective B157
walking A9
wall panelling B128

-
- wall space B222
 - war, fear of atomic B190
 - war, global A35
 - war, nuclear A66,B74
 - warfare B74
 - warm standing water A59
 - warning sign, early A74
 - wash, pine-scented floor B169
 - waste dump, toxic A39,B181
 - waste incineration B126
 - waste incineration emissions A21
 - waste management A67,B119
 - waste management, hazardous B223
 - waste oil B223
 - waste, municipal A52,B241
 - waste, radioactive B61
 - wastes, chemical B23
 - wastes, liquid B60
 - wastes, radioactive A67,B61
 - water A56,B90
 - water contamination, drinking B200
 - water damage A9,A54,B76
 - water pipes A58,B189
 - water pollution B12,B116,B189
 - water quality B70
 - water supplies A6
 - water supply A58,A78,B181
 - water supply, clean B116
 - water system, hot B165
 - water systems A59
 - water systems, hot A54,B21
 - water taps, domestic B21
 - water vapour A5,B199
 - water, acidified lake A22,A58,B70
 - water, clean B229
 - water, drinking 5,A58,B27,B115,B201,B236
 - water, ground B19
 - water, standing B21
 - water, treatment of B21
 - water, warm standing A59
 - water, well B88
 - waterborne radon-222 B88
 - Waterloo County Board of Education B216
 - waters, recreational B21
 - way of thinking, limited B26
 - weak cancer promoters B132
 - weakest members of society B213
 - weakness B50
 - weapon, nuclear B61
 - weapons, nuclear A22,A67,B60,B222
 - weather conditions A6
 - weather, bad B21
 - weight B133
 - weight gain A77,B29
 - welding A13,B123
 - welfare A36,A43,A70,B25,B45,B201
 - welfare act, child B179
 - welfare assistance B63
 - welfare of the children B144
 - welfare systems B95
 - welfare, child victims of B45
 - welfare, social B116
 - well water B88
 - well-being B25,B34,B35,B68
 - well-being, emotional B47
 - well-being, mental B240
 - well-being, people's B222
 - well-being, psychological B176
 - well-conducted classroom B222
 - well-connected A72
 - well-educated B233
 - wellbeing B106
 - wellbeing of families B149
 - wellbeing, basic B149
 - wellness B119
 - wellness behaviors B22
 - wet environment, hot and B100
 - wet soil B21
 - wet sulfate depositions B70
 - wet-process machines A16
 - wet-process photocopiers A15,B137
 - wetting agents A55,B14
 - wheel of life B25
 - wheelchair accidents B59
 - wheelchairs A42,B211
 - wheeze A5,B55
 - wheezing A53,B80
 - which B225
 - white B26
 - white blood cell B53
 - white middle class B120
 - white neighborhoods B233
 - white-collar workers B206
 - whites, upper middle-class B9
 - whole-house ventilation B219
 - widow, legless A41
 - widowed A33,B122
 - wife battering B134
 - will, living B189
 - window designs, traditional B222
 - window, access to a B240
 - windowless classrooms B154,B222
 - windowless environments B85
 - windowless rooms A61
 - windows B84,B85,B158,B222
 - windows and lighting in offices B239
 - windows of frequency response B200
 - windows, open B172
 - windsor B161
 - wine manufacturers B240
 - wines A58
 - with a disability, individuals B95
 - with individuals, direct contact 12
 - with mental disabilities, people B39
 - with others, relationships B114
 - with partners, difficulties B49
 - with physical disabilities, coping B148
 - with supervisors, relations B206
 - withholding of affection A26
 - withholding of affection B14
 - without schooling, growing B92
 - wives A36,B45
 - woman B26
 - woman, battered B134
 - woman, professional B39
 - womb A24,B59
 - women 6,A32,A69,A80,B33,B35,B37,B38,B40,B50,B62,B69,B72,B75,B93,B99,B122,B133,B134,B136,B143,B182,B183,B208,B211,B218,B228
 - women office workers B78
 - women workers B209
 - women's economic status B33
 - women's health B35
 - women's health issues B122
 - women's liberation movement B13
 - women's mental health A34
 - women's rights B13
 - women's social status A34,B35
 - women's work in the home B33
 - women, battered A33,A34,B101
 - women, battering of 12
 - women, clerical B39
 - women, devaluing of A32
 - women, discrimination against B26,B97
 - women, economic role of B218
 - women, elderly B87
 - women, equal status for 3
 - women, equality between men and B13
 - women, equality for 19
 - women, functioning B211

- women, health-related problems of B239
women, illness among B143
women, information about B35
women, needs of B196
women, older A33,B122
women, poverty among 12
women, pregnancy problems in B206
women, pregnant A40,B5,B198
women, rights of A35
women, shelters for battered B35
women, status of B35,B134
women, young B145
wood combustion B141
wood products B14
wood shavings A33,B5
wood-chip panels B159
woodchip board, prefabricated B160
woodstoves A20
woodworking shops A12,B13
wool carpets B46
work B34,B45,B87,B95,B148,B197,B218,B231
work alternative B37
work arrangements A68
work conditions, unpleasant B44
work efficiency B119
work environment B66,B83,B121,B176
work environment survey B206
work environments B39,B170
work environments, office B207
work experience B26
work in the home, women's B33
work incentives B112
work loads, household B45
work loads, imbalance in B45
work opportunities B95
work options B93,B151
work outside the home B183
work overload A68,B44
work patterns B196
work performance B17
work preparation programs B216
work pressure A65,B66
work pressures 5
work productivity B240
work schedules B151
work stress B121
work time flexibility B45
work, computer-mediated home-based B39
work, conditions of B33
work, dignifying paid B219
work, distributions of A71,B149
work, domestic A34,B183
work, flexible parttime B93
work, home-based B37
work, inability to B58
work, monotonous A32,B209
work, paid A34
work, part-time A18,A36,B45
work, places of B76
work, productive B183
work, role at B44
work, scarcity of B151
work, social B34
work, steady B39
work, stress at B44
work, world of B45
work-related death B241
work-related diseases A75
work-relatedness of disease B241
work-time, flexible A36
work-associated pressures B34
worker disease susceptibility B129
worker exposure B185
worker participation A15,A68,B185
worker's control B122
worker, health social B54
workers A61,B10,B28,B113,B123,B138,B173,B191,B217,
B240,B241
workers compensation legislation B185
workers, blue collar 6
workers, chemical B114
workers, child care A32,B209
workers, clerical A32,B31,B69,B208
workers, construction A14,B134
workers, distribution B28
workers, health care B209
workers, home-based A18,B39
workers, industrial A13,B15,B93
workers, male B45
workers, manual B28
workers, native community health care B96
workers, nonsusceptible B240
workers, office B77,B148,B149,B207,B239,B240
workers, older B112
workers, railroad B223
workers, railway B15
workers, service B208
workers, susceptible A17
workers, white-collar B206
workers, women B209
workers, women office B78
workforce, paid B208
working alone B28
working class B114
working conditions B44,B206
working conditions, unhealthy A38
working environment B37
working environment, healthy B13
working environments B222
working-class men, oppression of B114
worklife choices A18,B87,B93
workload, heavy B209
workload, imbalanced A36
workplace 3,A13,B34,B78,B93,B125,B132,B151,B162,B174,
B175,B176,B185,B192,B206,B235,B241
workplace democratization A15
workplace exposure B5
workplace exposure, parent's B165
workplace exposures B177
workplace hazards B236
workplace health and safety A15,B239
workplace stress B34
workplace, accidents in the B192
workplace, mental fitness in the B93
workplace-related stress B122
workplaces 5
workshops B226
workshops, art A14
worktime B151,B196,B219
worktime, organization of B196
worktime, re-organizing B196
worktime, rearrangement of B196
world economic crisis B138
world experts A78,A85
world health order, new B117
world leaders 18
world markets 3,19
world of work B45
worldwide market 18
wound and injuries B164
x-ray emitters B20
x-rays A62,B136
xenobiotic chemicals B72
xenobiotics, mobilization and excretion of B188
xylene A14,B226
yeasts A77,B29
York Region B113
young B25,B49,B212
young adults B147
young children B64,B199,B205,B212,B238

young drivers B130
young infants B198
young people 12,A24,A27,B152
young people's liberation B220
young people, oppression of B220
young unskilled adults B151
young women B145
young, needs of the B213
younger adults B67
youth B130,B144
youth suicide 12
youth, false ideas about A27
youths, homosexual B173
zinc B189
zinc protoporphyrin B223

